

30. Which of the following patient will **NOT** require the use of a fluid intake and output chart?
- A. Patient with a kidney disorders
 - B. Post surgical patients
 - C. Patients with normal fluid intake
 - D. Patients on IV therapy
31. What is the primary purpose of monitoring fluid intake and output (I&O)?
- A. To maintain dehydration in an acutely ill patient
 - B. To assess and maintain the patient's fluid balance.
 - C. Monitor blood sugar and BMI levels.
 - D. Track the patient's weight and height
32. Which of the following is considered fluid intake for I&O charting?
- A. Urine output
 - B. Vomit.
 - C. IV fluids.
 - D. Wound drainage.
33. What should a nurse do if a patient's fluid intake significantly exceeds their output?
- A. Continue to monitor the patient and document the findings.
 - B. Immediately increase the patient's fluid intake.
 - C. Notify the healthcare provider immediately.
 - D. Decrease the patient's fluid intake.
34. Which of the following is an example of an insensible fluid loss?
- A. Urine output.
 - B. Vomit.
 - C. Sweat.
 - D. Wound drainage.
35. When calculating fluid balance, what is the formula?
- A. Intake + output.
 - B. Intake - output.
 - C. Output - intake.

D. Intake x output.

36. The unconscious patient may be at risk of developing any of the following complications except....

- A. Airway obstruction
- B. Aspiration
- C. Pressure ulcers
- D. Fluid balance

37. What position should be used in nursing an unconscious patient?

- A. Prone
- B. Recovery
- C. lithotomy
- D. Dorsal

38. Standard precaution is crucial during sputum specimen collection

- A. TRUE
- B. FALSE

39. Which of the following is fundamental requirement before performing last offices on a patient?

- A. Organise your requirement
- B. Discuss with family members any preference they may have
- C. Ensure death is verified
- D. Ensure privacy

40. Which of the following is key in order to prevent pressure ulcers in an unconscious patient?

- A. Turning patient
- B. Feeding
- C. Bed making
- D. Checking vital signs

41. During serving of meals proper positioning is key in order to prevent....
- A. pressure ulcers
 - B. Throat cancer
 - C. Choking
 - D. Throat pain
42. The following are key requirements of meal service in nursing care **except**.....
- A. dietary Needs and Restrictions
 - B. Meal Preparation and Presentation
 - C. Positive meal time environment
 - D. Preference of the dietary team
43. Nasogastric (NG) tube feeding may be an option to use in patients who are ..
- A. Difficulty swallowing (dysphagia) due to stroke
 - B. Increased appetite
 - C. Conditions that affect the digestive system, making it difficult to absorb nutrients.
 - D. When a person is in a coma or otherwise
44. The act of serving meals within a nursing context transcends the simple delivery of food.
It is a fundamental component
- A. True
 - B. false
45. The following are contraindications for oral medication administration **except**.....
- A. NPO orders
 - B. Risk of aspiration
 - C. Conscious patient
 - D. Difficulty swallowing
46. Assessment of the an conscious patient's skin requires that the nurse
- A. Keep the patient environment clean
 - B. Lifts the patient regularly
 - C. Stays auscultates the patient's skin

D. Regular inspection of the skin particularly bony prominences

47. State the **five moments** of hand hygiene as recommended by the World Health Organization.....

.....
.....
.....
.....
.....

.....(5 marks)

48. Mention two (2) skills nurses use when utilizing the nursing process to solve patient's problems

.....(2marks)

49. State the 10 rights of medication administration

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

..... (5marks)

i. Disinfection and Sterilization

EXAMINER: STELLA SAGEO