



METHODIST UNIVERSITY GHANA

DEPARTMENT OF NURSING AND MIDWIFERY

BSC. NURSING: END OF FIRST SEMESTER EXAMINATION: 2024/2025

**COURSE CODE/TITLE: MNUR 106 FIRST AID, TRAUMA AND EMERGENCY
NURSING (3 CREDIT)**

WEEKDAY AND WEEKEND

INSTRUCTIONS:

**ANSWER ALL QUESTIONS ON THE QUESTION PAPER BY CIRCIRLING THE
BEST ANSWER FROM A-D. EACH QUESTION CARRY ONE MARK**

TIME ALLOWED: 1 HOUR

EXAMINER: STELLA SAGOE

Page 1 of 15

1. The most appropriate first aid management of epistaxis is:
 - A. Pinch the bridge of the nose and tilt the head backward.
 - B. Tilt the head forward and breathe through the mouth.
 - C. Tilt the head backwards and pour cold water on the head.
 - D. Tilt the head backward and breathe through the nose
2. What would you suspect if, soon after applying a dressing to the arm, the fingers turn pale or bluish in colour, are cold to touch and the casualty says he cannot feel his fingers?
 - A. Blood circulation is impaired
 - B. There may be a fracture underneath
 - C. There is a foreign object in the wound
 - D. The casualty is showing signs of shock
3. You met a man by the roadside staring blankly, unresponsive, making twitching movements and foaming at the mouth. Based on assessment the man is likely to be experiencing:
 - A. Diabetes
 - B. Major epileptic seizures
 - C. Minor epileptic seizures
 - D. Stroke
4. How would you manage a patient who has sustained a second-degree burn?
 - A. Apply Vaseline
 - B. Burst all the blisters
 - C. Place the area under cold water
 - D. Remove charred clothing from the burned area
5. As a First Aider, what would you do to reduce pain and minimize swelling in a sprain of the ankle?
 - A. Apply cold compress
 - B. Gentle massage
 - C. Lower the affected limb
 - D. Provide warmth

6. Upon discovering a Road Traffic Accident at Madina market, your first priority would be to:
- A. Ask witnesses to help the casualty
 - B. Call the emergency unit immediately
 - C. Ensure safety
 - D. Reassure the casualty
7. While assessing a casualty, you observed the casualty has a weak rapid pulse, pale, cold and clammy skin. You suspect the person is experiencing?
- A. Angina
 - B. Asphyxia
 - C. Shock
 - D. Stroke
8. What should a first aider use to clean wound?
- A. Antiseptic cream
 - B. Disinfectant
 - C. Soap and water
 - D. Water
9. Which of these objects is likely to cause a lacerated wound?
- A. A barbed wire
 - B. A bullet
 - C. A knife
 - D. A nail
10. If blood continues to seep onto the surface of the bandage, you should:
- A. Apply more pressure dressing over the original dressing
 - B. Apply a tourniquet
 - C. Remove the soiled dressing and redress
 - D. Remove the bandage and apply direct pressure
11. Abdominal thrust is used in the First Aid treatment of someone suffering from:
- A. Asthma
 - B. A diabetic coma
 - C. Choking
 - D. Strangulation

12. What do you do if someone suddenly begins to bleed severely from his left foot?

- A. Apply pressure to the wound
- B. Call for help
- C. Tell the person to lick the blood
- D. Remove foreign object

13. What will be your immediate action when someone breaks an arm?

- A. Call for help
- B. Put a plaster on the arm
- C. Put the arm in a sling
- D. Wash the area with antiseptic solution

14. What is a scald?

- A. A burn by liquid or gas
- B. A burn by fire
- C. A break in the leg
- D. A sting of a bee

15. A patient has a suspected fracture of the spine. The first action should be:

- A. Place casualty in recovery position
- B. Seek urgent medical help
- C. Secure the patient neck from side to side
- D. Test for response in toes

16. Treatment of a closed fracture of the collar bone would be:

- A. Apply pressure bandage
- B. Apply tourniquet to affected area
- C. Apply ice pack and elevate the fractured bone
- D. Immobilize limb on injured side in an (elevated) sling

17. The most effective treatment for a suspected stroke is:

EXAMINER: STELLA SAGOE

Page 4 of 15

- A. Call emergency number immediately and reassure
 - B. Commence CPR immediately
 - C. Do nothing as you can cause further damage
 - D. Place casualty in recovery position
18. A casualty with a suspected broken rib is best treated by:
- A. Giving pain relievers
 - B. Position patient in the upright position
 - C. Position patient comfortably with affected side down
 - D. Nothing, very little can be done to immobilize rib damage
19. When treating a suspected fracture of the pelvis, you should:
- A. Get the victim to hospital immediately
 - B. Elevate the lower legs
 - C. Place casualty in the lateral position and monitor
 - D. Place pelvic area into body splint to reduce movement
20. To avoid the risk of infection when treating a casualty, you should:
- A. Avoid skin-to skin contact only when blood is present
 - B. Cover face with available clothing
 - C. Do nothing for fear of becoming infected
 - D. Put gloves on before treatment is commenced for each patient
21. A person swallowed a cleaning detergent, you should:
- A. Give large amounts of fluid
 - B. Induce vomiting
 - C. Ring the poison information centre