

D. Ring the product manufacturer and seek advice

22. A person has been found in the Clonic stage of an epileptic seizure, you should:

- A. Place a wooden peg in the mouth to stop the casualty biting their tongue
- B. Restrain them to prevent self-injury
- C. Prevent injury, check airway and breathing after the seizure
- D. Seek medical assistance

23. A casualty has had a tooth knocked out during a game of football; your treatment should

be:

- A. Clean the tooth thoroughly with antiseptic
- B. Place the tooth back in position if possible or place the tooth in a glass of milk
- C. Place the tooth back in position and have the casualty clench their jaw
- D. Wrap the tooth in sterile cotton wool

24. A casualty has sand in their eye, you should:

- A. Cover eye with Vaseline
- B. Gently flush the eye
- C. Remove it with clean forceps
- D. Pull the eyelid back and wipe with gauze

25. Upon examination of a casualty who has suffered a head injury, the first thing you would

check is:

- A. Can the casualty move their toes?
- B. Is casualty responsive
- C. Whether casualty is breathing
- D. Whether casualty is bleeding

26. The treatment for a soft tissue injury is:

- A. Apply splint to reduce movement
- B. Massage the area
- C. Rest, Ice, compression, Elevation
- D. Stretch the injury to increase blood flow to the injury

27. After treating a casualty for a leg fracture, it is important to:

- A. Elevate the leg to a vertical position
- B. Lay casualty on their back to reduce possibility of movement
- C. Treat for shock and check circulation in the leg
- D. Walk the casualty around to improve circulation

28. The Heimlich maneuver is recommended for airway obstruction in children over:

- A. 1 year of age
- B. 4 years of age
- C. 8 years of age
- D. 12 years of age

29. What would you do first, if you notice a person in a restaurant choking on a piece of steak?

- A. Grab the person from behind, clasp your hands in front of the chest and squeeze hard.
- B. Encourage the person to lean forward and try to cough up the steak.
- C. Rub the person's back and offer them a drink of alcohol
- D. Sit the person and firmly push on his chest

30. How would you manage a deep, bleeding cut on the palm of the hand?
- A. Hold the injured hand under running water
 - B. Apply direct pressure to the cut, elevate the arm, then bandage the hand.
 - C. Apply pressure immobilization bandage to the arm
 - D. Apply a tourniquet bandage to the arm
31. What is the management of a conscious casualty who has diabetes and low blood sugar?
- A. Give water
 - B. Give nothing
 - C. Give sugar
 - D. Give insulin
32. After tapping the shoulder of an unconscious person, he did not respond to you. What is your next plan of action?
- A. Do CPR
 - B. Give 2 blows
 - C. Call for help
 - D. Do nothing
33. What is the ratio of compressions to breaths when performing Cardiopulmonary Resuscitation (CPR) for an adult?
- A. 15 compressions: 1 breath
 - B. 20 compressions: 2 breaths
 - C. 30 compressions: 2 breaths
 - D. 30 compressions: 5 breaths
34. The first concern in the approach to any emergency is to check:
- A. Casualty's level of consciousness
 - B. The casualty's airway
 - C. Get the casualty to hospital as soon as possible
 - D. Danger to self, others and casualty
35. To establish if the casualty is conscious the rescuer should:

- A. Check for danger, response, airway, breathing and circulation
- B. Look, listen and feel for breathing
- C. Shake the casualty vigorously
- D. Using a loud voice ask for response and tap the casualty's shoulder

36. The most common cause of airway obstruction in an unconscious casualty is:

- A. dentures
- B. the jaw
- C. the teeth
- D. the tongue

37. A casualty who is not breathing normally and shows no "sign of life" requires:

- A. Cardiopulmonary Resuscitation
- B. Chest compression
- C. To be placed in the lateral position
- D. Placed in recovery position

38. During mouth to mask rescue breaths, if the chest does not rise, this could be due to:

- A. Mucus in the mouth
- B. Poor head tilt
- C. The casualty being on their back
- D. Tight clothing

39. What techniques help to determine if a casualty is breathing?

- A. Asking them if they are having problems breathing
- B. Checking the carotid pulse
- C. Look, listen and feel

D. Shaking the casualty

40. The four main chambers of the heart are:

- A. Right and Left Ventricle, Right and left Vena Cava
- B. Right and Left Ventricle, Right and Left Atrium
- C. Right and Left Atrium, Right and left Aorta
- D. Right and left Ventricle, Right and left valve

41. If a casualty starts to breathe during CPR, you should:

- A. Commence CPR
- B. Blow until the chest rises then blow again after
- C. Place the casualty in the recovery position
- D. Stop Rescue breaths

42. The ratio for CPR on an infant casualty when two rescuers are resuscitating is:

- A. 15 breaths followed by 2 compressions
- B. 2 breaths followed by 30 compressions
- C. 1 breath followed by 5 compressions
- D. 10 breaths followed by 10 compressions

43. When performing external cardiac compression on an adult, the sternum should be depressed:

- A. Approximately 1/3 depth of the chest
- B. Approximately 1/10 depth of the chest
- C. Approximately 1/2 depth of the chest
- D. Approximately 1/20 depth of the chest

44. You should continue CPR for a patient that is not breathing for: