

A. 5 minutes

B. 1 minute

C. 4 minutes

D. Continuously

45. What will the First Aider do for a person observed to be in shock?

A. Give water for thirst and allow the casualty to rest

B. Left alone to recover

C. Provide warmth and placed in Trendelenburg position

D. Sit the casualty up and allow the casualty to rest

46. The basic principles in the treatment of bleeding are:

A. Allow to bleed freely for 30seconds and apply tourniquet

B. Allow the bleed freely for one minute and apply ice packs

C. Direct pressure to wound, elevate limb, keep casualty rested

D. Direct pressure to wound and apply tourniquet

47. A casualty has a knife impaled in their forearm, you should:

A. Cover the knife with a towel so casualty cannot see it

B. Get the casualty to the hospital as soon as possible

C. Remove the knife to reduce pain

D. Stabilize the knife with a pad, bandage around the knife

48. When caring for an amputated part of the body, you should:

A. Apply a pressure bandage over the affected area

B. Place the body part in a water proof bag and then place the bag in ice bag

- C. Try to bandage the part back in place and elevate the legs in the Trendelenburg position
- D. Wash the body part and put it in a bag of ice

49. A casualty has an open wound to the abdomen 'Evisceration', your priorities are:

- A. To allow the wound to bleed freely to stop infection
- B. To cover the wound with moist bulky dressings and treat for shock
- C. Push all the contents back into the abdomen
- D. Call for medical assistance immediately

50. After treating a casualty for a leg fracture, it is important to:

- A. Elevate the leg to a vertical position
- B. Lay casualty on their back to reduce movements
- C. Treat for shock and check circulation in the leg
- D. Walk the casualty around to improve circulation

51. The elevation sling is used for:

- A. Collarbone and shoulder injuries
- B. Forearm fractures
- C. Upper arm fractures
- D. Wrist and ankle fractures

52. The purpose of the collar and cuff sling is to:

- A. Avoid putting pressure on the collarbone
- B. Support the upper arm without putting pressure on the elbow
- C. Prevent the forearm from pressing on the broken bone
- D. Reduce movements in the forearm

53. A conscious casualty has fallen heavily from a motorbike and has tingling in the hands and feet along with neck pain, you should:
- A. Elevate the legs and check airway, breathing and circulation
 - B. Immobilize the casualty
 - C. Remove the helmet and test circulation
 - D. Sit the casualty up and remove their helmet
54. A 22-years old girl was having an asthmatic attack. After assisting with 4 puffs of inhaler and finding no improvement, you should:
- A. Call an ambulance
 - B. Commence Rescue Breaths to assist with breathing
 - C. Lay the person down and raise the legs
 - D. Lay casualty on their back to reduce movement
55. How should you open the airway of an unconscious casualty?
- A. Head tilt and chin lift.
 - B. Head tilt and jaw thrust.
 - C. Jaw thrust.
 - D. Lift the chin
56. An effective way to manage a casualty with internal bleeding is to:
- A. Keep casualty warm, administer vitamin K
 - B. Keep casualty warm and provide warm drinks
 - C. Keep casualty warm, give nothing by mouth
 - D. Keep casualty warm in semi fowler's position

57. Anaphylactic shock occurs when the individual is exposed to allergen in the environment, which one of these is not an indication of anaphylaxis?

- A. Bleeding from the nostrils
- B. Breathlessness
- C. Swelling of the air passages
- D. Rashes on the skin

58. A casualty suspected of internal haemorrhage complains of severe thirst and dry mouth, the best action to take by the first aider is:

- A. Administer intravenous fluids.
- B. Serve a cold drink
- C. Serve an ice cube and reassure him
- D. Serve a warm beverage

59. A 15-year-old boy has just splashed a chemical on his face. After sending someone to call for an ambulance, you would:

- A. Cover the burned area.
- B. Flush the burned area with large amounts of water until the ambulance arrives.
- C. Have the victim stay calm until ambulance arrives.
- D. Immediately drive the victim to the hospital

60. What should a casualty with a severe allergy carry at all times?

- A. Adrenaline (Epipen)
- B. Aspirin
- C. Acetaminophen/Paracetamol.
- D. Insulin.