

STUDENT'S INDEX NUMBER:



METHODIST UNIVERSITY GHANA
(All Rights Reserved)

FACULTY OF SCIENCE
DEPARTMENT OF NURSING & APPLIED SCIENCES

END OF FIRST SEMESTER EXAMINATIONS, 2022/2023 ACADEMIC YEAR
BSc GENERAL NURSING (DANSOMAN CAMPUS), LEVEL 300

MNUR 301: ADVANCED NURSING I
(Examiner: Nadia Abbey-Klutsey)

2.00 pm Saturday, December 3, 2022

TIME ALLOWED: 2 HOURS

INSTRUCTIONS: ANSWER ALL QUESTIONS. ANSWER SECTION A ON THE QUESTION PAPER AND SECTION B IN THE ANSWER BOOKLET PROVIDED

STUDENT'S ID NUMBER:

SECTION A [40 Marks]

Answer All Questions in This Section By Circling The Appropriate Letter Corresponding To The Correct Answer

- 1) A client is nauseated, has been vomiting for several hours, and needs to receive an antiemetic (anti-nausea) medication. Which of the following is accurate?
 - A. An enteric-coated medication should be given.
 - B. Any medication will not be absorbed as easily because of the nausea problem.
 - C. A parenteral route is the route of choice.
 - D. A rectal suppository must be administered.
- 2) The new pediatric nurse has just given a suppository to a 5-year-old boy. He has a bowel movement 7 minutes post administration. Which action should the nurse take next?
 - A. Inform the physician that the child has had a bowel movement
 - B. Give an additional half of the prescribed dose
 - C. Give another full dose of the suppository
 - D. Examine the stool for the suppository
- 3) The nurse uses a mortar and pedestal to crush a medication before giving it to one of her clients. Which of the five rights is the nurse ensuring?
 - A. The right route
 - B. The right client
 - C. The right time
 - D. The right drug

- 4) In administering an enteric-coated medication like oral Naproxen to a 70 year old geriatric patient on the female ward with dysphagia, one junior staff suggested the tablet be crushed. Which of these answers best explains why the medication should not be crushed?
- A. Because the tablet cannot be mixed in food/fluid
 - B. Crushing the tablet may lead to diarrhoea
 - C. Crushing the medication results in a faster rate of absorption by body tissues which may cause irritations to stomach lining.
 - D. None of the above
- 5) The nurse is teaching a patient about how to take a sublingual nitroglycerin tablet. Which statement by the patient best demonstrates understanding of the teaching?
- A. "I will hold the tablet next to my skin."
 - B. "I will put the tablet inside my cheek."
 - C. "I will put the tablet under my tongue."
 - D. "I will place the tablet in the lower lid of my eye."
- 6) The nurse is caring for a critically ill patient. What are the contraindications for administering medications by the oral route for this patient?
- A. Fractured leg
 - B. Penicillin allergy
 - C. Diarrhea
 - D. Unconsciousness
- 7) When is it acceptable for the nurse to take a verbal order from the prescriber before giving a drug to a patient?
- A. During the night shift when the prescriber is not at the hospital
 - B. In an emergency situation such as a cardiac arrest
 - C. When a patient is experiencing severe pain
 - D. At any time it is necessary
- 8) The pediatric nurse must convince a 4-year-old boy to take his medication. Which phrase is the most acceptable?
- A. "It is time to take your medicine. It tastes just like candy!"
 - B. "If you don't take your medicine now, you will need to take a time-out."
 - C. "If you take your medicine, I will give you a soda pop"
 - D. "Your mother and I need you to take your medicine now."
- 9) Which parts of the blood can be transfused?
- A. Platelets
 - B. Red blood cells
 - C. Whole blood
 - D. All of the above
- 10) What is the most important role of the nurse in preventing drug errors?
- A. Always checking the patient's diagnosis before giving a drug
 - B. Always following the rights of drug administration
 - C. Being the one defence for detecting and preventing drug errors
 - D. Being most likely to detect a drug error that has occurred.

- 11) What are the common risks of donating blood?
- A. Contract common viruses
 - B. Low blood pressure
 - C. Bacterial infection
 - D. None of the above
- 12) When food is given in the stomach or intestines directly then its nutrition.
- A. intravenous
 - B. saline
 - C. parenteral
 - D. enteral
- 13) Which of the following statements regarding emergency oxygen therapy is **true**?
- A. Non-rebreathe oxygen mask is the delivery device of choice in critical illness
 - B. Oxygen should be used as a treatment for breathlessness
 - C. Nasal cannula is the oxygen delivery device of choice in critical illness
 - D. Oxygen therapy should always be prescribed
- 14) Linda a 49 years old woman, with uncontrolled blood pressure and diabetics was suddenly rushed to the emergency ward following an explained collapse at her work place this afternoon. In providing first aid to Linda, an intravenous (IV) line must be secured for IV fluids to be given. Which of the following cannulae has the fastest flow-rate?
- A. 18G (green)
 - B. 22G (blue)
 - C. 20G (pink)
 - D. 14G (brown)
- 15) What administration technique does the nurse use to give a 2 yr old child ear drops?
- A. Pull the earlobe down and back
 - B. Pull the earlobe up and out
 - C. Keep the earlobe straight
 - D. Hang the patient's head over the side of the bed.
- 16) The doctor writes an order for a medication. The nurse notes the order says to administer 12.5 mg PO. Pharmacy dispenses the nurse with 6.25 mg tablets. How many tablets will the nurse administer and what medication administration right is the nurse assessing?
- A. 2 tablets, right evaluation
 - B. 2.5 tablets, right calculation
 - C. 2 tablet, right dosage
 - D. 2.5 tablets, right time
- 17) The purpose of the medication administration rights is to help prevent medication errors.
- A. True
 - B. False
 - C. Maybe
 - D. Not sure
- 18) What blood type is known as the "universal donor"?
- A. Type A
 - B. Type B
 - C. Type AB
 - D. Type O

- 19) What blood type is known as the "universal recipient"?
- A. Type A
 - B. Type B
 - C. Type AB
 - D. Type O
- 20) Before a blood transfusion you educate the patient to immediately report all the following signs and symptoms during the blood transfusion that could represent a transfusion reaction with the exception of?
- A. Sweating
 - B. Chills
 - C. Hives
 - D. Tinnitus
- 21) What substances are generally added to total parenteral nutrition solutions?
- A. Antifungal agents, electrolytes and vitamins
 - B. Citrate preservatives, electrolytes and vitamins
 - C. Electrolytes, vitamins and trace elements
 - D. Electrolytes, antihistamine antagonists and vitamins
- 22) An HIV/AIDS emaciated patient on the female medical recovery ward, needs to be fed since he is on antiretroviral drugs. However, patient is not able to tolerate oral feeds and all other attempts to initiate other methods of feeding prove futile. The only option left was to feed directly through the veins. This method is feeding id known as nutrition.
- A. Parenteral
 - B. Saline
 - C. Enteral
 - D. Intravenous
- 23) In setting an IV line for a 22 year old woman on the ward by a nurse, what materials are necessary when preparing to start an IV?
- | | |
|------------------|-----------------------------------|
| i. Gloves | iv. Eye protection |
| ii. Gown | v. Mask |
| iii. IV catheter | vi. Primed saline or heparin lock |
- A. I, ii, and vi
 - B. I, ii, and iv
 - C. I, iii and vi
 - D. I, iv and v
- 24) While in the process of inserting an IV, which of the following is not acceptable?
- A. A device may be reused as long as it is in the same site as the original attempt.
 - B. Enter the skin directly over the vein
 - C. Enter the skin at 30 – 40 degree angle
 - D. Enter the skin adjacent to the vein and direct the needle into the side of the vein wall
- 25) A patient is ordered to receive 2 units of packed red blood cells. The first unit was started at 14:00 pm and ended at 18:00 pm. You send for the other bag of red blood cells. As the nurse you know it is priority to:
- A. Hang a new bag of dextrose to transfuse with the blood
 - B. Continue with the old blood transfusion tubing set
 - C. Obtain a new blood transfusion tubing set for this unit of blood
 - D. Obtain signed informed consent for the second unit of blood from the patient

- 26) Before starting a blood transfusion the nurse will perform a verification process with _____. This will include?
- A. any available personnel; physician's order, patient's identification, blood bank's information, expiration date of blood
 - B. licensed personnel only (another RN); physician's order, patient's identification, blood bank's information, patient's blood type and donor's type along with Rh factor, expiration date, assess the bag of blood for damage or abnormal substances
 - C. blood bank; patient's identification, blood bank's information, patient's blood type and donor's type along with Rh factor, expiration date, bag of blood for damage or abnormal substances
 - D. licensed personnel only (another RN); blood compatibility, physician order, expiration date
- 27) A patient started receiving their first unit of blood at 1000. It is now 1010 and the patient is reporting itching, chills, and a headache. In addition, the patient's temperature is now 99.8°F from 98°F. Your next nursing action is:
- A. Stop the transfusion
 - B. Notify the physician
 - C. Decrease the rate of the transfusion
 - D. Reassure the patient that this is normal and will resolve in 30 minutes.
- 28) Administrations of inhaled drugs are used in all the conditions below with the exception of
- A. Asthma
 - B. Lower gastrointestinal conditions
 - C. Upper respiratory tract infections
 - D. Chronic obstructive pulmonary disease
- 29) While in the process of inserting an IV, which of the following is not acceptable?
- A. Enter the skin at 30 – 40 degree angle
 - B. Enter the skin directly over the vein
 - C. Enter the skin adjacent to the vein and direct the needle into the side of the vein wall
 - D. A device may be reused as long as it is in the same site as the original attempt.
- 30) Alex a 32 year old was rushed to the emergency ward in an unconscious state. During triaging of this patient to know the level of consciousness using the Glasgow coma scale, all the following important with the exception of.....?
- A. eye opening
 - B. vocal response
 - C. verbal response
 - D. motor response
- 31) To prevent pressure ulcers, which of the following interventions are appropriate?
- A. Frequency of skin assessments may need to be increased if the residents status deteriorates.
 - B. Keep the head of the bed above a 30 degree angle at all times to reduce pressure and shearing force on the sacral area.
 - C. When positioning or lifting up the resident in bed, health care providers should not drag skin across linens to prevent skin injury cause by friction and shearing
 - D. Both a. and c.