



METHODIST UNIVERSITY GHANA

DEPARTMENT OF NURSING AND MIDWIFERY

BSC. NURSING: END OF FIRST SEMESTER EXAMINATION: 2024/2025

**COURSE CODE/TITLE: MNUR MNUR 302 ADVANCE NURSING TWO
(3 CREDITS)**

INSTRUCTIONS:

ANSWER ALL QUESTIONS IN THIS SECTION

TIME ALLOWED: 2HOURS

EXAMINER: NADIA ABBEY

Page 1 of 2

ANSWER ALL QUESTIONS IN THIS BY CIRCLING THE MOST APPROPRIATE ANSWER FROM (A-D). ONE MARK IS ASSIGNED TO EACH QUESTION

1. Which of the following is a contraindication to performing a thoracentesis?
 - A. Suspected pleural effusion
 - B. Bleeding disorder
 - C. Pneumonia
 - D. History of heart failure

2. You are providing care to a patient with a chest tube. On assessment of the drainage system, you note continuous bubbling in the water seal chamber and oscillation. Which of the following is the CORRECT nursing intervention for this type of finding?
 - A. Check the drainage system for an air leak.
 - B. Reposition the patient because the tubing is kinked.
 - C. Continue to monitor the drainage system.
 - D. Increase the suction to the drainage system until the bubbling stops

3. A nurse is preparing to care for a client who had undergone an above-knee amputation of the right leg. The nurse plans to allow which position for the client on the first 24 hours?
 - A. Supine position with the affected limb supported with pillows.
 - B. Prone position with the affected limb on a dependent position.
 - C. Trendelenburg's position.
 - D. Supine position with the affected limb flat on the bed

4. A patient is 8 hours post-opt from an colostomy placement. Which finding requires immediate nursing action?
 - A. The stoma is not draining any stool.
 - B. The patient states the site is tender.
 - C. The stoma is swollen and large
 - D. The stoma is black.

5. A patient, who has a colostomy, asks what type of foods they should avoid to decrease odorous gas. You would tell the patient to avoid:

- A. Onions, alcoholic beverages, eggs, and cabbage
- B. Beef, fried foods, lettuce, and rice
- C. Apple, pears, nuts, and wheat
- D. Potatoes, peas, carrots, and chicken

6. A patient with a chest tube has no fluctuation of water in the water seal chamber. What could be the cause of this?

- A. This is an expected finding.
- B. The lung may have re-expanded or there is a kink in the system.
- C. The system is broken and needs to be replaced.
- D. There is an air leak in the tubing.

7. You're assessing a patient who is post-opt from a chest tube insertion. On assessment, you note there is 50 cc of serosanguinous fluid in the drainage chamber, fluctuation of water in the water seal chamber when the patient breathes in and out, and bubbling in the suction control chamber. Which of the following is the most appropriate nursing intervention?*

- A. Document your findings as normal.
- B. Assess for an air leak due to bubbling noted in the suction chamber.
- C. Notify the physician about the drainage.
- D. Milk the tubing to ensure patency of the tubes.

8. While helping a patient with a chest tube reposition in the bed, the chest tube becomes dislodged. What is your immediate nursing intervention?*

- A. Stay with the patient and monitor their vital signs while another nurse notifies the physician.
- B. Place a sterile dressing over the site and tape it on three sides and notify the physician.
- C. Attempt to re-insert the tube.
- D. Keep the site open to air and notify the physician.

9. A tracheostomy tube is inserted in a patient who is in respiratory distress as a result of pneumonia. The family asks why the tube is inserted. What should the nurse include when explaining to the patient and family? The purpose of a tracheostomy tube is to:

- A. Decrease the client's anxiety by increasing the size of the airway
- B. Facilitate nursing care since tracheal tubes have fewer side effects than nasotracheal tubes
- C. Provide more controlled ventilation and ease removal of secretions the client is unable to handle
- D. Provide increased cerebral oxygenation, thereby preventing further respiratory depression

10. An adult is about to have a tracheostomy performed. Which action is of the highest priority for the nurse before the procedure is done?

- A. Drawing blood for serum electrolytes and blood gases
- B. Doing a surgical prep of the neck and upper chest wall
- C. Establishing means of postoperative communication
- D. Inserting an indwelling catheter and attaching it to dependent drainage

11. Which of the following is NOT typically used as part of the Glasgow Coma Scale (GCS) for assessing level of consciousness?

- A. Eye opening
- B. Verbal response
- C. Motor response
- D. Heart rate

12. Postural drainage is a technique designed to:

- A. Help remove mucus via gravity
- B. Help ease your back pain
- C. Position the body so that you have to breathe harder
- D. Move your head away from your chest

13. Which of the following should be assessed regularly in a client who has just undergone a bronchoscopy?

- A. Blood pressure and pulse rate
- B. Bowel sounds
- C. Gag and cough reflexes
- D. Skin temperature

14. The colour of normal CSF is

- A. Creamy yellow
- B. Slightly pink
- C. Clear straw coloured
- D. Crystal clear

15. Which of the following is not a complication of lumbar puncture?

- A. Local back pain
- B. Epidermoid tumour implantation
- C. Olfactory disturbances
- D. Post lumbar puncture headache

16. Which of the following is an absolute contraindication to lumbar puncture?

- A. Uncorrected coagulopathy
- B. Nuchal rigidity
- C. A and B
- D. None of the above

17. The client arrives in the emergency room with a penetrating eye injury from wood chips while cutting wood. The nurse assesses the eye and notes a piece of wood protruding from the eye. What is the initial nursing action?

- A. Perform visual acuity tests
- B. Apply an eye patch
- C. Remove the piece of wood using a sterile eye clamp
- D. Irrigate the eye with sterile saline

18. The client arrives in the emergency room after sustaining a chemical eye injury from a splash of battery acid. The initial nursing action is to:

- A. Begin visual acuity testing
- B. Cover the eye with a pressure patch
- C. Irrigate the eye with sterile normal saline
- D. Swab the eye with antibiotic ointment

19. When using a Snellen alphabet chart, the nurse records the client's vision as 20/40. Which of the following statements best describes 20/40 vision?

- A. The client has alterations in near vision and is legally blind
- B. The client can see at 20 feet what the person with normal vision can see at 40 feet
- C. The client can see at 40 feet what the person with normal vision sees at 20 feet
- D. The client has a 20% decrease in acuity in one eye and a 40% decrease in the other eye

20. Which of the following is a common indication for an antral washout?

- A. Acute pharyngitis
- B. Chronic maxillary sinusitis unresponsive to medical treatment
- C. Tonsillitis
- D. Middle ear effusion

21. A patient with a newly applied leg cast complains of severe, throbbing pain and numbness in their foot. What is the MOST IMMEDIATE concern?

- A. Skin breakdown
- B. Compartment syndrome
- C. Infection
- D. All of the above

22. Which of the following is a contraindication for casting?

- A. A recent fracture
- B. Severe soft tissue swelling
- C. An open fracture
- D. A closed fracture

23. Which of the following is a sign of cast-related neurovascular compromise?

- A. Mild itching under the cast
- B. Aching pain in the affected limb
- C. Cyanosis of the distal extremity
- D. A small pressure sore

24. A newly RN nurse is about to insert a nasogastric tube to a client with Guillain-Barre Syndrome. To determine the accurate measurement of the length of the tube to be inserted, the nurse should:

- A. Place the tube at the tip of the nose, and measure by extending the tube to the earlobe and then down to the xiphoid process.
- B. Place the tube at the tip of the nose, and measure by extending the tube down to the chin and then down to the top of the xiphoid process.
- C. Place the tube at the base of the nose, and measure by extending the tube to the earlobe and then down to the top of the sternum.
- D. Place the tube at the tip of the nose and measure by extending the tube to the earlobe and then down to the top of the sternum.

25. 'Spigotting' a tube means:
- A. Opening it on free drainage
 - B. Attaching it to a bag
 - C. Cutting the tube to make it shorter
 - D. Clamping or plugging the end of the tube
26. The nurse is caring for a client who develops compartment syndrome from a severely fractured arm. The client asks the nurse how this can happen. The nurse's response is based on the understanding that:
- A. A bone fragment has injured the nerve supply in the area
 - B. An injured artery causes impaired arterial perfusion through the compartment
 - C. Bleeding and swelling cause increased pressure in an area that cannot expand
 - D. The fascia expands with injury, causing pressure on underlying nerves and muscles
27. After performing gastric lavage, the nurse should assess the patient for signs of:
- A. Increased appetite
 - B. Abdominal distention
 - C. Hypertension
 - D. Elevated body temperature
28. Which of the following is a potential complication of gastric lavage?
- A. Improved gastric motility
 - B. Electrolyte imbalances
 - C. Increased blood pressure
 - D. Enhanced kidney function
29. You just inserted a Nasogastric tube. Which of the following is not a correct way to check correct placement of the tube?
- A. Obtaining a sample of GI contents through the tube by aspirating
 - B. Following the MD order for an X-ray to confirm placement
 - C. Checking pH of GI contents to be at 1 to 3.5
 - D. Administering a 50cc Water flush and assessing for patient coughing
30. Which of the following conditions might necessitate the placement of a gastrostomy tube?
- A. Lung cancer
 - B. Gallstones
 - C. Stroke with dysphagia
 - D. Urinary tract infection
31. After a fractional test meal, the nurse should assess the patient for signs and symptoms of:
- A. Constipation
 - B. Gastric ulcer perforation
 - C. Allergic reaction
 - D. Hypoglycemia
32. Which complication is commonly associated with an ileostomy?
- A. Constipation
 - B. Dehydration
 - C. Fecal impaction
 - D. Bowel obstruction
33. Before performing a paracentesis, what should the nurse confirm about the patient?
- A. Recent food intake
 - B. History of asthma
 - C. Allergies to seafood
 - D. Coagulation status

34. A client is leaving the unit to go to hemodialysis. In which way should the nurse manage the client's medications?
- A. Hold the client's antihypertensives because of the risk of hypotension during hemodialysis
 - B. Only subcutaneous and IV medications may be given prior to dialysis
 - C. Do not give any of the client's medications the morning of dialysis because they will all be removed during dialysis
 - D. Give all of the client's medications since they will be gone for several hours
35. Which vascular access option is considered the gold standard for long-term hemodialysis?
- A. Hemodialysis catheter
 - B. Arteriovenous graft (AVG)
 - C. Central venous catheter
 - D. Arteriovenous fistula (AVF)
36. The nurse is caring for a client who will be receiving a suprapubic catheter. What should the nurse mention to the client?
- A. These catheters are often preferred over an indwelling urethral catheter for long-term urinary drainage
 - B. Press the appropriate gender button, position the scanner head with the directional arrowing point to the head, press the scanner head onto the skin 1 to 1.5 inches above the symphysis pubis, aim the scanner head toward the coccyx and activate the scan, verify that screen crossbars fall within the bladder image, observe and record volume
 - C. The client will have to wear an external appliance to collect urine
 - D. Trauma-related muscle damage, menopausal hormonal changes, prolonged periods of immobility, use of an indwelling urinary catheter, and childbearing
37. What is the most common complication of thoracentesis?
- A. Pneumonia
 - B. Pulmonary edema
 - C. Pneumothorax
 - D. Infection
38. What is the purpose of performing a hysterosalpingogram?
- A. To diagnose possible reasons for infertility
 - B. To treat infertility
 - C. To treat PID (Pelvic Inflammatory Disease)
 - D. All answers are correct
39. A client sustained a radial fracture and a cast was just applied. The client states that there is unrelieved pain and numbness in the fingers on the affected side. Which intervention should be a priority?
- A. Notifying the healthcare provider for cast removal
 - B. Elevating the extremity
 - C. Preparing for fasciotomy
 - D. Performing frequent neurovascular checks
40. Which blood sample is typically used for routine blood glucose testing?
- A. Arterial blood
 - B. Venous blood

C. Capillary blood

D. Intravenous blood

41. What is the advantage of echocardiography over other cardiac imaging modalities such as CT or MRI?

A. Echocardiography is less expensive

B. Echocardiography does not use radiation

C. Echocardiography provides better resolution of bone structures

D. Echocardiography can image lung tissue

42. Which waveform represents the time interval between ventricular depolarization and repolarization on an ECG tracing?

A. PR interval

B. QT interval

C. ST segment

D. QRS complex

43. Which component of the heart's electrical activity does the P wave represent on an ECG tracing?

A. Ventricular depolarization

B. Atrial depolarization

C. Ventricular repolarization

D. Atrial repolarization

44. What is the purpose of coronary angiography during cardiac catheterization?

A. To measure blood pressure in the heart chambers

B. To assess heart valve function

C. To visualize blockages or narrowing in the coronary arteries

D. To monitor electrical activity of the heart

45. Which of the following complications is associated with cardiac catheterization?

A. Increased heart rate

B. Hypertension

C. Bleeding

D. Hypoglycemia

46. The T wave is also referred to as:

A. Atrial depolarization

B. Ventricular repolarization

C. Atrial repolarization

D. Ventricular depolarization

47. When a catheter is used to clear a narrowed or blocked artery, the procedure is called.....

A. Angioplasty

B. Angiography

C. Valvuloplasty

D. All of the above

48. P wave indicates

A. Depolarization of right ventricle

B. Depolarization of left ventricle

C. Depolarization of both atria

D. Atria to ventricular conduction time

49. Ventricular muscle depolarization is indicated by

A. PR interval

B. P wave

C. U wave

D. The QRS complex

50. Echocardiography can be used for even _____

A. remove embolus of the heart

B. correctly locate the blockages in the arteries and veins

C. assess the baby's heart, anatomy and function

D. measure heartbeat

51. Echocardiography is essentially _____

A. ultrasound of the heart

B. echoing sound of the heart

C. another name for a treadmill test

D. recording of heart sounds

52. Which of the following is true about a typical pleurocentesis procedure?

A. It lasts about 10-15 minutes.

B. It's done using a needle and, in some cases, an X-ray or ultrasound.

C. The patient is sitting up and leaning forward during it.

D. All of the above

53. What is monitored during and after a thoracentesis?

A. Breathing and blood pressure

B. Heart rhythm and temperature

C. Blood pressure and steadiness

D. Mental status and heart rate