

INDEX NUMBER.....DATE.....

METHODIST UNIVERSITY COLLEGE GHANA



FACULTY OF SCIENCE

DEPARTMENT OF NURSING & APPLIED SCIENCE

(DANSOMAN & TEMA COMPUS)

END OF FIRST SEMESTER EXAMINATION, 2024/2025

B.S.c REGISTERED GENERAL NURSING

COURSE CODE: MNUR 303

COURSE TITLE: MEDICAL NURSING II

TIME: 2 Hours

INSTRUCTIONS:

1. DO NOT OPEN THE BOOKLET UNTIL YOU ARE TOLD TO DO SO
2. READ THE QUESTIONS CAREFULLY AND PROVIDE ONLY RELEVANT ANSWERS
3. WRITE CLEARLY IN THE SPACES PROVIDED OR CIRCLE APPROPRIATELY USING A BLUE PEN
4. STOP WRITING AS SOON AS YOU ARE INSTRUCTED TO DO SO
5. REMAIN IN YOUR SEAT UNTIL THE PAPERS ARE COLLECTED
6. ANSWER SECTION B IN ANSWER BOOKLET

SECTION A

1. Peptic ulcer may be classified according to the following organs EXCEPT
 - A. Duodenal
 - B. Gastric
 - C. Ileal
 - D. Oesophageal
2. A type of chest pain characterized by severe incapacitating chest pain despite treatment.
 - A. Intractable Angina
 - B. Stable angina
 - C. Myocardial infarction
 - D. Vagina pectoris
3. Clinical manifestation of pulmonary congestion include the following EXCEPT.
 - A. Cough
 - B. Dyspnea
 - C. Nausea
 - D. Pulmonary crackles
4. Which of the following agents is described as H₂ – Receptor antagonist?
 - A. omeprazole
 - B. sulphathalarole
 - C. ranitidine
 - D. amoxicillin
5. The nursing care for a patient suffering from pneumonia does NOT include :
 - A. tepid sponging
 - B. ensuring a high fluid intake of preferred drinks
 - C. isolating the bacteria that caused the infection
 - D. control of pain to ensure comfort.
6. Myocardial infarction is characterized by a prolonged
 - A. increased supply of blood and oxygen to the tissues
 - B. substernal oppression or pain
 - C. radiating pains from the heart to the inner aspects of the arms.
 - D. emotional change due to fear of death.
7. The nursing care for a patient suffering from pneumonia does NOT include
 - A. tepid sponging
 - B. ensuring a high fluid intake of preferred drinks
 - C. isolating the bacteria that caused the infection
 - D. control of pain to ensure comfort.
8. Which of the following common signs or symptoms indicate that the patient is suffering from myocardial infarction?
 - A. The client experiencing nausea and vomiting
 - B. The client complains of pain while walking
 - C. Client's pain is not relieved by rest and nitroglycerin
 - D. Client's pain is described as sub-sternal and radiates to the left arm
9. Non –pharmacological treatment of peptic ulcer include the following EXCEPT
 - A. Cessation of smoking

- B. Avoid foods that aggravate the pain
- C. Relief of anxiety and stress
- D. Promote healing of the ulcer

10. In nursing a client who has a potential for haemorrhage, the nurses' goal include the following except

- A. Absence of all subjective and objective signs of gastrointestinal, vascular and soft tissue haemorrhage
- B. Early detection of bleeding
- C. Observing client's vital signs at frequent interval.
- D. Serving client's with iron-rich foods.

11. A patient with myocarditis is on digitalis. The nurse knows that patients with myocarditis are sensitive to digitalis, her priority intervention will be:

- A. To closely monitor for digitalis toxicity
- B. To closely monitor for pulse
- C. To closely monitor for blood pressure
- D. To call the doctor

12. Which of the following is NOT a common symptom of GERD?

- A. Dyspepsia
- B. Dysphagia
- C. Hyposalivation
- D. Regurgitation

13. A client with angina pectoris complains of chest pain while getting out of bed. The nurse must first:

- A. Get an order for pain medication
- B. Report the complaint to the physician
- C. Advise the client to continue to get out of bed
- D. Stop the client and encourage him to lie down in bed

14. In nursing management of a patient diagnosed of hypertension, the nurse's priority is

- A. to identify the primary cause and treat
- B. inform the patient that hypertension is incurable
- C. to inform the patient that hypertension is curable
- D. Restrict the patient from taking food constraining sodium.

15. In duodenal ulcer, the pain may wake the patient up in the

- A. middle of the night
- B. early morning
- C. late in the night
- D. late in the afternoon

16. After feeding a patient through a nasogastric tube, the nurse places the patient in which position?

- A. Supine
- B. Fowler's
- C. Sim's
- D. Trendelenburg

17. Dr. Akambong has determined that the client with hepatitis has contracted the infection from contaminated food. The nurse understands that this client is most likely experiencing what type of hepatitis?

- A. Hepatitis A
- B. Hepatitis B

- C. Hepatitis C
- D. Hepatitis D

18. Both sickle cell and thalassemia seems to confer relative resistant to:

- A. Syphilis
- B. Tuberculosis
- C. Malaria
- D. Leprosy

19. A characteristic crushing and constricting chest pain lasting at least 20 minutes due to decrease blood supply to the myocardium is known as?

- A. Heart failure
- B. Myocardial ischemia
- C. Heart attack
- D. Angina pectoris

20. Chronic bronchitis co-exists with emphysema when there is

- A. Decrease number of alveoli with increase demand of oxygen
- B. Enlargement of alveoli with destruction of their walls
- C. Over-secretion of mucus and increase expectoration of mucus.
- D. Is dilation of the bronchi and bronchioles.

21. A nurse should include which of the following non-pharmacological treatment plan for a client who has peptic ulcer.

- A. Continue to eat the same diet as before diagnosis
- B. Smoke only at bedtime
- C. Learn to use stress reduction techniques
- D. Take non-steroidal anti-inflammatory drugs for pain relief.

22. The best psychological management of medical conditions is to:

- A. Shout at patients and relatives
- B. Reassure the client and relatives
- C. Serve them with a strong coffee
- D. Tell relative to go home, leaving the patient at O.P.D.

23. Right ventricular failure (Right-sided-heart failure) is treated by the combination of

- A. morphine and frusemide
- B. rest, digitalis and diuretics
- C. morphine, antibiotics and ant diuretics
- D. morphine alone.

24. Nurses do not have the responsibility for determining drug dosage for their patients but they

- A. Do so in the absence of the physician
- B. Do need to be aware of the normal doses of drugs
- C. Prescribe emergency drugs
- D. Only serve drugs.

25. The primary action of the nurse in attending to an unconscious patient suffering from severe hypertension is to

- A. serve the prescribed drugs immediately
- B. check the vital signs immediately
- C. inform the physician immediately
- D. inform the relatives of the policy of the hospital

26. The basic factors that may cause peptic ulcer do NOT include one of the following:
- A. excessive secretion of gastric acid
 - B. inadequate protection of the lining of stomach and duodenum against digestion by a base and pepsin
 - C. helicobacter pylori infection
 - D. drug eg. NSAID'S
27. Before a nurse serves digoxin to a client, she should ...
- A. Check the pulse rate
 - B. Check the blood pressure
 - C. Check the urine output
 - D. Check the respiratory rate
28. The collapse of the lungs is known as?
- A. Alveoli enlargement
 - B. Atelectasis
 - C. Sleep apnea
 - D. Telesis
29. A client diagnosed of thrombophlebitis 24 hours ago suddenly complains of chest pain, shortness of breath and visible anxiousness. The nurse must check the client for other signs and symptoms of
- A. myocardial infarction
 - B. pneumonia
 - C. pulmonary embolism
 - D. pulmonary oedema
30. In Pulmonary disease (COPD), which nursing action best promotes adequate gas exchange?
- A. Administering a sedative as prescribed
 - B. Encouraging the client to drink three glasses of fluid daily
 - C. Keeping the client in semi-Fowler's position
 - D. Using a high-flow Venturi mask to deliver oxygen as prescribed
31. In a patient diagnosed with leukemia, the leucocyte count usually
- A. Rises above normal range
 - B. Is within normal range
 - C. Is below normal range
 - D. Is difficult to be determined
32. If a patient has difficulty in breathing while lying down flat, the condition is called:
- A. Apnoea
 - B. Orthopnoea
 - C. Dyspnoea
 - D. Hyperpnoe
33. A client was brought to an emergency department with angina pectoris, the priority of the nursing care is to:
- A. assemble oxygen apparatus at the bedside
 - B. put in place a bed table
 - C. make ECG machine ready
 - D. put the client on a low bed

34. Nurse Yakubu is caring for a male client with a diagnosis of Peptic ulcer. The nurse monitors the client knowing that this client is at risk for which vitamin deficiency?

- A. Vitamin A
- B. Vitamin B12
- C. Vitamin C
- D. Vitamin E

35. Nurse Sibdow is caring for a client with pneumothorax who has had a chest tube inserted, notes a continuous bubbling in the suction control chamber. What action is appropriate.

- A. Check for an air leak
- B. Do nothing, because this is an expected finding
- C. Immediately clamp the chest tube and notify the physician
- D. Increase the suction pressure

36. In pericarditis, which of the following clinical features differentiates it from other cardiopulmonary disease?

- A. Chest pain that worsens on inspiration
- B. Pericardial friction rub
- C. Anterior chest pain
- D. Weakness and irritability

37. Which of the following does NOT exacerbate the condition of a patient suffering from heart failure?

- A. recent upper respiratory infection
- B. nutritional anaemia
- C. peptic ulcer disease
- D. atrial fibrillation

38. A client states that angina pain increases after activity. The nurse should realize that angina pectoris is a sign of:

- A. Mitral insufficiency
- B. Myocardial ischaemia
- C. Myocardial infarction
- D. Coronary thrombosis

39. A nurse reviews the physician's prescription for a client with heart failure. The nurse expects to note which of the following?

- A. Cardizem
- B. Digoxin
- C. Propranolol
- D. Metronidazole

40. The molecule involved in oxygen transport within the red cell that gives blood its characteristic red colour is called.....

- A. fibrinogen
- B. haemoglobin
- C. prothrombin
- D. glucagon

41. Ama, 25 year old arrived in your ward with the diagnosis of peptic ulcer disease. What is the first nursing action to be instituted?

- A. Administer ordered medication
- B. Assess the level of abdominal pain
- C. Instruct patient to lie down
- D. Provide oral care

42. Which of the following measures will involve a non-pharmacological treatment of acute myocardial infarction?
- To ensure bed rest
 - To relieve pain
 - To treat complications
 - To prevent re-infarction
43. A client complains that low-salt food is very tasteless. The nurse's best response would be:
- Let's consider some alternatives
 - Salt can be harmful to your health
 - Ask the doctor to increase the intake for you occasionally
 - It must be difficult for you, but you will get used to it.
44. Mr. Rockson was admitted to male medical ward of BMC with a provisional diagnosis of malaria from the OPD. On the next day morning, the Doctor reviewed his condition with diagnostic investigations and confirmed the diagnosis to be pneumonia. Mr. Rockson is likely to be suffering from which type of pneumonia based on the duration of hospitalization?
- Bacterial pneumonia
 - Secondary pneumonia
 - Hospital acquired pneumonia
 - Community acquired pneumonia
45. They are vasoactive agents which relaxes vessels and increase oxygen supply to the myocardium.
- Calcium channel blockers
 - Propranolol
 - Nitrates
 - Diuretics
46. Peptic ulcer may be classified according to the following organs except
- duodenal
 - gastric
 - oesophageal
 - ileal
47. Status asthmaticus is defined as
- A chronic asthma that is often treated with drugs
 - A slowly but progressive asthma that is relieved by repositioning of the patient
 - A period when the patient has regain health after an attack
 - A severe asthma that has not responded to patient's normal
48. The basic factors that may cause peptic ulcer do NOT include one of the following:
- excessive secretion of gastric acid
 - inadequate protection of the lining of stomach and duodenum against digestion by a base and pepsin
 - helicobacter pylori infection
 - drug eg. NSAID'S
49. The success of pain management primarily depends on the
- type of drugs to be used
 - the type of pain and its location
 - the therapist and the patient
 - the patient's pain threshold

50. A client has been admitted with chest trauma after vehicle accident and has undergone intubation. The nurse notes absence of breath sound in the right upper lobe of the lung. The nurse immediately assess for other signs of:

- A. Pulmonary embolism
- B. Right pneumothorax
- C. Displaced endotracheal tube
- D. Acute respiratory distress syndrome

51. The hormone erythropoietin controls the production of

- A. platelets
- B. leukocytes
- C. red blood cells
- D. the release of prothrombin

52. The primary action of the nurse in providing care for a patient diagnosed of severe heart failure is

- A. to prescribe anti hypertensive drugs
- B. to restrict patient from performing an activity
- C. talk to patient's relative about the need to support patient to recover
- D. to identify the cause and treat promptly with drugs

53. Collection of pus in the pleural cavity is known as:

- A. Emphysema
- B. Empyema
- C. Pneumothorax
- D. Pulmonary embolism

54. A nurse is collecting data on a hepatitis patient, which of the following suggest a severe damage to the liver?

- A. Dark stools, yellow sclera and dark urine
- B. Clay stools, yellow sclera, and blood tinged urine
- C. Clay stools, pruritus and dark urine
- D. Dark stools, pruritus, and amber urine

55. Nurse Rockson is teaching a male client with chronic respiratory failure how to use a metered-dose inhaler correctly. The nurse instructs the client to:

- A. Hold the breath after inhalation
- B. Inhale quickly
- C. Inhale through the nose
- D. Take two inhalations during one breath

56. In nursing management of a patient diagnosed of hypertension, the nurse's priority is

- A. To identify the primary cause and treat
- B. Inform the patient that hypertension is incurable
- C. To inform the patient that hypertension is curable
- D. Restrict the patient from taking food constraining sodium.

57. Which of the following agents is described as proton pump inhibitor (PPI)?

- A. omeprazole
- B. sulphathalazole

- C. ranitidine
- D. amoxicillin

58. The priority nursing diagnosis of the acute phase of myocardial infarction is:

- A. Alteration in comfort (pain)
- B. Anxiety
- C. Ineffective coping
- D. Impaired tissue integrity

59. Left ventricular failure (left sided heart failure) is best treated with.....

- A. intravenous injection of digitalis and antibiotics
- B. intravenous injection of digitalis alone
- C. intravenous injection of morphine, aminophylline and frusemide
- D. intravenous injection of aminophylline and frusemide

60. In providing instructions to a client with peptic ulcer disease, the nurse tells the client to..

- A. Consume a bland diet only
- B. Eat foods that will increase gastrointestinal motility.
- C. Eat at least six small meals per day
- D. Eat everything that does not aggravate or cause pain.

SECTION B: ESSAY

QUESTION ONE

Miss Lydia, a 32 year old trader lives in a single room with her two children. She has been diagnosed as having Bacterial Pneumonia and was admitted to the female medical ward. On the third day of admission, she complained of pain occurring only on one side of the lung and worsens with deep breaths, coughing or sneezing.

- A. Briefly describe Pneumonia (5 mark)
- B. State FIVE (5) clinical manifestations of Bacterial Pneumonia (5 marks)
- C. Using the nursing process as a framework, formulate TEN (10) nursing interventions for ineffective airway clearance related to inability to cough effectively. (10 marks)

Total: 20 marks

QUESTION TWO

Mr. Edward, a 25 year young man is admitted to the Male Medical Ward of BMC with a diagnoses of Peptic Ulcer Disease (PUD) with Anaemia.

- a. What type of anaemia is he suffering from? (2 mark)
- b. Enumerate the combination of the triple therapy medications for peptic ulcer (6 marks)
- c. List four (4) clinical manifestations of anaemia (2 marks)
- d. Describe the nursing intervention for peptic ulcer (10 marks)

Total: 20 marks