

58. A 5-month old child develops severe diarrhoea and has been put on IV fluids. The nurse must observe the rate of flow to
- A. Avoid fluid getting into the tissue
 - B. Prevent cardiac overload
 - C. Prevent increased urinary output
 - D. Replace all fluids lost
59. In pyloric stenosis, food cannot move normally in the G.I tract from the
- A. Duodenum into the jejunum
 - B. Jejunum into the ileum
 - C. Oesophagus into the stomach
 - D. Stomach into the duodenum
60. Which of the following defects results in decreased pulmonary blood flow?
- A. Atrial septal defect
 - B. Tetralogy of fallout
 - C. Ventricular septal defect
 - D. Patent ductus arteriosus
61. Physiologic jaundice in a neonate can be caused by which of the following?
- A. Fetal-maternal blood incompatibility
 - B. Destruction of red blood cells as a result of antibody reaction
 - C. Liver's inability to hydrolyze and excrete bilirubin
 - D. Maternal error
62. Several complications can occur when a child receives a blood transfusion. Which of the following is an immediate sign/symptom of air emboli?
- A. Chills and shaking
 - B. Nausea and vomiting
 - C. Irregular heart rate
 - D. Sudden difficulty in breathing
63. Kernig's sign is a characteristic sign on
- A. Brain tumour
 - B. Cerebral palsy
 - C. Meningitis
 - D. Osteomyelitis
64. The Apgar score of a neonate 5 minutes after birth is 8. Which of the following is the nurse's best interpretation of this?
- A. Resuscitation is likely to be needed.
 - B. Adjustment to extra uterine life is adequate
 - C. Additional scoring in 5 more minutes is needed
 - D. Maternal sedation or analgesia contributed to low score
65. The primary purpose for regulating nursing practice by the state is to protect
- A. Practical nursing
 - B. Professional standards
 - C. The public
 - D. The employing agency
66. The main purpose of passing a urethral catheter after surgical repair of hypospadias is
- A. To decrease pain at the surgical site
 - B. To keep the new urethra from closing
 - C. To measure urine correctly

D. To prevent bladder spasm

67. Nursing care for a 7-year-old girl admitted with tetanus should be primarily directed towards

- A. Careful monitoring of urinary output
- B. Decreasing external stimuli
- C. Encouraging high intake of fluid
- D. Giving adequate diet

68. Which of the following clinical manifestations would be the most suggestive of acute appendicitis?

- A. Rebound tenderness
- B. Bright red or dark red rectal bleeding
- C. Abdominal pain that is relieved by eating
- D. Abdominal pain that is most intense at McBurney point

69. Which of the following is a symptom of increased intracranial pressure?

- A. Facial flushing
- B. Photophobia
- C. Tachycardia
- D. Vomiting

70. Hypospadias is a congenital malformation in which the urethra in the male

- A. Fails to develop
- B. Fails to open
- C. Opens on the lower surface of the penis
- D. Opens on the upper surface of the penis

71. A neural tube defect that is not visible externally in the lumbosacral area would be called:

- A. Meningocele
- B. Myelomeningocele
- C. Spina bifida cystica
- D. Spina bifida occulta

72. The defects associated with tetralogy of Fallot include:

- I. Ventricular septal defect
 - II. Mitral valve stenosis
 - III. Right ventricular hypertrophy
 - IV. Overriding aorta
- A. I and II only
 - B. I and III only
 - C. I, III and IV only
 - D. I, II, III and IV

73. The nurse understands that in patent ductus arteriosus, blood shunt

- A. Aorta into the pulmonary artery
- B. Inferior vena cava to left atrium
- C. Left atrium to right atrium
- D. Pulmonary artery to the aorta

74. The complications of poliomyelitis include

- A. Deafness and hypertension
- B. Deafness and muscle wasting
- C. Hypertension and muscle wasting
- D. Muscle wasting and respiratory paralysis

75. Which of the following statements best describes Hirschprung disease?
A. The colon has an aganglionic segment
B. There is a passage of excessive amounts of Meconium in the neonate
C. It results in excessive peristaltic movements within the gastrointestinal tract
D. It results in frequent evacuation of solids, liquid and gas.
76. It is important that a child with acute streptococcal pharyngitis be treated with antibiotics in order to prevent:
A. Otitis media
B. Diabetes insipidus
C. Nephritic syndrome
D. Acute fever
77. Which of the following problem is most often associated with myelomeningocele?
A. Hydrocephalus
B. Craniosynostosis
C. Biliary atresia
D. Esophageal atresia
78. Excessive vomiting can result in which of the following in an infant with pyloric stenosis?
A. Electrolysis
B. Hyponatremia
C. Metabolic acidosis
D. Metabolic alkalosis
79. In which of these communicable diseases are Koplik spots present?
A. German measles (Rubella)
B. Measles (rubeola)
C. Chickenpox (varicella)
D. Exantema subitum (roseola)
80. An infant with hydrocephalus is hospitalized for surgical placement of a ventriculo-peritoneal shunt. Postoperative nursing care should include which of the following?
A. Monitor closely for signs of infection
B. Pump the shunt reservoir to maintain patency
C. Administer sedation to decrease irritability
D. Maintain trendelenburg position to decrease pressure on the shunt
81. Which of the following will make you suspect that a child with measles is developing bronchopneumonia?
A. Increased pulse rate
B. Increased respiration
C. Redness of eyes
D. Wheezing
82. The mother of a ten-month old child with diarrhoea will be advised to
I. Breast feed child
II. Give coconut water
III. Serve light soup without pepper
VII. Starve child at least for one hour
A. I and II only
B. II and IV only
C. I, II and III only
D. II, III and IV only

83. A temporary gastrostomy is done on a 7-year old child with oesophageal atresia. Gastrostomy is performed to:
- A. Alleviate pain and correct the deformity
 - B. Prevent spread of the disease
 - C. Provide a method of feeding
 - D. Relieve ascites and reduce secretions
84. When caring for a child with probable appendicitis, the nurse should be alert to recognize that which of the following is a sign of perforation?
- A. Fever
 - B. Anorexia
 - C. Sudden relief from pain
 - D. Decreased abdominal distention
85. The following criterion is most likely to be used by the physician to select drug for a 6 year old with pneumonia
- A. Child tolerance of the drug
 - B. Physician's preference for the drug
 - C. Selectivity of the causative organism
 - D. Sensitivity of the causative organism
86. Which of the following terms refers to opacity of the crystalline lens that prevents light rays from entering the eye and refracting them on the retina?
- A. Myopia
 - B. Amblyopia
 - C. Cataract
 - D. Glaucoma
87. One of the complication of pinworm infections is
- A. Appendicitis
 - B. Hepatitis
 - C. Pneumonia
 - D. Stomatitis
88. Nursing care of the newborn with oral candidiasis (thrush) includes which of the following?
- A. Avoid use of pacifier
 - B. Remove characteristic white patches with a soft cloth
 - C. Continue medication for prescribed number of days
 - D. Apply medication to oral mucosa, being careful that none is ingested.
89. Stroking the neonate's cheek along the side of the mouth causes the infant to turn the head toward that side and begin to suck. This is which of the following reflexes?
- A. Perez
 - B. Sucking
 - C. Rooting
 - D. Extrusion
90. What is the earliest age at which a satisfactory radial pulse can be taken in children?
- A. 1 year of age
 - B. 2 year of age
 - C. 3 year of age
 - D. 4 year of age
91. A nursing responsibility when caring for an infant with suspected pyloric stenosis would be which of the following?

- A. Observe for projectile vomiting
- B. Monitor for signs of metabolic acidosis
- C. Administer large, infrequent feedings to allow for rest
- D. Place infant in supine position after feedings

92. Which of the following interventions may decrease the incidence of physiologic jaundice in a healthy term infant?

- A. Institute early and frequent feedings
- B. Place infant's crib near window for exposure to sunlight
- C. Bathe infant when axillary temperature is 36.30C
- D. Suggest that mother initiate breast-feeding when danger of jaundice is past.

93. Which one of the under listed cranial nerves is affected when facial paralysis occurs?

- A. 4th cranial nerve
- B. 5th cranial nerve
- C. 6th cranial nerve
- D. 7th cranial nerve

94. Haemolytic jaundice is characterized by;

- A. Abnormality in the red blood cells
- B. An increase in red and white blood cells
- C. Decreased numbers of red blood cells
- D. Over destruction of red blood cells

95. How long after birth does physiological jaundice usually occurs?

- A. 24 hours
- B. 48 hours
- C. 72 hours
- D. 1 week

96. Which of the following observations of the cerebrospinal fluid will confirm the diagnosis of meningitis?

- A. Amber coloured
- B. Blood stained
- C. Cloudy
- D. Crystalline clear

97. Which of the following is NOT an appropriate intervention to provide comfort for the child with itching associated with chickenpox?

- A. Encourage frequent warm baths
- B. Give aspirin for fever
- C. Give an antipruritic medication such as Benadryl
- D. Apply thick coat of calamine lotion over lesions

98. The nurse is caring for a neonate receiving an exchange transfusion for hemolytic disease. Assessment of the infant reveals slight respiratory distress, tachycardia and presence of rash. The nurse's first action should be which of the following?

- A. Notify practitioner
- B. Stop the transfusion
- C. Administer calcium gluconate
- D. Monitor vital signs electronically

99. Acute diarrhea is often caused by which of the following?

- A. Celiac disease
- B. Antibiotic therapy

- C. Immune deficiency
- D. Protein malnutrition

100. Which of the following is a major long-term problem for a child with cleft lip and palate?

- A. Emotional problems
- B. Abnormally small maxilla
- C. Faulty dentition
- D. Aspiration

101. The newborn with severe jaundice is at risk of developing which of the following?

- A. Encephalopathy
- B. Bullous impetigo
- C. Respiratory distress
- D. Blood incompatibility

102. Which of the following is the viral pathogen that frequently causes acute diarrhea in young children?

- A. Giardia
- B. Shigella
- C. Rotavirus
- D. Salmonella

103. An important nursing intervention when caring for a child who is experiencing a seizure would be which of the following?

- A. Describe and record the seizure activity observed
- B. Restrain the child when seizure occurs to prevent bodily harm
- C. Place a tongue blade between the teeth if they become clenched
- D. Suction the child during a seizure to prevent aspiration

104. The nurse should explain to the parents that their child is receiving lasix for severe congestive heart failure because it is:

- A. A diuretic
- B. A beta-blocker
- C. A form of digitalis
- D. An ACE inhibitors

105. At what age should the nurse expect the anterior fontanel to close?

- A. 2 months
- B. 2-4 months
- C. 6-8 months
- D. 12-18 months

106. The congenital abnormality in which a sac is formed on the umbilical area is

- A. Meningocele
- B. Omphalocele
- C. Pyloric stenosis
- D. Umbilical hernia

107. The mother of a ten-month old child with diarrhoea will be advised to

- I. Breast feed child
 - II. Give coconut water
 - III. Serve light soup without pepper
 - IV. Starve child at least for one hour
- A. I and II only
 - B. II and IV only

- C. I, II and III only
- D. II, III and IV only

108. Which of the following statements is the most descriptive of bronchial asthma?

- A. There is heightened airway reactivity
- B. There is decreased resistance in the airway
- C. The single cause of asthma is an allergic hypersensitivity
- D. It is inherited

109. A procedure in which a baby's blood is withdrawn and is replaced with fresh blood is known as

- A. Direct transfusion
- B. Exchange transfusion
- C. Replacement therapy
- D. Top up transfusion

110. Which of the following under listed statements is a symptom of meningitis?

- A. A negative Babinski response
- B. Areas of paraesthesia of lower extremities
- C. Peri-orbital oedema
- D. Resistance to neck flexion

111. The Glasgow coma scale consist of an assessment of :

- A. Pupil reactivity motor response
- B. Eye opening, verbal and motor response
- C. Level of consciousness, verbal response
- D. Intracranial pressure and level of consciousness

112. What care would you give to a baby undergoing phototherapy?

- I. Ensure adequate fluid intake
 - II. Ensure maximum skin exposure
 - III. Maintain the baby's temperature
 - IV. Protect the eyes from light
- A. I, II and III
 - B. I, II and IV
 - C. I, III and IV
 - D. I, II, III and IV

113. Kojo who has tetralogy of Fallot usually squats when he is dyspnoeic. The squatting relives the symptoms by

- A. Allowing maximal chest expansion and oxygenation
- B. Delaying venous return to the heart
- C. Increasing the production of red blood cells
- D. Increasing venous return to the heart

114. The nurse is preparing the parents of a newborn for home phototherapy. Instructions should include which of the following?

- A. Maintain infant in prone position
- B. Dress infant in light-weight clothing
- C. Make sure eyelids are open before applying patches
- D. Remove patches during feedings and other care giving activities

115. Before introducing a tube feeding to a baby the nurse should

- A. Irrigate the tube with water
- B. Place the baby in the Trendelenburg position
- C. Provide the baby with pacifier
- D. Slowly instill 10ml of formula

116. Which of the following is an important nursing intervention in the care of a child with bacterial conjunctivitis?

- A. Warm, moist compresses to remove crusts
- B. Oral antihistamines to minimize itching
- C. Continuous warm compresses to relieve discomfort
- D. Application of optic corticosteroids to reduce inflammation

117. What clinical manifestations would suggest hydrocephalus in a neonate?

- A. Bulging fontanel, dilated scalp veins
- B. Closed fontanel, high-pitched cry
- C. Constant low-pitched cry, restlessness
- D. Depressed fontanel, decreased blood pressure

118. Which of the following is not a symptom of bacterial meningitis in a child

- A. High temperature
- B. Irritability
- C. Vomiting
- D. Low pitched cry

119. A child with tetralogy of fallot becomes acutely ill with an increase cyanosis, tachycardia and tachypnoea. Which nursing action would be most effective to relieve cardiac overload?

- A. Have oxygen equipment available
- B. Maintain suction equipment available
- C. Place child in Knee-chest position
- D. Place child in Trendelenburg position

120. The following conditions are all neurological disorder in infant EXCEPT

- A. Exomphalus
- B. Hydrocephalus
- C. Myelomeningocele
- D. Spinal bifida occulta