
1. A 30-year-old woman presents with severe bleeding from a deep leg wound. What is the first thing you should do?

- A) Apply a tourniquet above the wound
- B) Elevate the leg
- C) Apply direct pressure to the wound
- D) Clean the wound with antiseptic

Answer: C

2. A person is in shock after an accident and has rapid, weak pulse, pale skin, and cold extremities. What type of shock are they most likely experiencing?

- A) Cardiogenic shock
- B) Hypovolemic shock
- C) Neurogenic shock
- D) Septic shock

Answer: B

3. A 50-year-old man has a deep laceration to his arm. The bleeding is profuse and pulsing. What type of bleeding is this?

- A) Venous bleeding
- B) Arterial bleeding
- C) Capillary bleeding
- D) Internal bleeding

Answer: B

4. A patient is showing signs of shock. You are unable to stop the bleeding, and the pulse is faint. What is the most immediate treatment to control shock?

- A) Administer oxygen
- B) Lay the patient flat and elevate the feet
- C) Cover the patient with a warm blanket
- D) Provide the patient with fluids orally

Answer: B

5. A person has a puncture wound on their hand from stepping on a nail. What should you do first?

- A) Clean the wound with alcohol
- B) Remove the nail
- C) Apply pressure to stop bleeding
- D) Apply an antiseptic and bandage

Answer: C

6. You are treating a patient with severe hemorrhage and shock. What is the best position to place the patient in to promote circulation?

- A) Trendelenburg position (feet elevated above head)
- B) Sitting upright with legs bent
- C) Lying flat with head elevated
- D) On their side, with legs straight

Answer: A

7. A 25-year-old woman presents with severe bleeding from a wound on her thigh. You have applied direct pressure, but the bleeding continues. What should you do next?

- A) Apply a tourniquet above the wound
- B) Continue applying pressure
- C) Give the patient water to drink
- D) Use a cold compress on the wound

Answer: A

8. A patient with a large abdominal wound is bleeding profusely. What should you NOT do?

- A) Cover the wound with sterile gauze
- B) Apply direct pressure
- C) Push protruding organs back into the abdomen
- D) Place the patient in the recovery position

Answer: C

9. A 60-year-old man is suffering from shock after losing a large amount of blood. His blood pressure is low, and his skin is clammy. What should be your first step?

- A) Apply a tourniquet to the limbs
- B) Keep the patient warm and comfortable
- C) Have the patient drink fluids
- D) Give the patient adrenaline

Answer: B

10. A person is bleeding heavily from a deep wound. You are unable to stop the bleeding with direct pressure. What should you do?

- A) Use a tourniquet above the wound
- B) Elevate the limb above the head
- C) Clean the wound with water
- D) Apply ice to the wound

Answer: A

11. A person is unconscious and breathing normally but has a large laceration on their forearm. What should you do first?

- A) Apply pressure to the wound
- B) Check for signs of shock
- C) Position the person in the recovery position
- D) Clean the wound with antiseptic

Answer: C

12. A 40-year-old man is having difficulty breathing after an accident. His skin is pale, and his heart rate is fast. What is the most likely condition he is in?

- A) Cardiac arrest
- B) Hypovolemic shock
- C) Pulmonary embolism
- D) Respiratory distress

Answer: B

13. You are treating a wound that has been exposed to dirt and debris. What is the best way to clean it?

- A) Use antiseptic solution
- B) Rinse the wound with clean water
- C) Apply pressure and do not clean the wound
- D) Use hydrogen peroxide

Answer: B

14. A person has an open wound with large amounts of blood oozing out. What should you do to prevent further blood loss?

- A) Apply a tourniquet
- B) Apply direct pressure to the wound
- C) Elevate the limb above the head
- D) Put the person in a sitting position

Answer: B

15. A 28-year-old woman has a deep cut on her leg, and she is in severe pain. What should you do first?

- A) Clean the wound
- B) Apply a sterile bandage
- C) Apply pressure to stop the bleeding
- D) Take her to the hospital immediately

Answer: C

16. A 40-year-old man has been in a car accident and is exhibiting signs of shock, including weak pulse and rapid breathing. What should you do next?

- A) Lay the patient down and elevate the legs
- B) Apply a cold compress to his head
- C) Give him a drink of water
- D) Attempt to stop any bleeding

Answer: A

17. A patient is showing signs of hemorrhagic shock. What is the most critical intervention?

- A) Administer oxygen
- B) Keep the patient calm
- C) Control the bleeding
- D) Administer pain relief

Answer: C

18. A person presents with a minor burn to the hand. What is the first step in first aid?

- A) Apply a bandage to the wound
- B) Immerse the hand in cold water
- C) Use a burn cream on the hand
- D) Cover the burn with sterile gauze

Answer: B

19. A person experiences a traumatic amputation of a finger. What should you do immediately?

- A) Apply pressure to stop the bleeding
- B) Elevate the limb above the head
- C) Apply a tourniquet above the wound
- D) Wrap the amputated finger in cloth and send it with the person to the hospital

Answer: A

20. A patient presents with severe abdominal pain, vomiting, and signs of shock. What should you suspect?

- A) Internal bleeding
- B) External hemorrhage
- C) Mild dehydration
- D) Appendicitis

Answer: A

21. A person has a large scalp wound with heavy bleeding. What should you do to control the bleeding?

- A) Apply direct pressure to the wound
- B) Place a bandage over the wound
- C) Apply cold to the wound
- D) Apply a tourniquet around the neck

Answer: A

22. A 32-year-old man is in shock after a traumatic injury and is becoming confused. What is the primary cause of his confusion?

- A) Lack of oxygen to the brain
- B) Blood loss
- C) Pain from the injury
- D) The shock is progressing

Answer: B

23. You are treating a patient in shock. What is the most important thing to monitor?

- A) Temperature
- B) Respiratory rate
- C) Pulse rate and blood pressure
- D) Level of consciousness

Answer: C

24. A person has a large, deep wound on their thigh with severe bleeding. What should be applied to control the bleeding?

- A) A tourniquet above the wound
- B) A clean dressing
- C) Pressure dressing
- D) A cold pack

Answer: A

25. A person has been in an accident and is bleeding heavily. You do not have access to bandages or gauze. What should you do?

- A) Use a clean cloth or clothing to apply pressure to the wound
- B) Elevate the limb above the head
- C) Use ice to slow the bleeding
- D) Wrap the wound with tape

Answer: A

26. A person has been exposed to a chemical and presents with a burn and difficulty breathing. What is the first step in treating them?

- A) Remove them from the source of exposure
- B) Administer oxygen
- C) Apply a burn cream
- D) Call emergency services

Answer: A

27. You are treating a patient who is in shock. What is the best way to raise blood pressure?

- A) Give the patient water
- B) Keep the patient warm and comfortable
- C) Elevate the legs above the head
- D) Offer the patient food

Answer: C

28. You are treating a person who has been severely bleeding from a wound. What should you do if direct pressure is not stopping the bleeding?

- A) Apply a tourniquet above the wound
- B) Apply a cold compress to the wound
- C) Keep applying pressure without any change
- D) Elevate the affected limb

Answer: A

29. A person has a deep wound that is not bleeding heavily, but there is a lot of dirt in it. What should you do first?

- A) Clean the wound with soap and water
- B) Apply direct pressure to stop bleeding
- C) Irrigate the wound with clean water
- D) Apply a bandage and leave the wound alone

Answer: C

30. A 60-year-old woman has fallen and is showing signs of shock. What is the first thing you should do?

- A) Administer pain relief
- B) Control the bleeding if any
- C) Provide fluids to the patient
- D) Lay her down and elevate the legs

Answer: D

31. You are treating a burn. What is the first thing you should do?

- A) Apply ointment
- B) Immerse the burn in cold water
- C) Wrap the burn with a sterile bandage
- D) Cover it with a burn gel

Answer: B

32. A person has a gunshot wound and is in shock. What should you do immediately?

- A) Administer pain medication
- B) Apply a sterile dressing and pressure to control the bleeding
- C) Try to remove the bullet
- D) Keep the person standing

Answer: B

33. A person with a bleeding wound is becoming drowsy and confused. What should you do next?

- A) Monitor their breathing
- B) Apply more pressure to the wound
- C) Administer fluids orally
- D) Lay the person down and elevate their legs

Answer: D

34. A 40-year-old man has a severe head injury with visible bleeding. What is the most important thing to do?

- A) Apply a bandage to the head
- B) Ensure the person is breathing
- C) Keep the person awake
- D) Try to stop the bleeding at all costs

Answer: B

35. A person has been in an accident and is in severe shock. What is the most appropriate first aid intervention?

- A) Elevate their head
- B) Apply pressure to the wound
- C) Keep them flat and warm
- D) Give them fluids

Answer: C

36. A person has severe bleeding from an arm injury. What should you do if you can't control the bleeding?

- A) Apply a tourniquet
- B) Elevate the arm
- C) Keep applying pressure
- D) Apply a bandage and wait for help

Answer: A

37. A person is suffering from shock and has an altered mental state. What should you do?

- A) Apply a tourniquet
- B) Monitor vital signs and keep the person warm
- C) Give them fluids
- D) Apply pressure to the wound

Answer: B

38. A person has a large wound on their chest and is struggling to breathe. What is the most important action?

- A) Apply a dressing to the wound
- B) Keep the person sitting upright
- C) Call emergency services
- D) Apply pressure to stop the bleeding

Answer: C

39. You are treating a patient who has a large laceration. What is the best method to stop the bleeding?

- A) Apply pressure directly to the wound
- B) Place the wound under cold water
- C) Elevate the area
- D) Apply ointment

Answer: A

40. A person is in shock, and their pulse is weak and fast. What is your next step?

- A) Provide pain relief
- B) Administer fluids orally
- C) Lay the person flat and elevate their legs
- D) Apply direct pressure to the wound

Answer: C

41. A person has suffered a puncture wound. What should be done to treat it?

- A) Remove the object causing the wound
- B) Apply direct pressure around the wound
- C) Clean the wound with antiseptic

D) Cover the wound with a bandage

Answer: B

42. You are treating a patient in shock. What should be monitored most closely?

A) Blood pressure and pulse

B) Respiratory rate

C) Pain level

D) Level of consciousness

Answer: A

43. A person presents with severe bleeding and rapid breathing. What is your first action?

A) Elevate their feet

B) Monitor their breathing

C) Apply pressure to the wound

D) Offer them a drink of water

Answer: C

44. A 35-year-old male is unconscious and has a significant head injury. What should you do?

A) Apply pressure to the wound

B) Check for signs of shock

C) Ensure their airway is clear and monitor breathing

D) Lay the patient flat and elevate their feet

Answer: C

45. A person has suffered a gunshot wound to the chest. What should be done first?

A) Call emergency services

B) Apply a dressing to the wound

C) Monitor vital signs

D) Apply pressure to the wound

Answer: A

46. A person presents with a large abdominal wound. What is your first action?

- A) Apply direct pressure to stop the bleeding
- B) Clean the wound
- C) Keep the person warm and flat
- D) Elevate the legs

Answer: A

47. A person is unconscious, and their wound is bleeding heavily. What should be done immediately?

- A) Apply a bandage to the wound
- B) Ensure the airway is open
- C) Apply direct pressure to the wound
- D) Elevate their legs

Answer: B

48. A person presents with a deep wound on their leg with excessive bleeding. What should you do if the bleeding continues after applying pressure?

- A) Apply a tourniquet above the wound
- B) Keep applying pressure
- C) Elevate the leg above the heart
- D) Cover the wound with a sterile bandage

Answer: A

49. A person has a large contusion with swelling and pain. What should you do?

- A) Apply a cold pack to the area
- B) Apply heat to the area
- C) Elevate the affected limb
- D) Apply a compression bandage

Answer: A

50. A person is suffering from shock after a severe fall. What is the best way to prevent further complications?

- A) Keep them warm and flat
- B) Give them fluids immediately
- C) Apply ice to the wound
- D) Keep them sitting up

Answer: A
