

KAUF UNIVERSITY COLLEGE

BSC NURSING, FIRST SEMESTER EXAMINATION, 2020 / 2021

COURSE: KNUR 213, FIRST AID BANDAGING

TIME ALLOWED: ONE HOUR 20 MINUTES

INSTRUCTION: CIRCLE THE BEST ANSWER ON THE QUESTION PAPER

OBJECTIVE QUESTIONS (CIRCLE THE CORRECT ANSWER ON THE QUESTION PAPER)

1. The most common sprained site is _____
a. Ankle joint b. Ligament c. Hamstring muscle d. Lumbar spine
2. A victim is coughing up blood with bleeding from the mouth and abdomen is tender to touch. Pulse is weak and rapid. The victim is having signs of _____
a. Massive head injuries b. Internal bleeding c. Drug overdose d. Possible poisoning
3. What does 'R' in the acronym 'RICE' stand for?
a. Rest b. Rehabilitation c. Resuscitate d. Reassessment
4. An injury to the _____ in the _____ is most commonly seen in Strain?
a. Cartilage, Musculoskeletal disease b. Ankle, joint c. Hamstring muscle, lower back d. Hamstring muscle, back of the thigh
5. What is the ratio of chest compression : rescue breath for adults?
a. 30 : 3 b. 2 : 30 c. 20 : 3 d. 30 : 2
6. When prioritizing the treatment of casualties which order must you follow?
a. Breathing – Bones/Burns – Bleeding – Other conditions
b. Bleeding – Breathing – Bones/Burns – Other conditions
c. Breathing – Bleeding – Bones/Burns – Other conditions
d. Bones/Burns – Breathing – Bleeding – Other conditions
7. _____ is used to determine the percentage of area of skin involved in determining the severity of burns.
a. Wallace figure of eight b. DRABC c. Wallace rule of nine d. The burns theory
8. In Strain casualties, _____ is applied to the injured area for _____ minutes at a time to avoid _____.
a. Ice pack, 20, frostbite
b. Ice pack, 5 to 10, complications
c. Warm compress, 20, swelling
d. Heat, 10, frothy
9. Shock is the body's response to _____.
a. Blood loss b. Emotional stress c. Severe allergic reaction d. All of the above
10. The following are applied in the management of Sprain except _____

- a. Use of bandages b. Heimlich maneuver c. Rehabilitation d. Over-the-counter drugs
11. What steps would you take to control bleeding from a nosebleed?
- a. Sit casualty down, lean forward and pinch soft part of nose.
 b. Sit casualty down, lean backward and pinch soft part of nose.
 c. Lie casualty down and pinch soft part of nose.
 d. Lie casualty down and pinch top of nose.
12. A state of partial consciousness is known as _____
- a. Coma b. Stupor c. Syncope d. Coma vigil
13. Which fracture has a high risk of infection?
- a. Fissured fracture b. Closed fracture c. Open fracture d. Complicated fracture
14. What is first aid?
- a. The immediate care given to an individual who has been injured or become ill prior to the arrival of qualified medical assistance
 b. The immediate care given only to unconscious individuals prior to the arrival of qualified medical assistance
 c. The immediate care given only to ill individuals prior to the arrival of qualified medical assistance
 d. The immediate care given only to injured individuals prior to the arrival of qualified medical assistance
15. Recovery from shock can be seen when _____
- a. The patient does not vomit
 b. The skin looks clammy and cyanosed
 c. The respiration is rapid and shallow
 d. The pulse becomes slower and stronger
16. What names are given to the three different depths of burns?
- a. Small, medium and large.
 b. Primary, secondary and tertiary degree.
 c. Minor, medium and severe.
 d. Superficial, partial thickness, full thickness.
17. There are three basic steps you can take in an emergency:
- a. Call, Check, Care b. Check, Call, Care c. Recognize, Decide, Call
 d. Decide, Execute, Call
18. The DRSABCD acronym is used in the management of _____
- a. Stroke b. Poisoning c. Choking d. shock
19. The first D in the acronym DRSABCD stands for _____
- a. Danger b. Defibrillation c. Death d. Drooling
20. The last D in the acronym DRSABCD stands for _____
- a. Danger b. Defibrillation c. Death d. Drooling
21. The letter A in the acronym DRSABCD stands for _____
- a. Accident b. Ambulance c. Alert d. Airway

22. A first aider aims at _____
 a. Preserving life
 b. Preventing illness from worsening
 c. promoting healing and recovery
 d. All of the above
23. You can assess one's level of consciousness by doing the following except _____
 a. Touching b. pouring water on the casualty c. pinching d. calling out his or her name
24. A type of fracture in which one fragment of bone goes into another and is firmly wedged together is known as _____
 a. Comminuted fracture b. Impacted fracture c. Greenstick fracture d. Avulsed fracture
25. The process of sorting multiple casualties in the event of war or disaster.
 a. Trial b. Tasking c. Triage d. Disaster management
26. A casualty has cut her finger. What do you NOT do?
 a. Wash with main tap water b. Put a plaster on the wound
 c. Apply direct pressure if necessary d. Apply an antiseptic solution
27. Dressing and bandages are used to _____
 a. Reduce the victim's pain. b. Reduce internal bleeding.
 c. Help control bleeding and prevent infection. d. Make it easier to take the victim to the hospital
28. You are caring for a victim with a burned hand. Put the hand in cool water if, _____
 a. The burns are deep b. There are burns with open blisters
 c. The burns are minor with no open blisters d. All of the above
29. Arterial bleeding is _____
 a. Easy to control
 b. Dark red in colour
 c. Under pressure
 d. None of the above
30. What should you do for a casualty with a sprained ankle?
 a. Apply local heat b. Massage the swelling and secure firmly
 c. Rest with leg elevated d. Keep exercising it
31. Which of the following could indicate internal bleeding in the stomach?
 a. Bright red frothy blood coughed up b. Dark red blood vomited
 c. Smoky urine d. Black tarry motions from the rectum
32. What should you not do when you arriving at a scene?
 a. Reposition the casualty into the recovery position b. Check for pulse
 c. Clear the airway d. Sprinkle water on the casualty's face
33. A burn should be immersed or placed under cold water for at least _____
 a. 5 minutes b. 10 minutes c. 15 minutes d. 20 minutes
34. If blood sips thru a bandaged wound, _____

Indicate on the space provided if the sentence is True or False

46. _____ During burns or an injury, garments should be removed from the injured part first
47. _____ When giving first aid, the first aider could cause further injury to the victim
48. _____ The body temperature of a victim in shock usually rises
49. _____ Tourniquets are used as first trial to arrest bleeding
50. _____ Greenstick fracture is common in children under 12 years
51. _____ Antidotes are substances given to neutralize the effect of poison
52. _____ Burns are injuries that result from moist heat or liquid
53. _____ Disaster management can be pre-planned
54. _____ Protective gloves should be worn when dealing with only blood or blood products.
55. _____ CPR is not performed on children.
56. _____ Do not break blisters when a casualty has burns
57. _____ Anaphylactic shock is caused by severe allergic reactions
58. _____ A casualty with burns or haemorrhage can present with hypovolemic shock
59. _____ By-standers should never be involved in managing a scene
60. _____ Oligemic shock occurs when more blood is pumped to the heart

Match the following appropriately

- | | |
|-----------------|-------------------------|
| 61. Bleeding | Muscle pull |
| 62. Fainting | Shaking up of the brain |
| 63. Strain | Coughing of blood |
| 64. Epistaxis | Nose bleed |
| 65. Fragment | Blood in stool |
| 66. Concussion | Blood in urine |
| 67. Hemoptysis | Haemorrhage |
| 68. Hematemesis | Syncope |
| 69. Melaena | Pieces of broken bones |
| 70. Haematuria | Vomiting of blood |