

PART-I-Multiple choice questions

Instruction: Read each question carefully and provide your answer on the answer sheet provided. (1pt each)

1. The best position for the client who is admitted with risk of increased intracranial pressure from a concussion would be
 - A. Trendelenburg
 - B. Semi-fowler's.
 - C. Sim's lateral.
 - D Supine.
2. Which of the following findings would the nurse expect to see in a client diagnosed with metabolic acidosis?
 - A. Hypercalcemia
 - B. Hypermnatremia
 - C. Hyperkalemia
 - D Hypermagnesemia
3. The nurse counsels a client who has been prescribed a loop diuretic to supplement her diet with foods high in
 - A. sodium
 - B. Calcium
 - C. Magnesium
 - D potassium
4. Which of the following laboratory results would the nurse expect to observe in a client with metabolic alkalosis?
 - A. Ph 7.48; pCO₂ 43; HCO₃ 33
 - B. Ph 7.31; CO₂ 44; HCO₃ 20
 - C. Ph 7.16; CO₂ 57; HCO₃ 25
 - D Ph 7.18; CO₂ 41; HCO₃ 14

A client is admitted to the
Valium. Arterial blood gas
HCO₃ of 25. The nurse
A. Metabolic acidosis
B. Respiratory acidosis
C. Respiratory alkalosis
D Metabolic alkalosis

6. W
Mani

- er sheet
pressure
5. A client is admitted to the emergent department following ingestion of five tablets of Valium. Arterial blood gas analysis reveals a pH of 7.13; PO₂ of 80; a pCO₂ of 50; and an HCO₃ of 25. The nurse would interpret the results as
- A. Metabolic acidosis
 - B. Respiratory alkalosis
 - C. Respiratory acidosis.
 - D Metabolic alkalosis

6. When caring for a client with a tunneled central venous catheter, which of the following Manifestations requires immediate intervention by the nurse?

- A. Redness at the catheter site
- B. Tenderness along the track of the catheter
- C. A loose dressing
- D A small amount of blood in the dressing

7. A client with multiple IV sites is to begin receiving total parental nutrition (TPN). most appropriate IV access site would be the peripherally inserted central venous (PICC) catheter in the

- A. Femoral vein.
- B. Cephalic vein.
- C. Subclavian vein.
- D Basilic vein.

8. What is the purpose of valves in the vein?

- A. To redirect blood flow to another vein
- B. To shut off blood flow to stop bleeding
- C. To prevent backflow of blood as it travels through the body
- D To help blood flow more forcefully through the body

9. What is the best choice of catheter to use in a patient with a history of thrombophlebitis?
- A. The largest one you feel you can successfully insert in the patient
 - B. The smallest one you can find
 - C. The largest gauge your facility has
 - D. The smallest gauge to accommodate therapy

10. What is the first step in preparing the IV site?

- A. Wash your hands
- B. Put on gloves
- C. Clean the site with an antiseptic
- D. Apply a tourniquet

11. Which of the following is causes of phlebitis; except?

- A. Using needle too small
- B. Prolonged use of the same IV sites
- C. Injury during venipuncture
- D. Using needle too large

12. Which of the following is systemic complication of intravenous therapy?

- A. Phlebitis
- B. Infiltration
- C. Embolism
- D. Thrombosis

13. Which of the following is True about total amount of fluid present in human body?

- A. 30%
- B. 40%
- C. 60%
- D. 80%

14. Which of the following is not a sign of fluid overload?

- A. 120-125 ME/L
- B. 135-145 ME/L
- C. 150-160 ME/L
- D. 170-180 ME/L

15. Which of the following is not a sign of fluid overload?

- A. Decrease in urine output
- B. Edema
- C. Uncoordinated
- D. Fever

16. Which of the following is not a sign of fluid overload?

- A. Decrease in urine output
- B. Edema
- C. Uncoordinated
- D. Fever

14. Which of the following is normal sodium level?
- A. 120-125 ME/L
 - B. 135-145 ME/L
 - C. 150-160 ME/L
 - D. 170-180 ME/L
15. Which of the following is clinical manifestation of severe dehydration?
- A. Decrease skin turgor
 - B. Edema
 - C. Unconsciousness
 - D. Fever
16. Which of the following is normal potassium level?
- A. 1-2 mEq/L
 - B. 3-5 mEq/L
 - C. 6-7 mEq/L
 - D. 8-9 mEq/L
17. Which of the following is how much zinc supplement providing to dehydration patient?
- A. 10-20 mg
 - B. 30-40 mg
 - C. 40-50 mg
 - D. 60-70 mg
18. Which of the following treatment is providing severe dehydration?
- A. Oral water supplement
 - B. Providing ORS
 - C. Bed rest
 - D. Infuse isotonic crystalloid solution
19. Which of the following is contraindication to administration of oral dehydration therapy?
- A. Intestinal obstruction
 - B. Fever
 - C. Diarrhoea
 - D. Dysentery
20. Which of the following is paO_2 level in moderate hypoxia?
- A. 95-100
 - B. 60-80
 - C. 40-60
 - D. 20-30
21. Which of the following is normal sodium bicarbonate level?
- A. 10 to 20 mEq/L
 - B. 22 to 32 mEq/L

- C. 30 to 40 mEq/L
- D. 40 to 50 mEq/L

22. Which of the following condition paO_2 level more than 45 mm Hg?

- A. Respiratory Acidosis
- B. Respiratory Alkalosis
- C. Metabolic Acidosis
- D. Metabolic Alkalosis

23. Which of the following is student is nervous for examination and is breathing rapidly, what do you expect out of the following

- A. Metabolic acidosis
- B. Metabolic alkalosis
- C. Respiratory acidosis
- D. Respiratory alkalosis

24. Which of the following is cause of hypervolemia?

- A. Early renal failure
- B. Diarrhoea
- C. Vomiting
- D. Respiratory infection

25. Which of the following is clinical manifestation of hypovolemia?

- A. High BP
- B. Weight gain
- C. Oliguria
- D. Increased skin turgor

26. Which of the following is nursing management of hypovolemia patient, except

- A. Monitor daily weight
- B. Monitor vital signs
- C. Monitor urine output
- D. Observe for temperature

27. Which of the following is drug used to manage hypervolemia?

- A. Furosemide
- B. Diazepam
- C. Amoxicillin

28. Which of the following is **False** about controlling hypervolemia?

- A. Restricting sodium intake
- B. Monitoring parenteral fluid intake
- C. Supine position providing to patient
- D. Maintaining intake and output chart

29. Which of the following is normal calcium level?

- A. 1.5 to 2.5 mg/dl
- B. 3.5 to 5.5 mg/dl
- C. 6.5 to 7.5 mg/dl
- D. 8.5 to 10.5 mg/dl

30. Which of the following is normal magnesium level?

- A. 1.5 to 2.5 mEq/L
- B. 3.5 to 4.5 mEq/L
- C. 4.5 to 5.5 mEq/L
- D. 5.5 to 6.5 mEq/L

31. Which of the following is function of potassium?

- A. Maintaining acid base balance
- B. Transmission of nerve impulses to kidney
- C. Increases acidosis
- D. Increases heart impulses

32. Which of the following is clinical manifestation of hypokalemia?

- A. Breathing difficult
- B. Tachycardia
- C. Confusion
- D. Diarrhoea

33. A solution that is similar to the osmolarity of plasma that will cause no cell damage called?

- A. Hypotonic
- B. Normal saline
- C. Hypertonic
- D. Isotonic

34. All the following are important electrolyte in the body except:

- A. Potassium ions
- B. Carbon ions
- C. Chloride ions
- D. Sodium ions

35. Which of the following is intravenous fluid administered to correct hyponatremia?

- A. Dextrose
- B. Ringer lactate
- C. 0.9% sodium chloride
- D. 5% dextrose

36. Which of the following is causes of metabolic alkalosis, except?

- A. Vomiting
- B. Diarrhoea
- C. Loop diuretics
- D. Corticosteroid

37. Which of the following is nursing management of hyperchloremia patient?

- A. Monitoring vital signs of patient
- B. Provide bed rest
- C. Provide fowler position
- D. Provide Massage therapy

38. Which of the following therapy produces subconscious state to patient?

- A. Thermotherapy
- B. Laughter therapy
- C. Hypnosis therapy
- D. Manipulation therapy

39. Which of the following is sign of heat stroke, but not heat exhaustion?

- A. Fat heart beat
- B. Coma
- C. Dizziness
- D. Continuous heavy sweating

40. Urine analysis is also commonly referred to as:

- A. Urine drip
- B. Urine tap

C. Urine dip
D. Urine tap

41. Which of the following is nursing management of hypernatremia?

- A. Provide bed rest
- B. Encourage output
- C. Provide medication
- D. Encourage fluid intake

42. Which of the following is nursing management of hyperkalemia?

- A. Monitor vital signs
- B. Provide bed rest

- C. Urine slip
- D. Urine tip

41. Which of the following is nursing management of hypercalcemia patient?

- A. Provide bed rest
- B. Encourage output
- C. Provide medication
- D. Encourage fluid intake

42. Which of the following condition identified hypophosphatemia?

- A. Malnutrition
- B. Anemia
- C. Hyperthyroidism
- D. Measles

43. Which of the following hormone is high during early period of sleep?

- A. Adrenocorticotrophic hormone
- B. Testosterone hormone
- C. Prolactin hormone
- D. Thyroid hormone

44. Which of the following is a function of sleep?

- A. Necessary for hormone decreases
- B. Necessary for chemical
- C. Necessary for Heart
- D. Necessary for protein synthesis

45. Which of the following stages of Non random eye movement dreaming occur?

- A. Stage-I
- B. Stage-II
- C. Stage-III
- D. Stage-IV

46. Which of the following nursing intervention helps for sleeping to patient?

- A. Provide calm and quiet environment
- B. Provide light wave
- C. Make noise
- D. Provide drinks

47. Which of the following drug providing to patient not getting sleep?

- A. Antibiotic drug
- B. Vitamin drug
- C. Sedative drug
- D. Amoxicillin drug

48. Which of the following is wet dream sleep disorders?

- A. Bruxism
- B. Nocturnal erections
- C. Nocturnal enuresis
- D. Somnambulism

49. Which of the following is initial insomnia?

- A. Difficulty falling asleep
- B. Difficulty prolonged asleep
- C. Sleeping late
- D. Falling excess sleep

50. Which of the following is normal Hemoglobin level?

- A. 8-10 g/dl
- B. 13-18 g/dl
- C. 19-20 g/dl
- D. 20-25 g/dl

51. Which of the following is not contributing factor for hyponatremia?

- A. Low salt diet
- B. Renal disease
- C. Diuretics
- D. Infection

52. Which of the following is normal Red blood cell count?

- A. 1.5 to 3.5 mc/L
- B. 4.5 to 5.5 mc/L
- C. 6.5 to 7.5 mc/L
- D. 8.5 to 9.5 mc/L

53. Which of the following is ECG changes in hyper magnesium patient

- A. Prolonged PR interval
- B. Prolonged PQ wave
- C. Depressed T wave
- D. Tall QRS wave

- C. Avoid harm
- D. Increase anxiety

61. Which of the following is first technique of crisis intervention?

- A. Catharsis
- B. Clarification
- C. Suggestion
- D. Reinforcement

62. Which of the following is third step of crisis intervention?

- A. Assessment
- B. Planning
- C. Evaluation
- D. Implementation

63. Which of the following intravenous fluid not administration to the diabetic patient?

- A. Dextrose
- B. Normal saline
- C. 5% Normal saline
- D. Ringer lactate

64. Which of the following age group Physical maturity identified?

- A. School age
- B. Adolescent age
- C. Old age
- D. Adult age

65. Which of the following psychological identity observed during adult age?

- A. Personal identity
- B. Concern over next generation
- C. Changes in family situation
- D. Changes in financial situation

66. How much percentage above tie the tourniquet during intravenous cannula?

- A. 10 centimeter
- B. 15 centimeter
- C. 20 centimeter
- D. 25 centimeter

67. Which of the following is intravenous site for neonatal child?

- A. Hand

- B. Wrist
- C. Umbilical
- D. Scalp

68. Which of

- A. P

- B. Wrist
- C. Umbilical
- ☒ D. Scalp

68. Which of the following vein used for central venous catheter line?

- A. Antecubital Vein
- B. Peripheral vein
- ☒ C. Umbilical vein
- D. Jugular Vein

69. Intravenous fluid infusion of 1000ML, D5 to be infused for 24 hours, how many ml patient receive in 24 hours?

- A. 60 ML
- B. 41 ML
- ☒ C. 50 ML
- D. 71 ML

70. Which of the following is a symptom of metabolic acidosis?

- ☒ A. Dyspnea
- B. Infection
- C. Abdominal cramp
- D. Palpitation

71. Which one of the following is purpose of position?

- A. To decrease circulation
- B. To promote bed sore
- ☒ C. To improve circulation
- D. To provide discomfort

72. Which of the following is not a principle of position?

- A. To maintain good body mechanism
- B. To obtain assistance required
- ☒ C. To improve circulation
- D. To provide discomfort

73. Which of the following is position providing difficulty breathing patient?

- A. Supine position
- ☒ B. Fowler position
- C. Prone position
- D. Lithotomy position

74. Which of the following is use of prone position?
- A. Examination of anterior trunk
 - B. Examination of abdomen
 - C. Examination of spine
 - D. Examination of face
75. Which of the following is position providing to patient during enema insertion?
- A. Supine position
 - B. Lateral position
 - C. Knee chest position
 - D. Prone position
76. Which of the following is use of Sims position?
- A. Examination of heart
 - B. Examination of spine
 - C. Examination of rectum
 - D. Examination of abdomen
77. The nurse should position a client recovering from general anesthesia in?
- A. Side lying position
 - B. High fowler position
 - C. Supine position
 - D. Trendelenburg position
78. Which of the following position providing to patient during delivery?
- A. Knee chest position
 - B. Lateral position
 - C. Examination of rectum
 - D. Lithotomy position
79. A patient is scheduled for NG feeding, in which position should the nurse place
- A. Low fowlers position
 - B. Supine position
 - C. Side lying position
 - D. Trendelenburg position
80. Which of the following is principles of rehabilitation?
- A. Restore patient independence
 - B. Not active participation

- C. Demotivate patient
 - D. Patient dependent some one
81. Which of the following is rehabilitation helps to get job for person?
- A. Medical rehabilitation
 - B. Social rehabilitation
 - C. Vocational rehabilitation
 - D. Psychological rehabilitation
82. Which of the following is Not a role of rehabilitation nurse?
- A. Nursing skill
 - B. Therapeutic practice
 - C. Advice
 - D. Noncounselor
83. Which of the following is rehabilitation providing to psychiatric patient?
- A. Psychological rehabilitation
 - B. Speech rehabilitation
 - C. Physical rehabilitation
 - D. Occupational rehabilitation
84. Which of the following is normal sodium bicarbonate level?
- A. 12 to 22 mEq/L
 - B. 22 to 32 mEq/L
 - C. 32 to 42 mEq/L
 - D. 42 to 52 mEq/L
85. Which of the following is monitoring the patient receiving intravenous fluid?
- A. Heart rate
 - B. Mental status
 - C. Urine output
 - D. Capillary refill
86. Causes of metabolic alkalosis include all the following, except?
- A. Corticosteroid therapy
 - B. Hypokalemia
 - C. Diuretic therapy
 - D. Recurrent vomiting

87. Excessive citrate in transfused blood can cause which of the following abnormalities?

- A. Respiratory alkalosis
- B. Metabolic acidosis
- C. Metabolic alkalosis
- D. Respiratory acidosis

88. A needle insertion is required when accessing which of the following central venous devices (CVAD)?

- A. Non-tunneled device
- B. Peripherally inserted central catheter
- C. Implantable port
- D. Skin tunneled device

89. How long is the intended term of use for a PICC?

- A. Short
- B. Long
- C. Intermediate
- D. Very long

90. What is the biggest risk factor for central venous access devices (CVADs)?

- A. Anesthetic
- B. Infection
- C. Drug error
- D. Patency

91. In a normal ABG, a normal pH value is:

- A. 7.35 to 7.45
- B. 7.55 to 7.66
- C. 7.77 to 7.88
- D. 7.89 to 8.45

92. Which of the following regulates the secretion of erythropoietin hormone?

- A. Oxygen
- B. Proteins
- C. Number of red blood cells
- D. The concentration of hemoglobin

99. Which of the following is situational crisis?

- A. Flood
- B. Accident
- C. Behavioral changes
- D. Stress

100. Which of the following is adventitious crisis?

- A. Tension
- B. Sudden death
- C. Divorce
- D. War