

1. The best position for the client who is admitted with risk of increased intracranial from a concussion would be

A. Semi-Fowler's

B. Sims' lateral

C. Trendelenburg

D. Supine

2. Which of the following findings would the nurse expect to see in a client diagnosed with metabolic acidosis

A. Hypercalcemia

B. Hyponatremia

C. Hyperkalemia

D. Hypermagnesemia

1. The nurse counsels a client who has been prescribed a loop diuretic to supplement her diet with foods high in

A. Sodium

B. Calcium

C. Magnesium

D. potassium

2. Which of the following laboratory results would the nurse expect to observe in a client with metabolic alkalosis?

A. Ph 7.48; pCO₂ 43; HCO₃ 33

B. Ph 7.31; CO₂ 44; HCO₃ 20

C. Ph 7.16; CO₂ 57; HCO₃ 25

D Ph 7.18; CO₂ 41; HCO₃ 14

3. A client is admitted to the emergency department following ingestion of five tablets of Valium. Arterial blood gas analysis reveals a pH of 7.13, PO₂ of 80, a pCO₂ of 50, and HCO₃ of 25. The nurse would interpret the results as

A. Metabolic acidosis

B. Respiratory alkalosis

C. Respiratory acidosis.

D Metabolic alkalosis

4. When caring for a client with a tunneled central venous catheter, which of the following Manifestations requires immediate intervention by the nurse?

- A. Redness at the catheter site
- B. Tenderness along the track of the catheter
- C. A loose dressing

D A small amount of blood in the dressing

5. A client with multiple IV sites is to begin receiving total parental nutrition (TPN). Most appropriate IV access site would be the peripherally inserted central venous (catheter in the

- A. Femoral vein.
- B. Cephalic vein.
- C. Subclavian vein.

D Basilic vein.

6. What is the purpose of valves in the vein?

- A. To redirect blood flow to another vein
- B. To shut of blood flow to stop bleeding

C. To prevent backflow of blood as it travels through the body

D To help blood flow more forcefully through the body

10. What is the first step in preparing the TV site?

A. Wash your hands

B. Pat on gloves

C. Clean the site with an antiseptic

D Apply a tourniquet

11. Which of the following is causes of phlebitis, except?

A. Using needle too small

B. Prolonged use of the same IV sites

C. Injury during venipuncture

D Using needle too large

12. Which of the following is systemic complication of intravenous therapy?

A. Phlebitis

B. Infiltration

C. Embolism

D Thrombosis

13. Which of the following is True about total amount of fluid present in human body?

A. 30%

B. 40%

C. 60%

D. 80%

14. Which of the following is normal sodium level?

A. 120-125 MEL

18. 135-145 ME/l

B. 150-160 MEL

C. 170-180 ME/l

15. Which of the following is clinical manifestation of severe dub

A Decrease skin turgor

B. Edema

D. Unconsciousness

E. Fever

16. Which of the following is normal potassium level?

A. 1-2 mEq/L

B. 3-5 mEq/L

C. 6-7 mEq/L

D. 8-9 mEq/L

17. Which of the following is how much zinc supplement providing to dehydration patient?

A. 10-20 mg

B. 30-40 mg

C. 40-50 mg

D. 60-70 mg

18. Which of the following treatment is providing severe dehydration?

A. Oral water supplement

B. Providing ORS

C. Bed rest

D. Infuse isotonic crystalloid solution

19. Which of the following is contraindication to administration of oral dehydration the

A. Intestinal obstruction

B. Fever

C. Diarrhoea

D. Dysentery

20. Which of the following is paO_2 level in moderate hypoxia?

A. 95-100

B. 60-80

C. 40-60

D. 20-30

21. Which of the following is normal sodium bicarbonate level?

A. 10 to 20 mEq/L

B. 22 to 32 mEq/L

39. Which of the following is sign of heat stroke, but not heat exhaustion?

A. Fat heart beat

B. Coma

C. Dizziness

D. Continuous heavy sweating

40. Urine analysis is also commonly referred to as:

A. Urine drip

B. Urine tap

C. Urine slip

D. Urine tip

41. Which of the following is nursing management of hypercalcemia patient

A Provide bed rest

C. Encourage output

D. Provide medication

E. Encourage fluid intake

42. Which of the following condition identified hypophosphatemia?

A. Malnutrition

B. Anemia

C. Hyperthyroidism

D. Measles

43. Which of the following hormone is high during early period of sleep?

A. Adrenocorticotrophic hormone

B. Testosterone hormone

C. Prolactin hormone

D. Thyroid hormone

44. Which of the following is a function of sleep?

A. Necessary for hormone decreases

B. Necessary for chemical

C. Necessary for Heart

D. Necessary for protein synthesis

45. Which of the following stages of Non random eye movement dreaming occur?

- A. Stage-1
- B. Stage-II
- C. Stage-III
- D. Stage-IV

46. Which of the following nursing intervention helps for sleeping to patient?

- A. Provide calm and quiet environment
- B. Provide light wave
- C. Make noise
- D. Provide drinks

47. Which of the following drug providing to patient not getting sleep?

48 発生 of the dream lep disorders?

- A. Bro

B. Nocturnal

C. Somnambulism

49. Which of the following is initial is

A. Difficulty falling deep

Difficulty prolonged asleep

B. Sleeping late

C. Falling excess sleep

50. Which of the following is normal Hemoglobin level?

A. 8-10 g/dl

B. 13-18 g/dl

C. 19-20 g/dl

D. 20-25 g/dl

51. Which of the following is not contributing factor for hyponatremia?

- A. Low salt diet
- B. Renal disease
- C. Diuretics
- D. Infection

52. Which of the following is normal Red blood cell count?

- A. 1.5 to 3.5 mc/L
- B. 4.5 to 5.5 mc/L
- C. 6.5 to 7.5 mc/L
- D. 8.5 to 9.5 mc/L

53. Which of the following is ECG changes in hyper magnesium patient

- A. Prolonged PR interval
- B. Prolonged PQ wave

C. Depressed T wave

D. Tall QRS wave

All the following are physical signs of physical child abuse EXCEPT.....

A. Bruises

B. Being sad

C. Cuts

D. Untreated wounds

54. Which of the following is not a physical abuse?

A. Beating

B. Corporal punishment

C. Child labour

D. Punching

55. Yaw Mensah was sent at 12:00 noon to buy salt for lunch, on his way he saw his friends playing football and he decided to join them. He reported with the salt at 6:30pm. His mother confined him in a dark room for two hours. Identify the type of abuse his mother subjected him to.

A. Commercial abuse

B. Emotional abuse

C. Neglect

D. Physical abuse

56. All the following are reasons for follow up EXCEPT.....

A. Attending to areas of developmental need

B. Making sure the perpetrator is prosecuted

C. Monitoring of child's overall progress

D. Performing further medical examination as appropriate

57. The wide range of specimens that could be collected to assist the criminal investigations process is called.....specimen

A. Blood product

B. Body tissues

C. Forensic

D. Histological

61. Which of the following is first technique of crisis intervention?

A. Catheterization

B. Clarification

C. Suggestion

D. Reinforcement

62. Which of the following is third step of crisis intervention?

A. Assessment

B. Planning

C. Evaluation

D. Implementation

63. Which of the following intravenous Fluid not administration to the diabetic patient?

A. Dextrose

B. Normal saline

C. 5% Normal saline

D. Ringer lactate

64. Which of the following age group Physical maturity identified?

A. School age

B. Adolescent age

C. Old age

D. Adult age

65. Which of the following psychological identity observed during adult age?

- A. Personal identity
- B. Concern over next generation
- C. Changes in family situation
- D. Changes in financial situation

66. How much percentage above tie the tourniquet during intravenous cannula?

- A. 10 centimeter
- B. 15 centimeter
- C. 20 centimeter
- D. 25 centimeter

67. Which of the following is intravenous site for neonatal child?

- A. Hand

74. Which of the following is use of prone position?

- A. Examination of anterior trunk
- B. Examination of abdomen
- C. Examination of spine
- D. Examination of face

75. Which of the following is position providing to patient during enema insertion?

- A. Supine position
- B. Lateral position
- C. Knee chest position
- D. Prone position

76. Which of the following is use of Sims position?

- A. Examination of heart
- B. Examination of spine

C. Examination of rectum

D. Examination of abdomen

77. The nurse should position a client recovering from general anesthesia in?

A. Side lying position

B. High fowler position

C. Supine position

D. Trendelenburg position

78. Which of the following position providing to patient during delivery?

A. Knee chest position

B. Lateral position

C. Examination of rectum

D. Lithotomy position

79. A patient is scheduled for NG feeding, in which position should the nurse place

1. Low fowlers position
2. Supine position
3. Side lying position
4. Trendelenberg position

90. What is the biggest risk factor for central venous access devices (CVADs)?

- A. Anesthetic
- B. Infection
- C. Drug error
- D. Patency

91. In a normal ABG, a normal pH value is.

- A. 7.35 to 7.45
- B. 7.55 to 7.66

C. 7.77 to 7.88

D. 7.89 to 8.45

92. Which of the following regulates the secretion of erythropoietin

A. Oxygen

B. Proteins

C. Number of red blood cells

D. The concentration of hemoglobin