
a
Each point: one donor. Low = the donor's cell-type pattern looks the same in both datasets; high = the two datasets disagree
Green x Mathys
Green x Multiome
Mathys x Multiome
Multiome x PsychAD
ainSCOPE CMC x PsychA
n=227
rho=-0.28, p=2e-05
n=105
rho=-0.08, p=0.39
n=97
rho=+0.21, p=0.041
n=76
rho=+0.17, p=0.13
n=54
rho=+0.03, p=0.82
Donor-level cross-dataset disagreement
(mean |CLR difference|, all shared cell types)
1.6
1.2
0.8
-4
-2
0
2
log2(median UMI, dataset B / dataset A)
b
Green x Mathys
n=227 beta=+0.095 [+0.048, +0.142] p=9e-05
Green x Multiome
n=105 beta=+0.035 [-0.039, +0.108] p=0.35
Mathys x Multiome
n=97 beta=+0.000 [-0.095, +0.095] p=1
Multiome x PsychAD
n=76 beta=+0.089 [-0.012, +0.189] p=0.082
BrainSCOPE CMC x PsychAD
n=54 beta=-0.020 [-0.091, +0.050] p=0.57
Pooled (fixed-effects)
n=559 beta=+0.053 [+0.021, +0.085] p=0.0013
-0.1
0.0
0.1
0.2
Slope of discordance on log2(|depth gap| + 1)