

a
ρ = 0.71, p = 4e-24
n = 150 supertypes
Deconvolution accuracy of the supertype
(median out-of-fold r, bulk vs snRNA-seq)
0.00
0.25
0.50
0.0
0.2
0.4
0.6
snRNA-seq reliability of the supertype
(median ICC across matched-donor pairs)
Neuronal
Non-neuronal
Reactive (-SEAAD)
b
Sst Chodl
Lamp5 Lhx6
OPC
Astrocyte
Sst
Lamp5
L6 IT Car3
Pvalb
L5/6 NP
Sncg
Immune
L6b
Chandelier
Oligodendrocyte
L6 CT
L4 IT
L2/3 IT
L6 IT
Endothelial
VLMC & Perivascular
Vip
Pax6
L5 IT
L5 ET
-0.5
0.0
0.5
1.0
1.5
Triangulation reliability
(ROSMAP triad, bulk leg; axis clipped)
unresolved supertype
resolved supertype
c
CommonMind × PsychAD MSBB
HBCC × PsychAD HBCC
MSBB × PsychAD MSBB
ROSMAP × Green
ROSMAP × Mathys
ROSMAP × Multiome
Out-of-fold r, deconvolved vs snRNA-seq
0.75
0.50
0.25
0.00
-0.25
Subclasses
Other supertypes
CORE supertypes