

# Supplementary material (Online Resource 1): the derivations

to the article “Universe24: an integer cellular automaton read through clicks, the formulas it shares with nature and the ones it adds”

Alon Gonen 

Independent researcher, Haifa, Israel.

Corresponding author(s). E-mail(s): [alon@defounder.ai](mailto:alon@defounder.ai);

Journal: Foundations of Physics

This supplement gives the algebraic steps of every result the paper marks DERIVED or COMPUTED without a proof in its text, Derivations S.1 to S.56, numbered in their own order, each naming the paper’s section it serves; the paper’s claims list stands first, every claim with its status, its fence, what it rests on and its derivation, then the Nomenclature. The derivations keep the words of the long version of the paper, mapped to the paper’s in the glossary below; the long version [1] carries sixty-five derivations in its own numbering, and a derivation this supplement does not carry is cited as “S. $n$  of the long version” by that number. Theorems 1 to 3 of the paper are proved in S.45 and Theorem 4 in S.5 and S.6; the one-line theorems carry their proof in their sentence. Each derivation names its inputs from the paper, its steps, and, where a number was computed, the computation and its check. The notation is the paper’s: Rule3’s coefficients at a Node of clock pace  $p_0$  and Link paces  $p_a$  are  $R_a = 2 \text{ num } p_a^2$  on the two arrivals of the axis  $a$  where the axis’s two Links share one pace (the six Port coefficients  $R_{ij}$  of Eq. (1) of the paper in general),  $S = 12 \text{ den } \Gamma^2 - 12(\text{den} - \text{num}) p_0^2 - 4 \text{ num } \sum_a p_a^2$  on the Node’s own level, and the wall  $w = 6 \text{ den } \Gamma^2$ ; the content read at the Node is  $x = c/\Gamma$ , also written  $U$ ;  $g = (\text{den} - \text{num})/\text{den} = 1 - \cos \omega_0$ . The lattice’s own order: every derivation below is exact algebra of the line where it says so; where a continuum method is used on the lattice (the far field  $3/(4\pi r)$  of a holder’s rest, S.25; the retarded solution of the wave equation, S.40; the shell theorem, S.29; the ray integrals along the straight path, S.20 and S.30; the radial grid, S.52), it holds at long wavelength and large  $r$  with the lattice’s correction of relative order  $(\text{Link}/r)^2$  or  $k^2/12$  along an axis, the order the law’s lines name; and the theorems of mathematics cited by name (the discrete Fourier transform’s resolution  $2\pi/n_w$  in S.8, Watson’s constant

$G_0(0) = 0.252731$  [2] in S.36, the post-Newtonian periapsis formula  $6\pi U_1/(1 - e^2)$  in S.19 [3]) are standard results used without proof.

### ***The use of a large language model.***

The derivations of this supplement, like the article, were worked out with a large language model (Claude, Anthropic, through Claude Code, 2026) used as a tool under the author’s direction; the model is not an author, and the author reviewed every step and takes full responsibility for the content, as the article’s Statements and Declarations state.

## **The words: the long version’s and the paper’s**

The derivations keep the words of the long version of the paper; the paper’s word stands beside each of them here.

*family* a field, one kind of physical value with its pair and its shape (the paper’s Sections 2 and 4.1)

*record* a configuration, a field’s levels over the lattice

*holder* a potential field, a field that holds a value for others to read: the sign holder is the charge potential, the massless holder gravity’s potential field, the binding holder the binding potential, a gapped holder a potential field with a gap

*content* the potential level, what a holder holds at a Node,  $x = c/\Gamma$ , also written  $U$

*share* the quantum share, a configuration’s count in a region

*credit, the credit rule* the counting rule, which credits one quantum to one NodeDetector at a click

*books* the count register of the click layer, one integer per configuration

*draw* the NodeDetector’s random draw

*window* the counting window

*lay, laid* the initial data, the configurations’ initial writing

*root* the source Node of a configuration’s initial data

*blind, the blind row* the prediction written before the computation

*front* a region’s boundary, the Ports through which its inflow is read; light’s front and the erasing front keep their names

*tension* the stress, the momentum flux of a wave

*body* a bound state, bound or free, a configuration standing in a potential well or falling free

*the taker* the configuration whose transition the click reads

*the wall* the integer bound of a level,  $6\Gamma^3$  at the lattice’s one denominator

*the pixel* the compact branch of the bound state, the width of one Node

*the hold* the potential’s write, the long version’s name of the act

*the event lattice* the lattice

*packet* a wave packet

*the declared universe* the declaration of a world: its fields with their pairs and shapes, its holders, its NodeDetectors and its initial data

## **The claims of the paper**

Every claim of the paper, with its status (the key of Section 1.3 of the paper), its fence, what it rests on and its derivation; the paper carries the same status in the parenthesis that closes each claim’s sentence. A calibration is a number taken from nature; experiment, a number the model does not fix.

**Table 1:** The claims of the paper, each with its status, its fence, what it rests on and its derivation or section

| The claim                                                                                                                                                                                                                                                                                                                                                      | Status                                                                                                                                                                                       | Fence                               | Rests on                                                                                       | Where                                         |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|------------------------------------------------------------------------------------------------|-----------------------------------------------|
| Universe24 is the meeting of the past with the future; a click is that meeting, the one measurement; only clicks are seen                                                                                                                                                                                                                                      | assumption, the paper's first statement                                                                                                                                                      | clicks                              | the model's statement                                                                          | Section 1.1                                   |
| Rule3's step is reversible exactly, integer for integer; the whole evolution goes back under the condition on the reads                                                                                                                                                                                                                                        | theorem for the step and where the writes change no level; the write an assumption                                                                                                           | lattice                             | Rule3                                                                                          | Section 3.1; S.45                             |
| The two slits' run forward and back returns every level over 40 intervals, before the counting window opens at 45                                                                                                                                                                                                                                              | theorem                                                                                                                                                                                      | lattice                             | Rule3's integer line on the declared universe                                                  | Section 3.1; S.23                             |
| Rule3 is covariant under the cube's group of 48; $D$ a scalar, $F$ a vector, $W$ a scalar under the 48, odd under the time reversal and the plane's reflection; no boost in the group; The rule's acts form no product of levels, on the infinite lattice, for one field at fixed coefficients; Three exact massive rotations, $[1, 2]$ , $[0, 1]$ , $[-1, 2]$ | theorem                                                                                                                                                                                      | lattice                             | Rule3                                                                                          | Sections 2.1, 3.2 and 3.6; S.4, S.45          |
| A conserved form at static paces with the weights $1/p_i^2$ , its current with the Link's factor, and the momentum flux as the wave's stress                                                                                                                                                                                                                   | theorem at static paces, weighted; derived for the momentum flux                                                                                                                             | lattice                             | Rule3                                                                                          | Section 3.3; S.5, S.7, S.32, S.45             |
| The count is the configuration's quantum share; the rational total is non-negative under the two conditions, the floors' total within the number of Nodes of it                                                                                                                                                                                                | theorem under the two conditions                                                                                                                                                             | lattice                             | Rule3                                                                                          | Section 3.4; S.5, S.6                         |
| The Wronskian is conserved exactly in the rational line under the condition on the paces; the integer step adds the remainders' term; antimatter is the opposite sense                                                                                                                                                                                         | theorem for the conservation; derived for the opposite sense, the sign declared                                                                                                              | lattice; clicks for antimat-<br>ter | Rule3, the rotation                                                                            | Section 3.5; S.9, S.33                        |
| An event spreads by Rule3 at one Link per interval, the causal bound; light's speed $1/\sqrt{3}$ Link per interval, its dispersion, none along the diagonal; the two fronts, the octahedron and the inscribed sphere; the lattice Klein-Gordon equation with Schrödinger's slow limit; $E = m^* c_m^2 = \omega_0$                                              | postulate 4, an assumption, the step's reach (the line reads the six Ports and nothing farther); derived for the speed, the fronts and the rest, $E = m^* c_m^2$ by construction and no test | lattice; clicks for $E$             | Rule3 for the pairs                                                                            | Sections 2.2 and 2.5.1; S.1, S.18, S.21, S.53 |
| The paces compose, $p_0 = \Gamma(1 - 1/\Gamma)^c$ ; the Link clocked twice by its reading Node's clock, $hN = 1$ ; the odd Link part, gravity's holder's three odd lines sourced by the count's flux and read at the Port as the phase $\theta_{ij} = 4V_{ij}\omega_0$                                                                                         | assumption, fitted; the composition derived from it; assumption for a moving source, the coefficient derived under it                                                                        | lattice                             | fixed by the bending's measurement, the perihelion choosing the reading and testing the choice | Section 2.3; S.34; S.56                       |
| Frame dragging's form $\delta\omega/\omega_0 = +4\mathbf{V} \cdot \mathbf{v}$ ; $\alpha_1 = \alpha_2 = 0$ ; $\eta_N = 0$ at nature's gap                                                                                                                                                                                                                       | derived under the one assumption, to first order in $\mathbf{V}$ , slow matter                                                                                                               | clicks                              | the one assumption for a moving source; S.56                                                   | Section 5.1; S.56 (f), (g)                    |

| The claim                                                                                                                                                                                                                               | Status                                                                                                                                                                                                                                                    | Fence                                                                                | Rests on                                     | Where                                                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|----------------------------------------------|----------------------------------------------------------|
| Light undragged by a rotating mass; a fast record's coupling falling as $1/\gamma_L$                                                                                                                                                    | derived under the one assumption; untested                                                                                                                                                                                                                | lattice                                                                              | the one assumption for a moving source; S.56 | Section 5.1; S.56 (h)                                    |
| The exponential metric, $\gamma = \beta = 1$ in $U$ ; the bending $4U_b$ , Shapiro's delay, the perihelion, the redshift's factor, the shadow's $2em$ ; $\gamma_K$ , $\beta_K$ and $\alpha_K$ from the pair alone, closing with the gap | derived under the one assumption; the bending's measurement its calibration, the perihelion the test of the end's choice, the rest its consequences, the shadow bounded by measurement and not confirmed, conditional on the interior's fate              | clicks; the computed pair's values lattice                                           | $hN = 1$ , Eq. (2)                           | Sections 5.1 and 5.2; S.15, S.19, S.20, S.30, S.31, S.51 |
| The guard is the Courant-Friedrichs-Lewy condition [4]; under the composed paces no Link closes in the rational paces; a frozen Node is dark                                                                                            | theorem for the edge; derived for the closing; assumption for the frozen Node's energy                                                                                                                                                                    | lattice                                                                              | Rule3's coefficients                         | Section 2.3; S.3, S.39                                   |
| A click is the meeting of two quanta, decided by the counting rule's one draw, written at one Node; one click per quantum (antibunching); no signal; the write is the NodeDetector's act; the direction of time belongs to the clicks   | assumption for the draw, the write and the counting rule; derived under the credit's one draw, the NodeDetector's declaration, for exclusivity; derived for antibunching and the one form; theorem for the step; the arrow the NodeDetector's declaration | clicks; lattice for the write and the arrow                                          | the meeting, the NodeDetector's declaration  | Section 2.5, the steps paragraph; S.46                   |
| Born's rule is the NodeDetector's declaration; the uncertainty relation is its declarations, the region, the counting window and the random draw; the draw's frequencies within $2^{-b}$ , $b$ the declared bound's exponent            | declaration; derived under it                                                                                                                                                                                                                             | clicks                                                                               | the counting rule                            | Section 2.5; S.8, S.46                                   |
| Only a click moves a quantum between fields; between clicks every count is constant                                                                                                                                                     | derived, two derivations                                                                                                                                                                                                                                  | clicks                                                                               | Rule3's lines                                | Section 2.5; S.46                                        |
| Light is the charge potential's own configuration; a steady rotation writes no light, a breathing bound state at its beat; a uniformly moving one none in the long-wavelength band                                                      | derived                                                                                                                                                                                                                                                   | lattice (the Wronskian and the source); clicks (the far NodeDetector's rate) lattice | the write                                    | Sections 2.5 and 4.3; S.40                               |
| A potential field's reach from its gap, Yukawa's form; Poisson's rest $3s/(4\pi r)$ ; gravity enters as a slowed clock                                                                                                                  | derived                                                                                                                                                                                                                                                   | lattice                                                                              | Rule3 for a gapped pair                      | Section 4.1; S.11, S.37                                  |

| The claim                                                                                                                                                                                                                                                                                                                        | Status                                                                                                                                       | Fence                                                                                               | Rests on                              | Where                                           |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|---------------------------------------|-------------------------------------------------|
| The vacuum level 60 at $\Gamma = 6,000$ , above the lattice's floor and the swing read on the lattice; no click bounds it above; gravity's events at light's speed in every level                                                                                                                                                | declaration above two floors; derived for the speeds                                                                                         | lattice for the level and its floors; clicks for the speeds                                         | every field reads the potential level | Section 4.1; S.12, S.48                         |
| The binding potential's pair and range and gravity's divisor, chosen where nature's scale forced them                                                                                                                                                                                                                            | the pair and the divisor an assumption, fitted, with a reservation; the range derived from the pair (S.11)                                   | lattice                                                                                             | the declared universe                 | Section 4.1; S.11                               |
| The bound state's fixed point and functional; the stable bound state along the dilation to first order in the well; the two branches                                                                                                                                                                                             | theorem for the equation; derived for the stability, the integer version open; computed for the branches                                     | lattice                                                                                             | the functional                        | Section 4.2; S.13, S.45                         |
| No standing bound state is claimed from a run; the pixel's budget $(\text{Link}/R)^2$ , $\sqrt{n}/A$                                                                                                                                                                                                                             | a lattice reading, named                                                                                                                     | lattice                                                                                             | the runs                              | Section 4.2                                     |
| Bohr's form with the nucleus's angle alone; the self-read gives the law's form at large $Z$ only; hydrogen's lines under line (1); the criterion, what nature splits is a bound state                                                                                                                                            | derived in the slow limit; line (1) of S.52 an assumption, fitted; line (2) an assumption; the criterion the model's decision, an assumption | clicks                                                                                              | the charge potential                  | Section 4.2; S.43, S.52                         |
| The adiabatic invariant and Planck's relation; the drift under a moving well; the clock's rate $1 - f(\omega_0)U$ ; the resonance at $2\omega_b$ under the pace read alone (none under the rotation, S.41) and Rabi's rate, two derivations; a bound state counts photons at Millikan's slope $T$ ; no recoil in the law's write | derived (the clock's rate under the one assumption; the invariant computed)                                                                  | clicks; the drift lattice                                                                           | Rule3 at the bound state's Nodes      | Section 4.3; S.14, S.15, S.16, S.41             |
| A cluster: the three levels at a Node as the nesting, three kinds of binding, the click the only thing between clusters; dark matter in kind                                                                                                                                                                                     | definition, derived; the source a hypothesis                                                                                                 | lattice, clicks                                                                                     | every bound state is bound            | Section 4.4; S.29                               |
| Friedmann's dust equations and Hubble's law at $z \ll 1$ ; the deceleration $q_0 = \Omega_m/2 > 0$                                                                                                                                                                                                                               | derived, two derivations, under the declared homologous start                                                                                | clicks                                                                                              | Kepler's $G_K$                        | Section 4.4; S.29                               |
| The two potentials and the Kepler map; the moving clock at $c_m$ with its fourth order; de Broglie; the fall and the equivalence between fields, $\eta = 32.3 \omega_c^2$ ; the lens with its chromatic term; Coulomb over $\gamma_L$ (the Lorentz factor) and magnetism                                                         | derived; for Coulomb and magnetism theorem for the conservation, derived for the force's form under the rotation, not under the plain read   | clicks; the index lattice; for Coulomb and magnetism lattice for the identity, clicks for the force | the band, the paces, the rotation     | Section 5.1; S.18, S.21, S.22, S.28, S.33, S.44 |

| The claim                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | Status                                                                                                                                    | Fence                                                    | Rests on                                    | Where                                     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------|---------------------------------------------|-------------------------------------------|
| $G$ as a reading, Eq. (11);<br>$\alpha_{1aw} = (3\sqrt{3}/8\pi) k/(\Gamma E_s)$ ; charge universality by construction; the two forces' ratio $(m_p/m_e)^2$ ; the units; the Link's and the gap's bounds                                                                                                                                                                                                                                                                                                                                | derived; $k$ declared; charge universality and the forces' ratio in the small-gap limit, by construction, no test; the bounds comparisons | clicks                                                   | the rotation, Poisson's rest, LHAASO, MAGIC | Section 5.1; S.24, S.25, S.26, S.27, S.42 |
| The gravity of light at $1 \times E/c^2$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | derived under the static kernel, an assumption named, the packet's field open; untested                                                   | clicks                                                   | the static kernel                           | Section 5.2; S.39                         |
| The two-path check is 0.989 for a strontium clock at one metre for one second                                                                                                                                                                                                                                                                                                                                                                                                                                                          | derived; untested                                                                                                                         | clicks                                                   | S.31's scaling of the beat                  | Section 5.2; S.39 (1)                     |
| The two slits: the two slits' 284.7 by the law's line, the design's blind row 273, the Huygens sum, light's line propagated from the two gaps on the design's lattice; the run $N = 285$ at both seeds within the integer step's walk, the seed-24 clicks 28 of 275 from the prediction's counts (275 the blind row's sum by its per-Node rounding, 273 its $N$ ), a comparison and not a statistical test; the inflow's peak at 69.0 against 62.0, the flight time 62.2                                                               | computed; the run a check                                                                                                                 | clicks; 674 photons at the light's own rotation, lattice | Rule3 on the declared universe              | Section 6; S.23                           |
| Bell's $E = \cos 2(a - b)$ ,<br>$S = 478/169$ , Tsirelson's $2\sqrt{1 + \rho^2}$ , Greenberger-Horne-Zeilinger's (GHZ) $M = -4$ ; every local counting rule gives at most 2; no signalling                                                                                                                                                                                                                                                                                                                                             | the click's one form a postulate; the square declared; derived under them; computed for the numbers; theorem for the local bound          | clicks                                                   | the counting rule, the meeting's square     | Section 7; S.10, S.38, S.47               |
| The misses, twelve by name: nine findings against nature, $q_0 > 0$ , the nuclear binding and its size law, no Fermi constant $G_F$ , the $Z$ and parity (one root), the atom's 1s under the self-read, the mass ratios and the cluster's defect, the fine structure, helium's correlation energy and positronium's binding, and the $\times$ polarisation, derived under the static read, which the odd write keeps; three places with no line, the neutron's lifetime, Pauli's exclusion and the loops ( $g - 2$ and the Lamb shift) | derived, the comparisons supporting findings; no line for the three                                                                       | clicks; lattice for helium and positronium               | Rule3 and the declared entries              | Section 8.1; S.35, S.50, S.52             |
| Neither layer keeps a total energy                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | theorem for the step's conservations; assumption for the clicks'                                                                          | lattice, clicks                                          | Rule3, the counting rule                    | Section 2.4; S.49                         |
| The universe's physical coefficients $E_g$ , $k$ and the gaps are each read from one measurement of nature; the binding pair and range, $T$ , the integer bound and the Link unit are declared; no number comes from the lattice itself                                                                                                                                                                                                                                                                                                | calibration, experiment for $E_g$ , $k$ , the gap and the Link's length; hypothesis for any such number                                   | clicks                                                   | nature's measurements                       | Section 8.2; S.48                         |

| The claim                                                                                                                                                                                                                                                                                                 | Status                                                                                                                                                          | Fence                                                                                                             | Rests on                      | Where                                                                 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------|-------------------------------|-----------------------------------------------------------------------|
| The declarations: the entries, the NodeDetectors, the rates, the conversion's table, the initial data, the receding face, the one draw, the erasing front and the empty wave beyond it                                                                                                                    | assumption for every declaration                                                                                                                                | lattice, clicks                                                                                                   | the law's list                | Section 8.2                                                           |
| The hypotheses under their own names: the conversion click, the dimension-3 field with its confinement in the model's words, the carrier, the nuclear potentials, the parts as particles, the masses as fields, only a click moves a quantum in nature (the single atom's lifetime its test)              | hypothesis; no number of the paper rests on the last                                                                                                            | clicks                                                                                                            | stated under their names      | Section 9; the long version's Section 10.3 and S.36, S.37, S.45, S.53 |
| The click's algebra beyond the law: the count's change and the booking identity; the draw of whether and which future continues; the momentum's chain; under the twist proposition the wave vector conserved, the recoil $\propto 1/M$ , Compton with the factor $\omega_0/\tan \omega_0$ , nature's at 1 | derived for the count, the booking and the chain; assumption, the counting rule, for the draw; hypothesis, fitted, for the twist, derived under it for the rest | lattice; clicks for the booking; Compton's factor lattice at the computed pair, nature's by construction, no test | the law's definitions; no run | Section 2.5.2; S.55                                                   |

## Nomenclature

Every letter of the paper and this supplement, with each of its senses; a letter that serves twice is named at each first use as well; the glossary of the long version's words, with the paper's beside them, stands above.

*A* an amplitude, a mass number and Bell's side;  $A_0$ ,  $A_a$  the vector potential on the time step and the Link (S.33, S.41)

*a* a Node's level ( $a_t$ ,  $a_{\text{now}}$ );  $\text{arr}_j$  the level of the neighbour  $j$  in Rule3's line, the six Ports summed (Eq. (1)); a radius (the ball's  $a(t)$ , Bohr's  $a_0$ ), Bell's setting, the pressure's coefficient and the axis index where so named; the fall's acceleration (Section 5.1, S.18) and the semi-major axis

*B* a standing record's peak amplitude, the slope of the band-edge solution  $A + Bt$ , Bell's side and the bound mode's binding where so named

*b* the impact parameter, the leaving level in the booking identity, Bell's setting, the hol-low's coefficient and the axis index where so

named;  $b_L$  the light level read;  $b = 1,241$  the laid packet's peak level (Section 6); **b** the vector of before levels in S.6, with **n** the now levels;  $b_t$  a second standing record's level (S.17)

*C* a body's count in quanta; an amplitude or a coefficient in the supplement where so named;  $C_i$  the counts at the Nodes; the third Greenberger-Horne-Zeilinger (GHZ) side;  $C_1(k)$ ,  $C_0(k)$  the step symbol's coefficients in S.56 (d)

*c*,  $c_m$  light's speed, and the content in level units where so written ( $c$ ,  $c_{\text{vac}}$ ,  $c_{\text{law}}$ ); the band's kinetic scale (Section 2.2, used in 5.1);  $c_{\text{local}} = cN^2$  the local light speed at the composed paces (Eq. (9));  $c_K$  the lattice

Klein-Gordon equation's coefficient,  $c_s$  a family's own light speed ( $c_s^2 = \text{num}_s/(3 \text{den}_s)$ ),  $c_k$  a part's laid weight (Eq. (12)), the third GHZ setting and the well's coefficient where so named;  $c_n, c_a$  cosines in S.7 and S.10  
*D* the form of a record  
*d* a radius, the slits' distance and a dimension where so named;  $d_i$  a Node's weight,  $d_{jm}$  the dipole element,  $d = \text{den} - \text{num}$  (S.27);  $d_L$  the luminosity distance (S.29);  $d_1, d_2$  two bodies' distances from a Link (S.17)  
*E, E(a, b)* the energy, the conserved form of Eq. (4), the part's wall in Eq. (1) and the level weights  $E_s, E_g$  (gravity's) and  $E_r$ , the divisor of the potential field  $r$  (S.13); Bell's correlation,  $E_3$  the three-quantum Greenberger-Horne-Zeilinger correlation  $E_3 = \cos 2(a + b + c)$  (Section 7.2);  $E_0$  the field's amplitude (S.41)  
*e* Euler's number, the share  $e_i$ , a port's coefficient  $e_k$ , the unit charge (one quantum's Wronskian  $T/2$ , Section 5.1), an orbit's eccentricity, a record's ellipticity and the part label  $e$   
*F* the current, the functional, a force, each where so named  
*f* the redshift's factor  $f(\omega_0) = 2(1 - \cos \omega_0)/(\omega_0 \sin \omega_0)$ ,  $f_v$  the moving clock's factor, the sourcing family index in  $w_f q_f$ , a generic function, and the subscript of the frozen content  $U_f$   
*G* Newton's constant,  $G_K$  as the orbits read it,  $G_{\text{clock}} = 1/(4\pi\Gamma E_g)$  as the clock's potential reads it, before Kepler's factor (Eq. (11)), and  $G_0(r)$  the lattice's Green's function (S.37), and the Link's unit  $G, G_{ab}$  the stress reading;  $G_F$  the Fermi constant  
*g* the rest term's coefficient, the metric's components, the ground part  $g$  and Landé's  $g_L$ ;  $g_{ij}$  the sign holder's Link current (S.33); Earth's acceleration in  $gh/c^2$  (S.39), Yukawa's coupling  $g^2$  (S.37), the transfer generator  $g = \Omega_R$  (S.41) and the anomaly  $g - 2$   
*H* Hubble's rate ( $H_0$ );  $T_H$  Hawking's temperature  
*h, h\_a(i)* the ruler of Section 2.3 and Planck's constant where nature's is meant; the Link's stress reading (the long version's tension; S.7); a hazard per window,  $(1 - h)^n$  (S.49 (25) of

the long version); the height in  $gh/c^2$  (S.39);  $h_{00}, h_{aa}, h_{xy}$  the metric perturbation's components (S.50 (1)), beside  $h = p_0/p_a$  the Link factor  
*I* the adiabatic invariant  $A^2 \sin \omega/T$ , the number of quanta; the integral  $I(z)$  of the running exponent (S.13)  
*i* the Node index ( $i, j$  its neighbour) and the imaginary unit in  $e^{i\theta}$ ; a bound mode's index where paired with  $j$   
*J* the joint share  $J(o_A, o_B)$  of Bell's pair over the two sides' ports  $o_A, o_B$ ,  $J_a$  the sign's Node current on the axis  $a$ , and the atom's Hartree self-term where so named  
*j* the neighbour Node index, a bound mode's index, a spatial index ( $g_{0j}, V_j$ ), nature's current density where so named  
*K* Friedmann's constant, the start's kick and a pressure's weight where so named;  $K_{jm}$  the two modes' coupling (S.41)  
*k*  $k_L$  light's wave number and  $\mathbf{k}_L$  its wave vector (S.55); the wave number and the sign holder's write weight;  $k_{\text{in}}$  the wave number a configuration takes on entering a potential level  $U$  at fixed rotation (Eq. (10));  $k_w$  the write's and  $k_r$  the read's weight,  $k = k_w k_r$  (S.26); the part's label in Bell's sums (Eq. (12));  $k_B$  Boltzmann's constant; the 1s cloud's curvature in atomic units in S.52 (b)  
*L* and  $\ell$   $L_{\text{box}}$  the declared box's side in Links (S.55); both a holder's level in levels,  $L$  also a length in Links where so named;  $\ell_{\text{Link}}$  the Link's length in metres (S.54)  
*l* the ray's path element  $dl$  and a level increment in Cauchy's equation  $P(U + l) = P(U)P(l)$   
*M* the taker's count at a click (Section 2.5.2, S.55); the mass in  $GM$  and Mermin's  $M$ ;  $\mathbf{M}$  Rule3's step matrix, with  $\mathbf{D}$  the weights' diagonal and  $\mathbf{K} = \mathbf{DM}$  (S.56 (a));  $M_1, M_2$  the parts' products (S.10)  
*m* the rest term of the lattice Klein-Gordon equation,  $m^*$  the inertia,  $m = GM/c^2$  in Links in the shadow's radius and Rule3's margin  $m(c)$ ; matter's numerator in  $[m, \text{den}]$ ; a bound mode's index ( $\phi_m, \omega_m$ ) and  $m_\ell$  the orbital quantum number (S.43)  
*N* the books' count and the lapse  $N = p_0/\Gamma, e^{-U}$  to  $O(U/\Gamma)$ ; a number of Nodes

$(N_x \times N_y \times N_z)$ , intervals or bodies where so named,  $N_R$  a region's count  
 $n$  a count of intervals or quanta where so named; light's index  $n = e^{2U}$ ;  $\mathbf{n}$  a unit direction,  $n_a$  its components;  $n = 1$  nature's survival exponent (S.49 of the long version); a mode's index ( $E_n, \omega_n$  of the modes)  
 $O, o_A, o_B, O_h$  the cube's symmetry group of order 48,  $O(3)$  the rotation group of a three-line record,  $O(\cdot)$  the order symbol;  $o_A, o_B$  the two sides' ports in Bell's joint share (Eq. (12))  
 $P$  the momentum density, a probability, the guard's edge on a pace and the momentum  $P = cN^2k$  at the composed paces;  $\mathbf{P}$  a signed permutation (S.4),  $P(U)$  the pace as a function of the content (S.34)  
 $p_0, p_i, p_a$  the paces and  $q_{ij}$  the Link's factor, with  $(p, q)$  Bell's setting pair;  $\mathbf{p}$  the momentum (Section 5.1),  $p$  an orbit's semi-latus rectum (S.19) and nature's pressure (S.39)  
 $Q, Q_R$  the two-level form and  $q_{ij}$  the Link's factor; a region's inflow;  $Q(t)$  a second moment (S.40)  
 $q, q_{ij}$  the Link's factor, a body's charge  $q = \pm Ce$ , Bell's setting in  $(p, q)$ , the momentum transfer of a hard click, the second argument of the two-level form  $Q(p, q)$ , and  $q_0$  the deceleration parameter;  $q_f$  the sourcing field  $f$ 's quantity at the Node in the one write, a form  $D_f$ , a Wronskian  $W_f$  or a stress  $T_{aa,f}$  (Eq. (1))  
 $R$  the read coefficient of Rule3, a body's radius in Links and the binding range; a NodeDetector's region from Section 2.5 on; Rydberg's unit in  $E_{\text{tot}}/(Z^2R)$  (S.43, S.52), a scattering rate where so named;  $\mathbf{R}(\theta)$  the turn's matrix  
 $r$  Rule3's remainder and a radius; the parts' amplitude ratio in Bell's  $\rho = 2r/(1+r^2)$ ,  $r_P$  Pekar's constant,  $r_p$  the proton's radius;  $r_a$  the taker's books' remainder (S.55)  
 $S$  Rule3's self coefficient in Eq. (1), Bell's  $S$  in Section 7 and  $S_t$  the step map;  $S_6$  the six-neighbour sum (S.5),  $S_4 = \sum_a n_a^4$  (S.54); a source's total (S.40)  
 $s$  a source's write per interval over the holder's divisor (the far level  $3s/(4\pi r)$ , Section 5.1, S.20, S.25), a share, the refinement scale, the family index and a scaling power where so named;  $s(x) = 2\sin(x/2)$

(S.28, S.31, S.34),  $s_{\parallel}, s_{\perp}$  a separation's components (S.39)  
 $T$  the invariant's unit and  $T_{aa}$  the tension;  
 $T_b$  a beat's length (S.17)  
 $t$  the interval (the time index):  $a_t, S_t$ , the click  $(R, t)$ ;  $t_a(i, j)$  the Link's tension read into  $q_{ij}$   
 $U$  the potential, the clock's fraction  $c/\Gamma$ ;  
 $U_K = G_K M/(c^2 r)$  the potential the orbits of matter read, Kepler's (Eq. (8))  
 $\mathcal{U}$   $\mathcal{U}_t = e^{-i\theta_t}$  the time turn (S.41)  
 $u$  the direction call's working integer, the turned level  $u_i$ , the separation in units of the width (S.13's integration variable), the expansion variable  $1 - \cos k$ , the band-edge solution  $u_t$  and a mode's projection  $u_n$  where so named; the gauge-turned record  $u_t$  in S.41 (a)  
 $V, V_r$  a holder's well energy at the dilation 1,  $V_j$  a post-Newtonian vector potential, Yukawa's  $V(r)$ ,  $V_{ij} = -V_{ji}$  the level of gravity's odd line along a Link, the mean of its two ends' levels, sourced by the count's flux (S.56), the slow limit's potential term and the Link-phase perturbation operator where so named  
 $v$  a velocity (the group velocity, a source's speed,  $\mathbf{v}$ ), the level Rule3 would write at the Node in  $\Delta(wQ) = w(x-v)(x-b)$ , the turned level  $v_i$  and a mode's projection  $v_n$  where so named  
 $W$  the Wronskian,  $W_c$  and  $W_{\text{rec}}$  the credit's units; a holder's declared write weight ( $W, W_1, W_2$  of the nuclear holders) where so named;  $W_j$  a post-Newtonian vector potential (S.35)  
 $w$  the wall  $w = 6 \text{ den } \Gamma^2$ , Rule3's divisor;  
 $w_f$  a sourcing family's weight;  $w_p$  a packet's width  
 $X$  the state space and the quadrature reference record with  $Y'$  where so named  
 $x$  a content fraction, a level or a coordinate where so named;  $x_t$  the lattice's state at the interval  $t$  (Section 2.5); the draw's state in S.46  
 $Y, Y_i$  the line's read  $\sum_b R_b(\text{now}_{i+b} + \text{now}_{i-b}) + S_i \text{ now}_i$ ;  $Y'$  the quadrature reference record

$y$  a lattice coordinate or row, the before level in the direction call  $(x, y, \rho)$ , and the second vector of levels in  $Q(\mathbf{x}, \mathbf{y})$   
 $Z$  the nuclear charge number; the  $Z$  boson ( $M_Z$ )  
 $z$  the redshift and the complex level where so named; the direction call's quotient (Theorem 1);  $z = \kappa R$  in S.13  
 $\alpha$  the fine-structure constant, the redshift's  $\alpha_U$ ,  $\hat{\alpha}$  the bending angle (Eq. (9)) and the frame's  $\alpha_1, \alpha_2$ ;  $\alpha_k$  a part's amplitude at one side's port (Eq. (12)),  $\alpha_K$  the post-Newtonian parameter against Kepler's potential,  $\alpha_g(s)$  gravity's coupling; a phase in S.17 with  $\beta$   
 $\beta$  the post-Newtonian parameter and a speed over light's;  $\beta_k$  a part's amplitude at the other side's port (Eq. (12)); a standing record's phase in S.17  
 $\chi$  the ratio of the two energy changes under the write (S.26); the post-Newtonian superpotential where so named  
 $\Delta$  the six-neighbour second difference and, as  $1/m^*$ , the band's curvature; a detuning ( $\Delta = \omega_L - 2\omega_b$ ), a spread ( $\Delta\omega, \Delta y, \Delta U$ ) and the centred difference  $\Delta_a$  where so written  
 $\delta$  the remainders' change  $r - r'$ , the settings' difference  $a - b$  (S.10), a rotation's shift from rest, a small change, the breathing's depth, the detuning per interval, Kronecker's  $\delta_{ab}$ , and the partons' fixed profiles  $\delta_0, \delta_1$ , each where so named  
 $\epsilon$  the step's remainder term  $(r - r')/w$  and a resonance's modulation depth where so named;  $\epsilon_4 = (3 \sum_a n_a^4 - 1)/36$  the band's quartic coefficient (S.54),  $\epsilon_{ij}$  the two modes' modulation,  $\epsilon_a(i)$  the light's profile (S.41); the mode's frequency over  $Z^2 R$  in S.43 (c); the odd real coefficient of a one-part line in S.56 (d)  
 $\eta$  the Eötvös parameter;  $\eta_N$  the Nordtvedt parameter (Section 5.1)  
 $\Gamma$  the Node clock, the lattice's clock ( $\Gamma = 6,000$ ), the clock integer of the pace's base  $1 - 1/\Gamma$   
 $\gamma$  the post-Newtonian parameter,  $\gamma_K$  against Kepler's potential, Lorentz's factor ( $\gamma_L$ , the coupling's fall at speed in S.56;  $\gamma_s$  a family's own), and  $\gamma_{GW}, \gamma_{EM}$  the gravitational wave's and the electromagnetic wave's Shapiro-delay parameters

$\hbar$  Planck's reduced constant, nature's;  $T$  stands for it in the lattice's units  
 $\kappa$  a holder's inverse reach  
 $\Lambda$  the fringe's period in rows (S.8, S.46); the cosmological constant in the  $\Lambda$ CDM model (cold dark matter with a cosmological constant)  
 $\lambda$  the wavelength and a Lagrange multiplier; the ratio  $U/U_K$  (S.19), the tail's level ratio (S.11), a gauge parameter (S.35); a root of the step's symbol (S.56 (d));  $\bar{\lambda}_e$  the electron's reduced Compton wavelength (Section 5.1)  
 $\mu$  a growth rate and  $GM/c^2$  where so written; the reduced inertia of a pinned pair where so named  
 $\nabla$  the gradient;  $\nabla^2$  the Laplacian  
 $\nu$  the pair's ratio num/den =  $\cos \omega_0$ ; the neutrino in the pair ( $\nu, e$ )  
 $\Omega$  a transition's rotation and the moving centre's rotation  $\Omega(k)$ ;  $\Omega_R$  Rabi's rate,  $\Omega_w$  the weak drive's,  $\Omega_m$  and  $\Omega_\Lambda$  the density parameters  
 $\omega$  a record's rotation per interval, the frequency:  $\omega(k)$  the band,  $\omega_0$  the rest rotation (the gap),  $\omega_b$  a bound body's,  $\omega_s$  the family  $s$ 's rest rotation (its mass),  $\omega_e$  the electron's and, in a transition, the excited part's against the ground part's  $\omega_g$ ,  $\omega_L$  the light's,  $\omega_h$  the holder's own events',  $\omega_{zp}$  the nucleus's zero-point,  $\omega_i, \omega_j, \omega_n$  the bound modes';  $\omega_n$  also the neutron's gap,  $\omega_p$  the proton's  
 $\Phi$  Newton's potential  $\Phi = -c^2 U$ ; the plaquette flux of the orbital momentum;  $\Phi_t$  the time turn's accumulated phase  
 $\phi$  the body's normalised record or mode ( $\phi_i, \phi_m, \phi_\lambda$ ); a phase in  $\cos(kx + \phi)$ ; a retarded potential in S.39 (2)  
 $\Pi$   $\Pi = E\mathbf{v}$  the covariant momentum variable (S.51)  
 $\pi$  the number  $\pi$   
 $\psi$  the ray's angle to the radius,  $nr \sin \psi = b$ ; the slowly varying envelope of the slow limit  
 $\rho$  Bell's  $2r/(1 + r^2)$  and the dust density; the direction call's carried remainder (Theorem 1);  $\rho_i = w W_i$  the continuity equation's density (S.33);  $\phi^2$  of unit mass (S.13)  
 $\sigma$  a holder's source per interval, a sign and a spread where so named; a cube map (S.4)

$\tau$  the lifetime (the hazard  $1/\tau$ ),  $\tau_{\text{hold}}$  the hold of the two paths,  $\tau_D$  the dark periods' declared mean,  $\tau_{\text{int}}$  the interval in seconds  
 $\Theta$   $\Theta_t$  the midpoint phase of the adiabatic ansatz (S.14)  
 $\theta$  the sign holder's turn angle ( $\theta_t, \theta(r), \theta_0$  the nucleus's),  $\theta_L$  the light's angle on the Link,  $\theta_a$  the Link phases,  $\theta_{ij} = -\theta_{ji}$  the Port's phase from the odd line's level along the Link (S.56),

the pairs' common phase and nature's GHZ angles where so named; Compton's scattering angle in S.55  
 $\vartheta$  the modulation's phase in  $\cos(\omega_L t + \vartheta)$  (S.16)  
 $\xi$  the post-Newtonian (Whitehead) parameter  
 $\zeta$  the detuned slow amplitude (S.16);  $\zeta_1$  a post-Newtonian parameter

## The model (Section 2 of the paper)

**Derivation S.1** (the band at a pace, Eq. (3) of the paper) Inputs: *Rule3's line at paces where the two Links of an axis share one  $R_a$ ,  $2 \text{ num } \Gamma^2$  at the vacuum's and  $2 \text{ num } p_a^2$  in general,  $w a_{\text{next}} = \sum_a R_a (\text{arr}_{+a} + \text{arr}_{-a}) + S a_{\text{now}} - w a_{\text{before}}$ , with the remainders dropped (they are bounded by  $w$  and, below one level per Node per interval, leave a plane wave's rotation unmoved at amplitudes large against a level, the paper's clause of Section 6.2, below one level per Node per interval; at an amplitude of one level the integer orbit is its own cycle).* Steps: for a plane wave  $a_t(\mathbf{x}) = A \cos(\mathbf{k} \cdot \mathbf{x} - \omega t)$  the two arrivals of the axis a sum to  $2 \cos k_a a_{\text{now}}$  and  $a_{\text{next}} + a_{\text{before}} = 2 \cos \omega a_{\text{now}}$ , so

$$\begin{aligned} 2 \cos \omega &= \frac{2 \sum_a R_a \cos k_a + S}{w} \\ &= \frac{4 \text{ num } \sum_a p_a^2 \cos k_a + 12 \text{ den } \Gamma^2 - 12(\text{den} - \text{num}) p_0^2 - 4 \text{ num } \sum_a p_a^2}{6 \text{ den } \Gamma^2}, \end{aligned}$$

that is  $\cos \omega = 1 - g(p_0/\Gamma)^2 - \frac{\text{num}}{3 \text{ den}} \sum_a (p_a/\Gamma)^2 (1 - \cos k_a)$ , which is Eq. (3). At the vacuum's pace,  $p_0 = p_a = \Gamma$ , this is  $\cos \omega = \text{num}/\text{den} - \frac{\text{num}}{3 \text{ den}} \sum_a (1 - \cos k_a) = \frac{\text{num}}{3 \text{ den}} \sum_a \cos k_a$ ; at  $k = 0$ ,  $\cos \omega_0 = \text{num}/\text{den}$ . Written as  $2(1 - \cos \omega) = 2(1 - \text{num}/\text{den}) + \frac{\text{num}}{3 \text{ den}} \sum_a 2(1 - \cos k_a)$ , the band is the discrete Klein-Gordon form with  $c_K^2 = \text{num}/(3 \text{ den})$  and  $m^2 = 2(1 - \text{num}/\text{den})$ , an identity of Eq. (3). Light's speed and dispersion: for  $[1, 1]$  along one axis  $\cos \omega = (2 + \cos k)/3$ ; expanding,  $1 - \omega^2/2 + \omega^4/24 = 1 - k^2/6 + k^4/72$ , so  $\omega = (k/\sqrt{3})(1 - k^2/36 + \dots)$  and the group velocity  $d\omega/dk = \sin k/(3 \sin \omega) = (1/\sqrt{3})(1 - k^2/12 + \dots)$ : the speed  $1/\sqrt{3}$  at long wavelength and the group speed falling as  $1 - 0.083 k^2$ . Along the diagonal  $k_x = k_y = k_z = k$  the band is  $\cos \omega = \frac{1}{3} \cdot 3 \cos k = \cos k$  exactly, with no dispersion; one Link per interval along the axes is the octahedron  $|x| + |y| + |z| = t$ , whose face  $x + y + z = t$  stands at the distance  $t/\sqrt{3}$ , light's  $1/\sqrt{3}$ . Check: at  $k = 0.05, 0.1$  and  $0.2$  the exact  $\sqrt{3} d\omega/dk - 1$  over  $k^2$  is  $-0.0833, -0.0834$  and  $-0.0835$ . Status: derived; lattice.

**Derivation S.2** (light's index at a content, Section 2.3) Inputs: *Eq. (3) with the paces of Eq. (2),  $p_0 = \Gamma(1 - 1/\Gamma)^L = \Gamma N$ ,  $N = e^{-x} + O(1/\Gamma)$  at the content  $x = L/\Gamma$  ( $L$  the holder's level, the  $c$  of the preamble), and  $p_a = p_0^2/\Gamma = \Gamma N^2$  for a reader at the weight 1 and no tension.* Steps: for light,  $g = 0$ , so  $\cos \omega = 1 - \frac{1}{3} N^4 \sum_a (1 - \cos k_a)$ , and at long wavelength  $\omega = N^2 |\mathbf{k}|/\sqrt{3}$ . Light's speed at the content  $x$  is  $N^2 = e^{-2x}$  times its vacuum speed, so its index is  $n = e^{2x} = 1/(1 - 2x) + O(x^2)$  and  $n - 1 = 2x + O(x^2)$ : the Link's pace is the clock twice, and so is light's slowing. Status: derived; lattice.

**Derivation S.3** (the guard's edge and the horizon, Section 2.3) Inputs: *Eq. (3).* Steps: the lowest value of  $2 \cos \omega$  on the band, its top  $\omega$ , is at  $k_a = \pi$  on every axis, where  $2 \cos \omega =$

$2 - 2g(p_0/\Gamma)^2 - \frac{4\text{num}}{3\text{den}} \sum_a (p_a/\Gamma)^2$ ; the recurrence  $a_{\text{next}} + a_{\text{before}} = 2\cos\omega a_{\text{now}}$  has no exponentially growing solution iff  $|2\cos\omega| \leq 2$ ; at the edges  $\pm 2$  a linearly growing solution exists ( $u_t = t$  at  $+2$  and  $u_t = (-1)^{t+1}t$  at  $-2$ , from  $u_0 = 0$ ,  $u_1 = 1$ ), which the guard admits, light's checkerboard mode sitting at  $-2$ ; for a negative numerator the lowest  $2\cos\omega$  is at  $k = 0$  and the same bound holds with  $|\text{num}|$ . The guard is the fixed-coefficient one: with the coefficients changing in time the product of the one-step transfers can have a trace below  $-2$  while every instant's band obeys the bound, and a solution grows (S.16's parametric line is the case), so the guard bounds each instant's band and not a sequence of paces in time. With  $p_0 = p_a = p$  the condition  $2\cos\omega \geq -2$  reads  $(p/\Gamma)^2 (2g + 4\text{num}/\text{den}) \leq 4$  at wave number  $\pi$  on every axis and  $2g(p/\Gamma)^2 \leq 4$  at  $k = 0$ , that is  $(p/\Gamma)^2 \leq 2\text{den}/(\text{den} + \text{num})$  for  $\text{num} \geq 0$  and  $2\text{den}/(\text{den} - \text{num})$  for a negative numerator, so the edge is  $P^2 = 2\text{den}\Gamma^2 \div (\text{den} + |\text{num}|)$ , which is  $\Gamma^2$  for light (at  $[-1, 2]$ ,  $\Gamma = 3$  and  $p = 4$ :  $P^2 = 12 < 16$ , the uniform mode's  $2\cos\omega = -10/3$  with the roots  $-3$  and  $-1/3$  growing while the checkerboard's  $\cos\omega = 1/9$  stands). No horizon: under the composed paces  $p_0 = \Gamma(1 - 1/\Gamma)^c > 0$  at every content and  $p_a = p_0^2/\Gamma > 0$ , so  $R_a = 2\text{num}p_a^2$  never vanishes and no Link closes in the rational paces, on the domain  $0 < \text{num} \leq \text{den}$ ; the quadratic truncation  $p_a = \Gamma(1 - 2x)$  vanished at  $x = 1/2$ , a horizon of the truncation. The frozen clock: at a content  $c$  the clock runs at  $N = (1 - 1/\Gamma)^c$  of the vacuum's,  $e^{-c/\Gamma}$  to the unit, so a deep well freezes the Node's Link pace  $p_i = p_0^2/\Gamma$  to 0 at the content  $(\Gamma/2) \ln 2\Gamma$  (S.39), the clock pace there  $\sqrt{\Gamma/2}$ , and the Node's share is undefined there. Status: theorem for the edge and the extreme mode; derived for the horizon of the quadratic form, a feature of the truncation; lattice.

**Derivation S.4** (the cube's group of 48 and its action, Section 2.1) Inputs: Rule3's line, whose six arrivals enter through the six Port coefficients  $R_{ij}$ , the two of an axis one  $R_a = 2\text{num}p_a^2$  with no coefficient naming an axis beyond its own pace; the readings  $D$ ,  $F$  and  $W$  of Eq. (1). The group: a map of the six Ports onto themselves that keeps opposite Ports opposite is a signed permutation of the three axes,  $\mathbf{x} \mapsto \mathbf{P}\mathbf{x}$  with  $\mathbf{P}$  a permutation matrix with signs; there are  $3! \cdot 2^3 = 48$  of them, the group  $O_h$  of the cube, of which the 24 with  $\det \mathbf{P} = +1$  are the rotations and the other 24 reflections and rotatory reflections; with the translations of the lattice they form the space group of the simple cubic lattice. Covariance: let  $\sigma$  be such a map and let the turned lattice carry the levels  $a_{\sigma(i)}^\sigma = a_i$  and the paces  $p_{\sigma(a)}^\sigma(\sigma(i)) = p_a(i)$ . The arrivals of the axis  $\sigma(a)$  at  $\sigma(i)$  in the turned lattice are the arrivals of the axis  $a$  at  $i$  in the original, in either order, and  $R_a$  is even in the axis's sign; so the sum  $\sum_a R_a(\text{arr}_{+a} + \text{arr}_{-a})$  is the same number at  $\sigma(i)$  as at  $i$ ,  $S$  and  $w$  are the same, and the division gives the same quotient and remainder: the step commutes with  $\sigma$ , and a lattice turned by  $\sigma$  runs as the turned lattice. A translation commutes for the same reason. The readings' kinds:  $D_i$  is one number at a Node with no axis in it, carried to  $\sigma(i)$  unchanged, a scalar. The currents  $F_{ij}$  are attached to the Links and carried with them, and the momentum density  $P_a(i) = (F_{i,i-a} - F_{i,i+a})/\text{num}$  (this derivation's orientation, positive for a record moving toward  $+a$ ) changes sign where  $\sigma$  reverses the axis  $a$  and is carried to the image axis otherwise, a vector;  $W_i$  carries no axis and is carried unchanged, a scalar under the cube's 48, odd under the plane's internal reflection  $\text{im} \mapsto -\text{im}$  and under the time reversal. The click  $(R, t)$  names a region and an interval and is carried to the turned region, a scalar. The time reversal: it exchanges next and before and keeps now;  $D_i = \text{now}_i^2 - \text{next}_i \text{ before}_i$  is symmetric in that pair and is unchanged;  $F_{ij}$  and  $W_i$ , antisymmetric in the two adjacent levels they read, are carried to minus the next interval's readings,  $F_{ij}(t) \mapsto -F_{ij}(t+1)$  and  $W_i(t) \mapsto -W_i(t+1)$  (the reversed pair at  $t$  is (now, next)), the reversed current being the forward one read backward; and it carries the step to its inverse by Theorem 1. The rational rotation: the two parts of a plane record obey the same linear line with the same coefficients, so any linear combination of them does; the combination, with  $p, q$ ,

$r$  here the integers of the rotation and not the paces,  $(\text{re}', \text{im}') = ((p \text{ re} - q \text{ im})/r, (q \text{ re} + p \text{ im})/r)$  with  $p^2 + q^2 = r^2$  carries solutions to solutions, and its Wronskian is  $(p^2 + q^2)/r^2$  times the old one, unchanged. What is not in the group: a boost changes the record's wave number and rotation together, which no permutation of Ports does, and a rotation by an angle other than the cube's carries a Node off the lattice; so neither is a symmetry, and Lorentz's factor, where the paper finds it, is a property of the clicks (Derivation S.21). Status: theorem; lattice.

## Exact properties (Section 3)

**Derivation S.5** (the share's change is the currents, Eq. (5)) Inputs: the share at the vacuum's paces,  $e_i = 3 \text{ den}(\text{now}_i^2 + \text{before}_i^2) - \text{num now}_i S_6(\text{before})_i$ , with  $S_6$  the six-neighbour sum; Rule3 at the vacuum's paces, where  $S = 0$ ,  $R_a = 2 \text{ num } \Gamma^2$  and  $w = 6 \text{ den } \Gamma^2$ , so that  $3 \text{ den}(\text{next}_i + \text{before}_i) = \text{num } S_6(\text{now})_i + 3 \text{ den } \epsilon_i$  with  $\epsilon_i = (r_i - r'_i)/w$  the remainder term of the share's change (the paper's Section 3.4) and  $w = 6 \text{ den } \Gamma^2$  the wall at these paces, so that the remainders' term of this line,  $(r_i - r'_i)/(2\Gamma^2)$ , is  $3 \text{ den } \epsilon_i$  (from  $6 \text{ den } \Gamma^2(\text{next} + \text{before}) = 2 \text{ num } \Gamma^2 S_6(\text{now}) + r - r'$ ). Steps: after the step the pair (now, before) becomes (next, now), so

$$e'_i - e_i = 3 \text{ den}(\text{next}_i^2 - \text{before}_i^2) - \text{num}[\text{next}_i S_6(\text{now})_i - \text{now}_i S_6(\text{before})_i].$$

With  $3 \text{ den}(\text{next}_i + \text{before}_i) = \text{num } S_6(\text{now})_i + 3 \text{ den } \epsilon_i$  the first term is  $(\text{next}_i - \text{before}_i)(\text{num } S_6(\text{now})_i + 3 \text{ den } \epsilon_i)$ , and the bracket cancels its  $\text{next}_i$  part:

$$\begin{aligned} e'_i - e_i &= \text{num}[\text{now}_i S_6(\text{before})_i - \text{before}_i S_6(\text{now})_i] + (\text{next}_i - \text{before}_i) 3 \text{ den } \epsilon_i \\ &= \sum_{j \sim i} \text{num}(\text{now}_i \text{ before}_j - \text{before}_i \text{ now}_j) + (\text{next}_i - \text{before}_i) 3 \text{ den } \epsilon_i, \end{aligned}$$

the six currents  $F_{ij}$  into the Node plus the remainder term. At any paces: with the weighted share  $\tilde{e}_i = [w(\text{now}_i^2 + \text{before}_i^2) - S_i \text{ now}_i \text{ before}_i - \text{now}_i \sum_j R_{ij} \text{ before}_j]/(2p_i^2)$  and the line without its division,  $w \text{ next}_i^* = \sum_j R_{ij} \text{ now}_j + S_i \text{ now}_i - w \text{ before}_i$ , the difference  $e_i(\text{next}^*, \text{now}) - e_i(\text{now}, \text{before})$  is  $\sum_j (q_{ij}^2/\Gamma^2) \text{ num}(\text{now}_i \text{ before}_j - \text{before}_i \text{ now}_j)$  with no anisotropy term, the  $S_i$  terms and the Node's terms cancelling as in Theorem 3's proof, since  $R_{ij}/(2p_i^2) = \text{num } q_{ij}^2/\Gamma^2$  is the same from both ends. The integer step's  $\text{next} = \text{next}^* + (r - r')/w$  adds  $-(\text{next}_i - \text{before}_i)(r'_i - r_i)/(2p_i^2)$ , because  $w(\text{next}_i + \text{next}_i^*) - S_i \text{ now}_i - \sum_j R_{ij} \text{ now}_j = w(\text{next}_i - \text{before}_i)$ . Check: exact fractions on a periodic chain of five Nodes at  $[2, 3]$ , the two other axes folded (four self-Ports per Node),  $\Gamma = 10$ , with contents 0 to 3, Link factors 5 to 10 and random levels: the identity holds at every Node, with and without the division; and a second witness, a uniform  $\text{now} = 3$  over  $\text{before} = 0$  at  $\Gamma = 10$ ,  $q = 5$ , reads 81 per Node, half of Eq. (4)'s 162, since  $e_i$  carries  $1/(2p_i^2)$  where Eq. (4) carries  $1/p_i^2$ . Status: two derivations; theorem; lattice.

**Derivation S.6** (the total share is non-negative, Section 3.4) Inputs:  $e_i$  as above, at the vacuum's coefficients, on any lattice whose Links read the same from both ends and whose every beyond reads 0 (a closed, a periodic or an open-faced lattice);  $S_6$  symmetric with row sums at most 6 (six Ports, a Port onto a face reading 0 contributing nothing on either side), so its spectrum lies in  $[-6, 6]$ ; a declared beyond that is not 0 adds linear terms and is outside the theorem (a universe whose massless potential rests at a declared level beyond the open faces among them). Steps:  $\sum_i e_i = 3 \text{ den}(|\mathbf{n}|^2 + |\mathbf{b}|^2) - \text{num } \mathbf{n} \cdot S_6 \mathbf{b}$ , and  $|\mathbf{n} \cdot S_6 \mathbf{b}| \leq \|S_6\| |\mathbf{n}| |\mathbf{b}| \leq 6 |\mathbf{n}| |\mathbf{b}| \leq 3(|\mathbf{n}|^2 + |\mathbf{b}|^2)$ , so  $\sum_i e_i \geq 3(\text{den} - |\text{num}|)(|\mathbf{n}|^2 + |\mathbf{b}|^2) \geq 0$  for  $|\text{num}| \leq \text{den}$ . Equality needs  $|\text{num}| = \text{den}$  and  $\mathbf{b}$  an eigenvector of  $S_6$  at  $\pm 6$  with  $\mathbf{n} = \pm \mathbf{b}$

(**n**, **b** the vectors of the now and before levels): on an even periodic lattice the static uniform level of light or its checkerboard,  $\text{now} = (-1)^{x+y+z}$  with  $\text{before} = -\text{now}$ , whose shares are 0. A Node's share: at fixed  $\text{before}_i$  and  $S_6(\text{before})_i = s$ ,  $e_i$  is a quadratic in  $\text{now}_i$  with the minimum  $3 \text{den} \text{before}_i^2 - \text{num}^2 s^2 / (12 \text{den})$ , negative iff  $|s| > 6(\text{den}/|\text{num}|)|\text{before}_i|$  for  $\text{num} \neq 0$  (at  $\text{num} = 0$  the share is  $3 \text{den} (\text{now}_i^2 + \text{before}_i^2)$  and never negative); a plane record has  $|s|/|\text{before}_i| = |2 \sum_a \cos k_a| \leq 6$  and is never negative, a lay's edge can be. A region: pairing each boundary Link's cross term and bounding  $2|\text{now}_i \text{before}_j| \leq \text{now}_i^2 + \text{before}_j^2$  (the inside  $\text{now}_i^2$  absorbed by the interior row-sum deficit  $(6 - d_i)/2$ , the outside  $\text{before}_j^2$  kept),  $\sum_{i \in R} e_i \geq 3(\text{den} - |\text{num}|) \sum_R (\text{now}_i^2 + \text{before}_i^2) - \frac{|\text{num}|}{2} \sum_{\text{boundary Ports}} (\text{now}_j^2 + \text{before}_j^2)$  over the outside neighbours  $j$  (the numerator's modulus, so that the bound holds at a negative numerator too): a region's share is negative at most by half a level-square per boundary Port in the current's units. Status: theorem; lattice.

**Derivation S.7** (the tension of a plane wave, Section 3.3) Inputs: the stress reading  $G_{aa}(i) = \text{now}_i(\text{now}_{i+2a} - \text{now}_i) - \text{now}_{i+a}(\text{now}_{i+a} - \text{now}_{i-a})$  and the cross reading  $G_{ab}(i) = \text{now}_i \text{now}_{i+a+b} - \text{now}_{i+a} \text{now}_{i+b} + \text{now}_{i+a} \text{now}_{i-b} - \text{now}_i \text{now}_{i+a-b}$ ,  $G_{aa}$  at  $b = a$ . Steps: for  $\text{now}_i = A \cos(kx_i + \phi)$  along the axis, with  $c_n = \cos(kx_i + nk + \phi)$ ,  $G = A^2[c_0 c_2 - c_0^2 - c_1^2 + c_1 c_{-1}]$ . The identities  $c_0 c_2 = \frac{1}{2}[\cos(2kx + 2k + 2\phi) + \cos 2k]$ ,  $c_1 c_{-1} = \frac{1}{2}[\cos(2kx + 2\phi) + \cos 2k]$ ,  $c_0^2 = \frac{1}{2}[1 + \cos(2kx + 2\phi)]$  and  $c_1^2 = \frac{1}{2}[1 + \cos(2kx + 2k + 2\phi)]$  cancel the phase-dependent parts, leaving  $G = A^2(\cos 2k - 1) = -2A^2 \sin^2 k$  at every Node and interval. So the Link's booking is  $-\text{num} G = 2 \text{num} A^2 \sin^2 k$  and the Node's tension  $T_{aa} = -\text{num} h_a = \text{num} A^2 \sin^2 k$  with  $h_a = G/2$  on a plane wave, where the two ends' tensions are equal, and for a wave with the components  $k_a, k_b$  the cross reading is  $G_{ab} = -2A^2 \sin k_a \sin k_b$ . Status: derived; a theorem for the line's identity (the plane wave's stress reading phase-free at every Node), derived for the momentum flux (S.45); lattice.

## Measurement (Section 2.5)

**Derivation S.8** (the region's reading factor and the draw's scatter, Section 2.5) Inputs: the screen's reading factor over  $n$  consecutive rows of a fringe of period  $\Lambda$  (Section 6.2); the draw's binomial scatter (Section 2.5). Steps: a region of  $n$  consecutive rows sums a fringe  $\cos(2\pi y/\Lambda)$ ,  $\Lambda$  here the fringe's period, to  $\text{Re} \sum_{y=0}^{n-1} e^{2\pi i y/\Lambda} = \text{Re} \frac{1 - e^{2\pi i n/\Lambda}}{1 - e^{2\pi i/\Lambda}}$ , whose modulus is  $|\sin(\pi n/\Lambda)/\sin(\pi/\Lambda)|$ , the sum being  $e^{i\pi(n-1)/\Lambda} \sin(\pi n/\Lambda)/\sin(\pi/\Lambda)$ ; over  $n$  rows the fringe's modulation is kept with the signed factor  $\sin(\pi n/\Lambda)/(n \sin(\pi/\Lambda))$  (negative, the fringe inverted, for  $\Lambda < n < 2\Lambda$ ), 0.91 at  $n = 4$ ,  $\Lambda = 16$ , and 0 at  $\Lambda = n$ . A frequency read as a click rate over a window of  $n_w$  intervals resolves  $\Delta\omega \geq 2\pi/n_w$ , the window's own resolution.  $N$  quanta drawn independently by the shares  $P_i$  give each region a binomial count,  $NP_i \pm \sqrt{NP_i(1 - P_i)}$ . Check:  $\sin(\pi/4)/(4 \sin(\pi/16)) = 0.70711/(4 \times 0.19509) = 0.9061$ . Status: derived; clicks.

**Derivation S.9** (the Wronskian's conservation and its current, Section 3.5) Inputs: the two parts of a plane record step by the same read matrix **M**, with **DM** symmetric for **D** = diag( $1/p_i^2$ ) (the paper's Theorem 3),  $\mathbf{re}_{\text{next}} = \mathbf{M} \mathbf{re}_{\text{now}} - \mathbf{re}_{\text{before}}$  and the same for **im**. Steps: let  $W = \mathbf{re}_{\text{now}}^\top \mathbf{D} \mathbf{im}_{\text{before}} - \mathbf{im}_{\text{now}}^\top \mathbf{D} \mathbf{re}_{\text{before}}$ . After the step it is  $\mathbf{re}_{\text{next}}^\top \mathbf{D} \mathbf{im}_{\text{now}} - \mathbf{im}_{\text{next}}^\top \mathbf{D} \mathbf{re}_{\text{now}} = (\mathbf{M} \mathbf{re}_{\text{now}} - \mathbf{re}_{\text{before}})^\top \mathbf{D} \mathbf{im}_{\text{now}} - (\mathbf{M} \mathbf{im}_{\text{now}} - \mathbf{im}_{\text{before}})^\top \mathbf{D} \mathbf{re}_{\text{now}}$ ; the **M** terms are  $\mathbf{re}_{\text{now}}^\top \mathbf{M}^\top \mathbf{D} \mathbf{im}_{\text{now}} - \mathbf{im}_{\text{now}}^\top \mathbf{M}^\top \mathbf{D} \mathbf{re}_{\text{now}}$ , which vanish since  $\mathbf{M}^\top \mathbf{D} = \mathbf{D} \mathbf{M}$  is symmetric, and the rest is  $W$  again. Per Node the same cancellation leaves the Link terms: the

weighted Wronskian  $W_i/p_i^2$  changes by  $(2 \text{ num}/w) (q_{ij}^2/\Gamma^2) (\text{im}_i \text{re}_j - \text{re}_i \text{im}_j)$  summed over the Node's six Ports at the levels now, the Wronskian's current, which for the integer  $W_i$  at the vacuum's paces is  $(\text{num}/3 \text{ den}) (\text{im}_i \text{re}_j - \text{re}_i \text{im}_j)$  per Port. Check: a periodic chain of three Nodes at  $\Gamma = 2$  with the two other axes folded,  $[1, 1]$ ,  $\text{re}_{\text{now}} = (0, 3, 0)$ ,  $\text{im}_{\text{now}} = (3, 0, 0)$  and zeros before steps to  $(1, 4, 1)$  and  $(4, 1, 1)$  with no remainder;  $W_0$  goes from 0 to 3,  $W_0/p_0^2$  by  $3/4$ , and the current is  $(2/24) \cdot 9 = 3/4$ ; exact fractions on a chain of five Nodes with contents and Link factors confirm the identity at every Node. For the circular record  $\text{re} = A \cos \omega t$ ,  $\text{im} = eA \sin \omega t$  at one Node, with  $e$  here the ratio of the two parts' amplitudes,  $W = A \cos \omega t \cdot eA \sin \omega(t-1) - eA \sin \omega t \cdot A \cos \omega(t-1) = -eA^2 \sin \omega$ , constant. Domain: the identity is the line's,  $\mathbf{re}_{\text{next}} = \mathbf{M} \mathbf{re}_{\text{now}} - \mathbf{re}_{\text{before}}$  exactly. The integer step's division adds  $\epsilon = (r - r')/w$  to each part, and the remainder term  $\epsilon_{\text{re}} \text{im}_{\text{now}} - \epsilon_{\text{im}} \text{re}_{\text{now}}$  is not zero in general: at the vacuum's pace  $S = 0$  and  $w_{\text{next}} = 12 \text{ num } \Gamma^2 a_{\text{now}} - w_{\text{before}}$  for a uniform level, so at  $[2, 3]$  the parts  $(1, 3)$  with zeros before step to  $(\lfloor 4/3 \rfloor, \lfloor 4 \rfloor) = (1, 4)$  and  $W$  goes from 0 to  $1 \cdot 3 - 4 \cdot 1 = -1$ . An exactly conserved integer charge is open. Status: theorem, its witnesses computed; lattice.

**Derivation S.10** (Bell's correlation under the declared credit, Section 7.2) Inputs: two parts with the products  $M_1, M_2$  of their amplitudes at the two sides' ports; the port coefficients  $e(+) = (\cos a, \sin a)$ ,  $e(-) = (-\sin a, \cos a)$  on side A and the same with  $b$  on side B; the joint share  $J(o_A, o_B) = (\sum_k e_k(o_A) e_k(o_B) M_k)^2$  over the two sides' ports  $o_A, o_B$ ;  $E = \sum_{o_A, o_B} s(o_A) s(o_B) J(o_A, o_B) / \sum_{o_A, o_B} J(o_A, o_B)$  with  $s(\pm) = \pm 1$ . Steps: write  $c_a = \cos a$ ,  $s_a = \sin a$  and the same for  $b$ . The four sums are  $M_1 c_a c_b + M_2 s_a s_b$ ,  $M_1 s_a s_b + M_2 c_a c_b$ ,  $-M_1 c_a s_b + M_2 s_a c_b$  and  $-M_1 s_a c_b + M_2 c_a s_b$  for  $(+, +)$ ,  $(-, -)$ ,  $(+, -)$  and  $(-, +)$ . Their squares sum to  $(M_1^2 + M_2^2)(c_a^2 c_b^2 + s_a^2 s_b^2 + c_a^2 s_b^2 + s_a^2 c_b^2) = M_1^2 + M_2^2$ , the cross terms cancelling in pairs. The signed sum is  $(M_1^2 + M_2^2)(c_a^2 - s_a^2)(c_b^2 - s_b^2) + 8M_1 M_2 c_a s_a c_b s_b$ , so

$$E(a, b) = \cos 2a \cos 2b + \rho \sin 2a \sin 2b, \quad \rho = \frac{2M_1 M_2}{M_1^2 + M_2^2},$$

and the marginal  $P(A+) = [(M_1 c_a c_b + M_2 s_a s_b)^2 + (-M_1 c_a s_b + M_2 s_a c_b)^2] / (M_1^2 + M_2^2) = (M_1^2 c_a^2 + M_2^2 s_a^2) / (M_1^2 + M_2^2)$ , independent of  $b$ . For equal parts  $\rho = 1$  and  $E = \cos 2(a - b)$ , the known form for parallel polarisations, derived here under the declared credit by Lagrange's identity. The integer settings: with  $e(+) = (p, q)$  and  $e(-) = (-q, p)$  unnormalised and equal parts,  $J(+, +) = J(-, -) = (p_A p_B + q_A q_B)^2$  and  $J(+, -) = J(-, +) = (p_A q_B - q_A p_B)^2$ , whose sum is  $2(p_A^2 + q_A^2)(p_B^2 + q_B^2)$  by Lagrange's identity, so  $E = [(p_A p_B + q_A q_B)^2 - (p_A q_B - q_A p_B)^2] / [(p_A^2 + q_A^2)(p_B^2 + q_B^2)]$ , a rational number. At  $a = (1, 0)$ ,  $a' = (1, 1)$ ,  $b = (12, 5)$ ,  $b' = (5, 12)$ :  $E(a, b) = (144 - 25)/169 = 119/169$ ,  $E(a, b') = (25 - 144)/169 = -119/169$ ,  $E(a', b) = (17^2 - 7^2)/338 = 120/169$ ,  $E(a', b') = 120/169$ , and  $S = E(a, b) - E(a, b') + E(a', b) + E(a', b') = 478/169 = 2.8284$ . At the ideal angles  $S = \sqrt{2}(1 + \rho)$ ; at the integer settings  $S = (238 + 240\rho)/169$ . The two local credits: with the pair's phase  $\theta$  uniform, a credit of each side by the shares of its ports gives  $E = \langle \cos 2(a - \theta) \cos 2(b - \theta) \rangle_\theta = \frac{1}{2} \cos 2(a - b)$  and  $S = \sqrt{2}$ ; a credit by the sign of the larger port gives  $E = \langle \text{sgn} \cos 2(a - \theta) \text{sgn} \cos 2(b - \theta) \rangle_\theta = 1 - 4|a - b|/\pi$ , the sawtooth, whose sum at the four settings is exactly 2. Check: the averages over  $\theta$  on a grid of  $2 \times 10^5$  points give  $S = 1.4142$  and  $2.0000$ , and the contrast credit, one side's count weighted by its contrast  $|\cos 2(a - \theta)|$  and the other side's sign always counted, gives  $E = \cos 2(a - b)$  exactly once divided by one side's efficiency  $2/\pi$ , by  $\langle \cos x \text{sgn} \cos(x - 2\delta) \rangle = (2/\pi) \cos 2\delta$ ,  $S = 2.8284$  ( $478/169 = 2.828402$  against  $2\sqrt{2} = 2.828427$ ); the symmetric readings do not  $|\cos|$  on both sides:  $E(\pi/8) = 0.88$ ,  $S = 3.52$ ). Status: derived under the declared credit; clicks.

## The families (Section 4.1)

**Derivation S.11** (the reach of a holder with a gap, Section 4.1) Inputs: *a holder's rest outside its source, the line with  $a_{\text{next}} = a_{\text{before}} = a_{\text{now}} = a$ :  $6 \text{ den } a = \text{num } S_6(a)$  at the vacuum's pace, that is  $\text{num } (S_6 - 6) a = 6(\text{den} - \text{num}) a$ . Steps: for a rest falling along one axis as  $e^{-\kappa r}$  and uniform across,  $S_6(a) = (2 \cosh \kappa + 4) a$ , so  $6 \text{ den} = \text{num } (2 \cosh \kappa + 4)$  and  $\cosh \kappa = 3 \text{ den}/\text{num} - 2$ , for  $0 < \text{num} \leq \text{den}$  (the positive tail; the axial characteristic equation is  $\lambda + \lambda^{-1} = 6 \text{ den}/\text{num} - 4$  for the ratio  $\lambda$  of neighbouring levels, and at a negative numerator its decaying root is negative, an alternating tail,  $\lambda = -8 + \sqrt{63} = -0.0627$  at  $[-1, 2]$  with  $\cosh \kappa = 8$  for  $\kappa = -\ln |\lambda|$ ; at  $\text{num} = 0$  the static line forces  $a = 0$  outside the source, no tail). For  $\text{num}$  just under  $\text{den}$ ,  $\cosh \kappa = 1 + \kappa^2/2 + \dots$ , so  $\kappa^2 = 6 \text{ den}/\text{num} - 6 = 6(\text{den} - \text{num})/\text{num}$  and the range is  $R = 1/\kappa = \sqrt{\text{num}/(6(\text{den} - \text{num}))}$  Links: 20.00 at  $[2400, 2401]$ ;  $[1, 2]$  has  $\cosh \kappa = 4$ ,  $e^{-\kappa} = 0.127$ . The gap is also the rest rotation of the holder's own travelling events,  $\cos \omega_h = \text{num}/\text{den}$ ,  $\omega_h^2 = 2(\text{den} - \text{num})/\text{den} + \dots$ , whose Compton length at light's speed,  $c/\omega_h = \sqrt{\text{den}/(6(\text{den} - \text{num}))} = R\sqrt{\text{den}/\text{num}}$ , is the range to the gap's first order: Yukawa's identity with no input. Status: derived; lattice.*

**Derivation S.12** (the two speeds, Section 4.1) Inputs: *S.2's  $N^2 = e^{-2x}$  and the law's read of the content by every family. Steps: light at a content  $x$  travels at  $N^2/\sqrt{3} = (1-2x)/\sqrt{3} + O(x^2)$ ,  $N^2 = e^{-2x}$  (Derivation S.2). Under the law's line the massless holder reads the content as every family does, its own level among it, so its travelling events have the same paces and the two arrive together at every content, the difference of their Shapiro delays 0. Were the massless holder to read no content, its events travel at  $1/\sqrt{3}$  and over  $L$  Links the two arrive  $L\sqrt{3}(e^{2x} - 1) = 2xL\sqrt{3} + O(x^2)$  intervals apart at long wavelength, against GW170817's bound [5] on  $\gamma_{\text{GW}} - \gamma_{\text{EM}}$  (the gravitational wave's against the electromagnetic wave's), between  $-2.6 \times 10^{-7}$  and  $1.2 \times 10^{-6}$ ; under the line the measurement bounds no vacuum content. Status: derived; clicks.*

## Bodies (Section 4)

**Derivation S.13** (the bound body: the fixed point and the functional, Section 4.2) Inputs: *Rule3's line at the paces of the wells the record sources; a standing record  $a_{i,t} = \phi_i \cos(\omega_b t)$  with one rotation at every Node; the clock's pace  $(p_0/\Gamma)^2 = 1 - 2\ell_i/\Gamma + \dots$  at a well  $\ell_i$  (in levels,  $\ell_i = \Gamma U_i$ ; Newton's potential is  $\Phi = -c^2 U$ , S.28). The fixed point's equation: with the remainders dropped the line gives at every Node*

$$2 \cos \omega_b \phi_i = \frac{1}{w} \left[ \sum_a R_a(i) (\phi_{i+a} + \phi_{i-a}) + S_i \phi_i \right] = (\mathbf{M}\phi)_i,$$

so  $\phi$  is an eigenvector of the read matrix  $\mathbf{M}$  and the standing record is its top mode, the maximiser of  $\phi^\top \mathbf{D}\mathbf{M}\phi$  on the weighted sphere  $\phi^\top \mathbf{D}\phi = 1$ ,  $\mathbf{D}\mathbf{M}$  symmetric under the law's read coefficient (the paper's Theorem 3) (the rule's own iteration of Section 4.2 repeats this read act). To the first order in  $\ell/\Gamma$ ,  $S_i/w = 2 - 2g(1 - 2\ell_i/\Gamma) - \frac{2\text{num}}{3\text{den}} \sum_a (p_a/\Gamma)^2 = 2 \cos \omega_0 + 4g\ell_i/\Gamma - \frac{2\text{num}}{3\text{den}} \sum_a (p_a/\Gamma)^2$  and  $R_a/w = \frac{\text{num}}{3\text{den}} (p_a/\Gamma)^2$ , so, keeping the band's term at the vacuum's Link pace (to first order in  $U$  for a body whose band term balances its well,  $\Delta\phi = O(U\phi)$ , so that the product of the two is of the second order; a free packet in a well, with  $\Delta\phi$  of the packet's own size, is outside this reduction),

$$(\mathbf{M}\phi)_i = 2 \cos \omega_0 \phi_i + \frac{4g}{\Gamma} \ell_i \phi_i - \frac{\text{num}}{3\text{den}} (-\Delta\phi)_i, \quad (-\Delta\phi)_i = 6\phi_i - \sum_a (\phi_{i+a} + \phi_{i-a}).$$

The functional: the well is sourced by the record itself,  $\ell_i = \sum_r \sigma_r (C/E_r) \sum_j G_{r,ij} \phi_j^2$  for a body of  $C$  quanta with the counts  $C_j = C \phi_j^2$ . Let

$$F[\phi] = \frac{\text{num}}{3 \text{den}} \sum_{\text{Links}} (\phi_i - \phi_j)^2 - \frac{2g}{\Gamma} \sum_r \sigma_r \frac{C}{E_r} \sum_{ij} \phi_i^2 G_{r,ij} \phi_j^2,$$

the long version's equation (12). Its gradient is  $\partial F/\partial \phi_i = \frac{2 \text{num}}{3 \text{den}} (-\Delta \phi)_i - \frac{8g}{\Gamma} \ell_i \phi_i$ , using the symmetry of  $G_r$  (each well counted once, Hartree's form, is what makes the factor  $2g$  in  $F$  give  $4g$  in the gradient). So  $\partial F/\partial \phi = 2\lambda \phi$  on the sphere is exactly the fixed point's equation with  $\lambda = 2 \cos \omega_0 - 2 \cos \omega_b$ : a body is a critical point of  $F$  on the sphere, and the virial and the second variation along the dilation are the paper's proposition (Section 4.2). The scaling powers: under  $\phi_R(\mathbf{x}) = R^{-d/2} \phi(\mathbf{x}/R)$  the sum over Links of  $(\phi_i - \phi_j)^2$  is the lattice form of  $\int |\nabla \phi|^2$  and scales as  $R^{-2}$ , the pressure  $K/R^2$ . The well energy  $\sum_{ij} \phi_i^2 G_{ij} \phi_j^2$  is  $\iint \rho(\mathbf{x}) G(\mathbf{x} - \mathbf{y}) \rho(\mathbf{y})$  with  $\rho = \phi^2$  of unit mass, which scales as  $R^{-(d-2)}$  for the massless kernel  $G \sim 1/r$  and for a screened kernel within its reach, and as  $R^{-d}$  for a screened kernel beyond its reach, where  $G$  acts as a point kernel and the energy is  $\int \rho^2$ : the powers  $s_r$  of the paper's proposition. The cloud's size: for the massless kernel in three dimensions  $F(R) = a/R^2 - c/R$  with  $a = \frac{\text{num}}{3 \text{den}} \int |\nabla \phi|^2$  and  $c = \frac{2g}{\Gamma} \frac{C}{E} \iint \frac{\rho \rho'}{4\pi |\mathbf{x} - \mathbf{y}|} \cdot 3$  at the unit size, so the minimum is at  $R = 2a/c \propto \Gamma E_g/(gC)$ : the radius falls as  $1/C$  and grows with  $\Gamma$ ; the continuum constant of the root-mean-square (rms) radius is  $(2\pi \cos \omega_0/(9g)) r_P \Gamma E_g/C$  with Pekar's  $r_P = 4.635$  [6] (the Choquard minimiser's rms radius, at Lieb's energy  $-0.1085$  [7] and the virial  $1/2$ ),  $(4\pi/9) r_P = 6.47$  at  $[2, 3]$  where  $\cos \omega_0/(2g) = 1$ , the factor  $\cos \omega_0/(2g)$  tending to  $1/\omega_0^2$  as the gap closes (4.5 at  $[9, 10]$ , 500 at  $[999, 1000]$ ). The hollow's barrier (Section 4.2): a hollow beyond the body's reach adds the point-kernel term  $-b/R^3$  (the power  $R^{-d}$  above at  $d = 3$ ),  $F(R) = a/R^2 - b/R^3 - c/R$ ;  $F'(R) = (cR^2 - 2aR + 3b)/R^4$  vanishes at  $R_{\pm} = (a \pm \sqrt{a^2 - 3bc})/c$ , two critical points where  $a^2 > 3bc$  and none where  $3bc > a^2$ ; at a root  $F'' = 2(aR - 3b)/R^5$ , positive at  $R_+ \geq a/c > 3b/a$  and negative at  $R_- = 3b/(a + \sqrt{a^2 - 3bc}) < 3b/a$ , so the wide critical point is the minimum and the narrow one the saddle. The running exponent of a screened kernel (the proposition's commentary): for a Gaussian body of width  $R$  the integrated well energy is  $(C/R) I(\kappa R)$  with  $I(z) = \int_0^\infty u e^{-u^2/4 - zu} du$ , so its exponent is  $s = 1 + z \langle u^2 \rangle / \langle u \rangle$  in that weight:  $1 + \sqrt{\pi} \kappa R + O((\kappa R)^2)$  at a small  $\kappa R$ ,  $3 - 3/(\kappa R)^2 + O((\kappa R)^{-4})$  at a large one, 2.130 at  $\kappa R = 1$  and 2.971 at 10, crossing 2 at  $\kappa R = 0.82$ ; the pair kernel's log-derivative at one separation,  $1 + \kappa R$ , is another number (three derivations). The equation is a theorem (the functional's variational derivative); the branch is computed in the long version's Section 7.3 [1], the budget in S.53; the fence is lattice, the integer version open. Status: theorem for the critical point, the fixed point computed and the continuum constants; the stability along the dilation derived; lattice.

**Derivation S.14** (the invariant and the count's drift, Eq. (7)) Inputs: the one-Node line  $a_{t+1} + a_{t-1} = 2 \cos \omega_t a_t$  with  $\omega_t$  changing slowly. Steps: at fixed  $\omega$  the form  $D_t = a_t^2 - a_{t+1} a_{t-1} = a_t^2 + a_{t-1}^2 - 2 \cos \omega a_t a_{t-1}$  is conserved and equals  $A^2 \sin^2 \omega$  for  $a_t = A \cos(\omega t + \phi)$ ; in the plane  $(a_t, a_{t-1})$  the step is a linear map of determinant 1 whose orbits are the ellipses  $D = \text{const}$ , of area  $\pi D / \sin \omega$ . A slow change of  $\omega$  keeps the area enclosed by the orbit, the action of the map, so  $D / \sin \omega = A^2 \sin \omega$  is the adiabatic invariant to the first order in the slow variation: the discrete form of  $E/\omega$ . Equivalently, the ansatz  $a_t = C \cos(\Theta_t + \phi) / \sqrt{\sin \omega_t}$  with the midpoint phase  $\Theta_t = \sum_{s < t} (\omega_s + \omega_{s+1})/2$ , advancing by  $(\omega_t + \omega_{t+1})/2$  between  $t$  and  $t+1$ , satisfies the line to the second order in  $\omega_{t+1} - \omega_t$  (the phase  $\sum_{s < t} \omega_s$  fails at the first order; two derivations). Check: the recurrence stepped 20,000 intervals with  $\omega$  lowered from 0.841 to 0.600

keeps  $D/\sin\omega$  within  $4 \times 10^{-5}$  of its value while  $D$  falls by a quarter. The drift: the share at a Node in quanta is  $(\Gamma^2/p_i^2) A^2 \sin^2 \omega_b/T$  (Eq. (5) at the Node's paces), so with  $A^2 \sin \omega_b$  fixed,  $e_i(t_2)/e_i(t_1) = [\sin \omega_b(t_2)/\sin \omega_b(t_1)] [p_i^2(t_1)/p_i^2(t_2)]$ , the drift ratio of the paper's Section 4.3, the Node's share of Eq. (5). Planck's relation: the share  $A^2 \sin^2 \omega/T$  at the vacuum's paces is the invariant  $I = A^2 \sin \omega/T$  times  $\sin \omega$ . The body's click: under the declaration that one click is one quantum's change in the absorbing body (Section 2.5), the NodeDetector's own record answers at its rotation by the two-mode line of S.16, one transition per click from the mode  $\omega_i$  to the mode  $\omega_j$ , and the share it takes is  $T(\sin \omega_j - \sin \omega_i) = T \sin(\omega_j - \omega_i) [1 - \omega_i \omega_j/2 + O(\omega^4)]$ , by  $\sin \omega_j - \sin \omega_i = 2 \cos \frac{\omega_i + \omega_j}{2} \sin \frac{\omega_j - \omega_i}{2}$ : one photon's share  $T \sin \omega_L$  at the line  $\omega_L = \omega_j - \omega_i$ , to a relative  $\omega_0^2/2 < 10^{-18}$  at nature's gap (S.27). So a body counts photons, one per quantum since  $2I = 1$  ( $I = 1/2$  per quantum under  $2A^2 \sin \omega = T$ , S.49 (4) of the long version), and reads  $\hbar\omega$  per click with  $T$  for  $\hbar$ ; the share is the energy a click carries, and through a deepening well the number of quanta,  $2I$ , is kept. Status: derived to the first order in the slow variation, the invariant computed; lattice and clicks.

**Derivation S.15** (the clock's click rate, Section 4.3) Inputs: Eq. (3) at  $k = 0$  with  $p_0^2 = \Gamma^2(1 - 2U + 2U^2)$ , the composed paces, to the second order in  $U$ , where they agree with the composed paces of S.2:  $\cos \omega(U) = 1 - g(1 - 2U + 2U^2) = \cos \omega_0 + 2gU - 2gU^2$ . Steps: to first order  $d(\cos \omega) = 2g dU = -\sin \omega_0 d\omega$ , so  $d\omega/\omega_0 = -[2g/(\omega_0 \sin \omega_0)] dU = -f(\omega_0) dU$  with  $f(\omega_0) = 2(1 - \cos \omega_0)/(\omega_0 \sin \omega_0)$ ;  $f = 1.063$  at  $[2, 3]$ , and as  $\omega_0 \rightarrow 0$ ,  $1 - \cos \omega_0 \rightarrow \omega_0^2/2$  and  $f \rightarrow 1$ , the rate  $1 - U$  of the clock's pace alone. The light clock: a packet between two declared faces  $L$  Links apart at the content  $x$  has the round trip  $2L/v_g = 2L\sqrt{3}/N^2$  (Derivation S.2), so its rate falls by  $2x$  to the first order; between two bodies, which stand  $L/h$  Links apart under the one assumption of Section 2.3, the round trip is  $2L\sqrt{3}/N$  (Derivation S.31). Status: derived under the one assumption; clicks.

**Derivation S.16** (the resonance at twice the rotation, Section 4.3) Inputs: the one-Node line with the rotation modulated by a level  $b_L \cos \omega_L t$  read into the content, the law's two-mode line; light's declared read is the holder's rotation (Sections 2.3 and 4.2), under which S.41 derives the transfer line by the two-mode coupling, this derivation's content-modulation line being the pace read's own, with no analogue under the rotation (S.41's Status):  $2 \cos \omega(t) = 2 - 2g(1 - 2x + 2x^2)$  with  $x = x_0 + b_L \cos(\omega_L t)/\Gamma$ , so  $d(2 \cos \omega)/dx = 4g(1 - 2x)$  and  $a_{\text{next}} + a_{\text{before}} = [2 \cos \omega_b + \epsilon \cos(\omega_L t + \vartheta)] a_{\text{now}}$  with  $\epsilon = 4g(1 - 2x_0) b_L/\Gamma$ . Steps: write  $a_t = \text{Re}[z_t e^{i\omega_b t}]$  with  $z$  slowly varying,  $z_{t \pm 1} = z_t \pm z'$ . Then  $a_{t+1} + a_{t-1} = \text{Re}[e^{i\omega_b t} (2 \cos \omega_b z_t + 2i \sin \omega_b z')]$ , and the product  $\epsilon \cos(\omega_L t + \vartheta) \text{Re}[z_t e^{i\omega_b t}]$  has, at  $\omega_L = 2\omega_b$ , the resonant part  $(\epsilon/2) \text{Re}[\bar{z}_t e^{i\vartheta} e^{i\omega_b t}]$ . Equating the slowly varying coefficients of  $e^{i\omega_b t}$ ,  $2i \sin \omega_b z' = (\epsilon/2) e^{i\vartheta} \bar{z}$ , so  $z' = -i\epsilon e^{i\vartheta} \bar{z}/(4 \sin \omega_b)$ ; this antilinear map  $z \mapsto \lambda \bar{z}$  with  $|\lambda| = \mu$  has the real eigenvalues  $\pm\mu$  on two lines of the  $z$  plane, for  $\omega_b$  away from  $\pi/2$  and  $\omega_L$  off every other resonance, where the non-resonant terms average out; at  $\omega_b = \pi/2$  the drive  $\omega_L = \pi$  aliases both sidebands onto the conjugate and the two-step transfer, of trace  $-2 - \epsilon^2 \cos^2 \vartheta$  at the phase  $\vartheta$  of the kick against the mode, grows at  $\text{arsinh}(\epsilon |\cos \vartheta|/2)$  per interval,  $\text{arsinh}(\epsilon/2)$  at  $\vartheta = 0$  and nothing at  $\vartheta = \pi/2$ , twice the formula, so the amplitude grows as  $e^{\mu t}$  along one line and the share, quadratic in the amplitude, as  $e^{2\mu t}$ . Detuned by  $\Delta = \omega_L - 2\omega_b$ , the substitution  $z = \zeta e^{i\Delta t/2}$  gives  $\zeta' = -i(\Delta/2)\zeta - i\mu e^{i\vartheta} \bar{\zeta}$ , whose exponents are  $\pm\sqrt{\mu^2 - \Delta^2/4}$ : growth only for  $|\Delta| < 2\mu$ , a full width  $\epsilon/\sin \omega_b$ . Check: the recurrence stepped 20,000 intervals at  $2 \cos \omega_b = 1.4317$  and  $\epsilon = 0.013$  grows at 0.004653 to 0.004656 per interval over seven modulation phases, the formula's first-order 0.004654 to four digits. The two-mode line: with two bound modes the same modulation multiplies  $a_t = \text{Re}[z_i e^{i\omega_i t} + z_j e^{i\omega_j t}]$ , and its product with  $\cos \omega_L t$  carries the frequencies  $\omega_i \pm \omega_L$ . The

term resonant with the mode  $j$  from the mode  $i$  is at  $\omega_L = \omega_j - \omega_i$ , a transfer between the two modes, and the term that drives both from the conjugate of the other is at  $\omega_L = \omega_i + \omega_j$ , of which the one-mode case above is the degenerate case  $i = j$  (at  $i = j$  the difference line is  $\omega_L = 0$ , a static shift and no line). The transfer's rate,  $\Omega_R = \epsilon_{ij}/(4\sqrt{\sin\omega_i \sin\omega_j})$ , follows by the same slowly varying coefficients with the two modes'  $2i \sin\omega_i z'_i$  and  $2i \sin\omega_j z'_j$  coupled through  $\epsilon_{ij}$ , a Rabi oscillation. Status: two derivations; derived; clicks.

**Derivation S.17** (the cross current of two standing records, Section 4.4) Inputs: two standing records  $a_t = A \cos(\omega_1 t + \alpha)$  and  $b_t = B \cos(\omega_2 t + \beta)$  at the two ends of a Link. Steps: the current  $\text{num}(a_t b_{t-1} - a_{t-1} b_t)$  is  $\text{num}AB[\cos(\omega_1 t + \alpha) \cos(\omega_2 t - \omega_2 + \beta) - \cos(\omega_1 t - \omega_1 + \alpha) \cos(\omega_2 t + \beta)]$ , a sum of cosines of  $(\omega_1 - \omega_2)t$  and  $(\omega_1 + \omega_2)t$  with the constant parts  $\frac{1}{2} \cos(\alpha - \beta + \omega)$  and  $-\frac{1}{2} \cos(\alpha - \beta - \omega)$  only when  $\omega_1 = \omega_2$ ; for  $\omega_1 \neq \omega_2$  its mean over  $n$  beats is 0 to the order of the fast term over the window,  $1/((\omega_1 + \omega_2)nT_b)$  with  $T_b$  the beat's length, so no quantum passes at the first order. The reach: two bodies' tails are  $a_1 e^{-\kappa_b d_1}$  and  $a_2 e^{-\kappa_b d_2}$ ,  $\kappa_b$  the bodies' own inverse reach at  $\omega_b$  ( $\cosh \kappa_b = 3 \cos \omega_b / \cos \omega_0 - 2$ ), at a Link  $d = d_1 + d_2$  Links from the two centres, so the current there is of the order  $a_1 a_2 e^{-\kappa_b d}$  per interval and falls under one quantum's measure of a current for a gapped family,  $W_c \sin \omega_0$ , at  $d = \ln(\text{num} a_1 a_2 / (3 \text{den} T \sin \omega_0)) / \kappa_b$ . Status: derived; lattice.

## The forms of nature (Section 5.1)

**Derivation S.18** (the fall and the Newtonian potential the orbits measure, Eq. (8)) Inputs: Eq. (3) at a content  $U$  for a reader at the weight 1,  $\cos \omega(k, U) = 1 - g(1 - 2U + 2U^2) - \frac{\text{num}}{3 \text{den}}(1 - 2U)^2(1 - \cos k)$  along the motion, the quadratic paces to the first order in  $U$ , where they agree with the composed paces of S.2. Steps: at small  $k$  and  $U$ ,  $\omega(k, U) = \omega_0(U) + k^2/(2m^*)$  with the inertia  $m^* = 3 \text{den} \sin \omega_0 / \text{num} = 3 \tan \omega_0$  (from  $\sin \omega_0 d\omega = \frac{\text{num}}{3 \text{den}} k dk$ ), so the group velocity is  $v = k/m^*$ . The rays of  $\omega(k, x)$ ,  $x$  the coordinate here, give  $dk/dt = -\partial\omega/\partial x = -(\text{den} \omega_0 / dU) \nabla U = (2g/\sin \omega_0) |\nabla U|$  toward the well (Derivation S.15), so the acceleration is

$$a = \frac{1}{m^*} \frac{dk}{dt} = \frac{\text{num}}{3 \text{den} \sin \omega_0} \cdot \frac{2g}{\sin \omega_0} |\nabla U| = \frac{2 \cos \omega_0 (1 - \cos \omega_0)}{3 \sin^2 \omega_0} |\nabla U| = \frac{2 \cos \omega_0}{3(1 + \cos \omega_0)} |\nabla U|,$$

which in the gap  $g = 1 - \cos \omega_0$  is  $[(1 - g)/(3(1 - g/2))] |\nabla U|$ : 0.2667 at  $[2, 3]$  and  $1/3$  as the gap closes. Nature's fall is  $a = c^2 |\nabla U_K|$  with  $U_K = G_K M / (c^2 r)$  and  $c^2 = 1/3$  here, so  $U_K = [2 \cos \omega_0 / (1 + \cos \omega_0)] U$ , Eq. (8), 0.800 at  $[2, 3]$ . Status: derived; clicks.

**Derivation S.19** (the post-Newtonian parameters against the Newtonian potential the orbits measure, Section 5.1) Inputs: Eq. (8),  $U = \lambda U_K$  with the ratio  $\lambda = U/U_K = (1 + \cos \omega_0)/(2 \cos \omega_0) = (\text{den} + \text{num})/(2 \text{num})$  for  $0 < \text{num} \leq \text{den}$  ( $\cos \omega_0 > 0$ , the map  $U_K/U = 2 \text{num}/(\text{den} + \text{num})$  inverted; at  $\text{num} = \text{den}$  the parameters are the limit 1, 1, 1 with no bound orbit; at  $[0, 1]$   $U_K = 0$  and the column is empty; a negative numerator defines none); the composed clock's metric  $N^2 = e^{-2U}$ ,  $h^2 = e^{2U}$  (S.34); the three forms in  $U$ : the bending  $4U_b$ , the periapsis  $6\pi U_1/(1 - e^2)$  with  $U_1 = \mu/a$  (the post-Newtonian formula, standard [3]), the redshift  $f(\omega_0)U$ . Steps: (the orbit step) a slow packet of matter moves by the geometric optics of its band at the composed paces, whose rays are the geodesics of the metric  $N^2 = e^{-2U}$ ,  $h^2 = e^{2U}$  in the time rescaled by  $1/\cos(\omega/2)$ , the band being Klein-Gordon in  $s = 2 \sin(\omega/2)$  (S.31); at the computed pair the rescaling moves Kepler's  $G_K M$  between orbits by  $\tan^2(\omega_0/2)U = 0.2U$ , so  $\gamma_K$  and  $\beta_K$  against  $U_K$  are defined once the calibrating orbit is named; at S.54's bound,  $\omega_e \leq 9.6 \times 10^{-15}$ , the rescaling  $\tan^2(\omega_e/2)$  is

1 to  $10^{-29}$  and the rays are the geodesics; its expansion,  $g_{00} = -(1 - 2U + 2U^2)$  to the second order,  $g_{ab} = (1 + 2U)\delta_{ab}$  to the first and  $g_{0a} = 0$  for the static source (the odd part the holder's odd lines carry, sourced by the count's flux, S.56), the orders the periapsis reads, is the post-Newtonian metric with  $\gamma = 1$  exactly and  $\beta = 1$  in  $U$  at the small gap, so the periapsis advances per orbit by Einstein's  $6\pi U_1/(1 - e^2)$  with  $U_1 = \mu/a$ , that is  $6\pi\mu/(a(1 - e^2))$ . With the two ends' product  $p_0(i)p_0(j)/\Gamma$  for the Link the exterior equation is  $\nabla \cdot (e^{-4U}\nabla U) = 0$ , so  $e^{-4U} = 1 - 4m/r$ ; then  $U = u + 2u^2 + O(u^3)$  with  $u = m/r$  and the clock's  $N^2 = e^{-2U} = 1 - 2u - 2u^2$ , so  $\beta_K = -1$  against  $-g_{00} = 1 - 2u + 2\beta u^2$ , and the advance  $(2 + 2\gamma_K - \beta_K)/3 = 5/3$  of Einstein's, 72 arcseconds per century against 43 (the paper's Section 2.3). (The three parameters against  $U_K$ .) the bending is  $4\lambda U_{K,b} = 2(1 + \text{den}/\text{num})U_{K,b}$ , and the post-Newtonian bending is  $(1 + \gamma)2GM/(c^2b) = (1 + \gamma)2U_{K,b}$ , so  $1 + \gamma_K = 1 + \text{den}/\text{num}$ ,  $\gamma_K = \text{den}/\text{num}$ . The periapsis is  $\lambda$  times Einstein's  $6\pi U_{K,1}/(1 - e^2)$ , and the post-Newtonian advance is  $(2 - \beta + 2\gamma)/3$  of Einstein's, so  $(2 - \beta_K + 2\text{den}/\text{num})/3 = (\text{den} + \text{num})/(2\text{num})$ , whence  $\beta_K = 2 + 2\text{den}/\text{num} - 3(\text{den} + \text{num})/(2\text{num}) = (\text{den} + \text{num})/(2\text{num})$ . The redshift is  $f(\omega_0)\lambda U_K$ , and  $f\lambda = \frac{2(1 - \cos \omega_0)}{\omega_0 \sin \omega_0} \cdot \frac{1 + \cos \omega_0}{2 \cos \omega_0} = \frac{\sin^2 \omega_0}{\omega_0 \sin \omega_0 \cos \omega_0} = \frac{\tan \omega_0}{\omega_0}$ , so the redshift's post-Newtonian parameter  $\alpha_K$ ,  $z = (1 + \alpha_K)U_K$ , has  $1 + \alpha_K = \tan \omega_0/\omega_0$ . At [2, 3]: 1.5, 1.25 and 1.329; all three tend to 1 as  $\omega_0 \rightarrow 0$ , and  $\gamma_K - 1 < 2 \times 10^{-5}$  requires  $\text{den}/\text{num} - 1 < 2 \times 10^{-5}$ . The push on a moving record, of which the bending is the  $\beta = 1$  case, is derived from the same dispersion in S.51; the orbit step above holds for the quadratic band, and at the computed pair the band's  $k^4$  term, of the same post-Newtonian order, adds 1/8 of the advance to first order for an orbit in a coordinate plane and 1/12 in the (111) plane, 13 to 15 and 8 to 10 percent by the ray integration at the orbits quoted, the spread the second order's in  $\mu/p$ , the band's fourth-order term of S.21 against its quadratic one, computed on the exact band twice ( $\mu/p$  from  $5 \times 10^{-4}$  to  $1.1 \times 10^{-2}$ ,  $p$  the orbit's semi-latus rectum in Links,  $e$  from 0.25 to 0.33; the pure-axis coefficient's 17 percent realised by no orbit), vanishing as  $2(1 - \cos \omega_0)/\cos \omega_0$  (two recomputations on the exact band). Status: derived under the one assumption; clicks; the computed pair's values lattice, the ray integrations computed.

**Derivation S.20** (the bending of light and Shapiro's delay, Section 5.1) Inputs: *light's index*  $n = 1 + 2U$  with  $U = \mu/r$  (Derivation S.2). Steps: (a) The delay. The delay along a straight path at the impact parameter  $b$  from  $-x_1$  to  $x_2$  is  $\int (n - 1) dl/c = 2\sqrt{3}\mu \int_{-x_1}^{x_2} dl/\sqrt{l^2 + b^2} = 2\sqrt{3}\mu [\text{arsinh}(x_2/b) + \text{arsinh}(x_1/b)]$ , which for  $x_1, x_2 \gg b$  is  $2\sqrt{3}\mu \ln(4x_1x_2/b^2)$ : Shapiro's delay with  $\gamma = 1$  in  $U$ , and with  $\gamma_K$  against  $U_K$  by the same factor as the bending, since the bending is the transverse derivative of the same integral. (b) The bending. With  $n - 1 = 2U = 3s/(2\pi\Gamma r)$ , where  $s$  is the source's quanta per interval over gravity's divisor  $E_g$  (S.25),  $r^2 = b^2 + l^2$ , the transverse gradient is  $\partial n/\partial b = -3sb/(2\pi\Gamma(b^2 + l^2)^{3/2})$ , and  $\int_{-\infty}^{\infty} b dl/(b^2 + l^2)^{3/2} = 2/b$ , so the turn is  $3s/(\pi\Gamma b) = 4U_b$  with  $U_b = 3s/(4\pi\Gamma b)$ , the index's  $2U$  giving twice the Newtonian  $2U_b$ : Eq. (9) of the paper. Status: derived; clicks; the bending the one assumption's calibration and no test.

**Derivation S.21** (the moving clock to the second order, Section 5.1) Inputs: *the exact band of a free record of the pair* [num, den] *along an axis, from Eq. (1) at the vacuum's pace (S.1):*  $\cos \omega = c_s^2(2 + \cos k)$  with  $c_s^2 = \text{num}/(3\text{den})$ ,  $\cos \omega_0 = 3c_s^2$ ; *the rotation at the moving centre,*  $\Omega(k) = \omega(k) - k\omega'(k)$ , *the phase rate less  $k$  times the group velocity*  $v = \omega'(k)$ , *taken as what a clock carried by the packet reads (a definition of the packet's clock, the second order a definition of  $c_m$ ).* Steps: write  $s = \sin \omega_0$ ,  $c = \cos \omega_0$  and  $u = 1 - \cos k = k^2/2 - k^4/24 + \dots$ ;

then  $\cos \omega = c - c_s^2 u$ , and with  $\omega = \omega_0 + \delta$ ,  $\cos \omega = c - s\delta - c\delta^2/2 + \dots$ , so  $s\delta + c\delta^2/2 = c_s^2 u$  and

$$\omega(k) = \omega_0 + \alpha k^2 - \beta k^4 + O(k^6), \quad \alpha = \frac{c_s^2}{2s}, \quad \beta = \frac{c_s^2}{24s} + \frac{c c_s^4}{8s^3}.$$

Hence  $v = \omega' = 2\alpha k - 4\beta k^3$ ,  $\Omega = \omega - k\omega' = \omega_0 - \alpha k^2 + 3\beta k^4$ , and inverting  $v(k)$ ,  $k^2 = v^2/(4\alpha^2) + \beta v^4/(4\alpha^5) + \dots$ , so

$$\frac{\Omega}{\omega_0} = 1 - \frac{v^2}{4\alpha\omega_0} - \frac{\beta v^4}{16\alpha^4\omega_0} + O(v^6) = 1 - \frac{v^2}{2c_m^2} - \frac{2\omega_0}{\sin 2\omega_0} \cdot \frac{v^4}{8c_m^4} + O(v^6),$$

$$c_m^2 = 2\alpha\omega_0 = \frac{c_s^2 \omega_0}{s} = \frac{\omega_0}{3 \tan \omega_0},$$

where the fourth-order coefficient is  $\beta/(16\alpha^4\omega_0)$  over Lorentz's  $1/(8c_m^4)$ , which reduces with  $3c_s^2 = c$  to  $\omega_0(s/c + c/s) = \omega_0/(\sin \omega_0 \cos \omega_0) = 2\omega_0/\sin 2\omega_0$ . So the clock of a free quantum is Lorentz's factor at the family's own light speed  $c_m$  to the second order in  $v$  exactly, and at the fourth order, along an axis, it carries the factor  $2\omega_0/\sin 2\omega_0$ , 1.69 at [2, 3] and 1 as the gap closes; along the unit vector  $\mathbf{n}$  the band's  $k^4$  term carries  $\sum_a n_a^4$ , so the factor is  $\omega_0[(\sum_a n_a^4) \tan \omega_0 + \cot \omega_0]$ , 1.2224 along a face diagonal and 1.0657 along the body diagonal, where the group velocity is parallel to  $\mathbf{k}$ : the full factor is reached only in the limit of a small rest rotation, the same limit in which the post-Newtonian parameters reach 1 (S.19). Check: with the exact band at [2, 3] the residual  $(f_v - 1 + v^2/(2c_m^2))/v^4$ ,  $f_v = \Omega/\omega_0$ , at  $v = 0.02$  is  $-3.37$  along an axis against 1.6926 times Lorentz's  $-1.988$ , that is  $-3.365$ , and  $-2.12$  along the body diagonal against  $1.0657 \times (-1.988) = -2.119$ . A body: a bound body's centre moves on the body's own band,  $\omega_b(k)$ , whose rest rotation and curvature are computed from the fixed point (S.13); the same steps on that band give the same form with the body's  $c_b^2 = \omega_b/m_b^*$ ,  $m_b^*$  its band inertia and  $c_b$  its kinetic scale as  $m^*$  and  $c_m$  are the free packet's, and the body's band is a computed input, not derived in closed form here. The largest group speed: on the full band  $v_g$  is largest at  $\mathbf{k} = (\pi/2, \pi/2, \pi/2)$ ,  $(\text{num}/\text{den})/\sqrt{3} = \cos \omega_0/\sqrt{3}$ , light's speed times  $\cos \omega_0$ , 0.3849 at [2, 3]. Status: derived, the second order a definition; lattice and clicks.

**Derivation S.22** (de Broglie's relation, Section 5.1) Steps: from Derivation S.18,  $v = k/m^*$  with  $m^* = 3 \tan \omega_0$  at the vacuum's content, so  $k = m^* v = \omega_0 v/c_m^2$  for a free quantum ( $\omega_b$ , a body's own rotation, in its place on a body's band, S.21) with  $c_m^2 = \omega_0/m^* = \omega_0/(3 \tan \omega_0)$ , the family's own light speed; 0.2508 at [2, 3]. Status: derived; lattice and clicks.

## The two slits (Section 6)

**Derivation S.23** (the fringe's spacing and the near-field minima, Section 6.2) Inputs: the gaps centred at the rows 18 and 30,  $d = 12$ ; the screen  $L = 24$  Links from the wall;  $\lambda = 8$  Links; gravity's weight  $E_g = 10^6$  in this universe, chosen so that the packet's own gravity does not focus it. Steps: the far-field spacing is  $\lambda L/d = 16$  Links. In the near field the minima are where the two gaps' paths to the screen differ by half a wavelength,  $\sqrt{L^2 + (y - 18)^2} - \sqrt{L^2 + (y - 30)^2} = \pm 4$ , whose solutions are  $y = 15.28$  and  $y = 32.72$ : the rows 15 and 33. An opaque region in one gap takes that gap's inflow, so the screen reads the other gap's envelope and no fringe, by the continuity of the one inflow through the two boundaries. Check: the two roots computed from the equation above; the law's real line on this universe gives the exact lay 1,810.4 quanta by Eq. (5) and 284.7 units through the screen over the window (the line's 284.7 was computed after the run; the one number that stood before the run is the blind row's 273), and 278.9 on the integer lay of 1,809 quanta (the paper's Section 6.2); the integer lay's 278.9 lies below

the exact lay's 284.7 because the before level's mismatch at the lay seeds an opposite-branch admixture that the receding face loses (the two roots  $e^{\pm i\omega}$  of the leapfrog line, the backward branch's amplitude from the before level's difference, 18.80 at most, and its exit through the receding face); the folded lattice returns the Node to itself through both  $z$  Ports, so the band is  $2 \cos \omega = \frac{2}{3}(\cos k_x + \cos k_y + 1)$  exactly at  $k_z = 0$ ; the peak of the screen's inflow is at the interval 69.0 on the lattice as declared, 62.0 with the lattice refined fourfold and 62.1 eightfold (the line evaluated); the gap NodeDetector's net inflow is  $-0.58$  over the window; the run's record is deposited with the paper (Statements and Declarations) and holds the two seeds' rows. The blind row's 273 is light's line in real arithmetic propagated on the design's lattice ( $64 \times 48$  Nodes with every face open, 1,857 quanta and the window 45 to 120) from the two gaps as its sources and summed at the screen over the window, Huygens's construction with the lattice's own line as its propagator, which the record names the Huygens sum; the line stepped over the whole lattice from the laid packet is a second computation, 277.8 at the meeting round's lay and 284.7 at this universe's exact lay; the run of Rule3's integer line on this universe gives  $N = 285$  at both seeds, 24 and 25, the seed-24 clicks' sum of absolute differences from the blind row 28 of 275 (275 the blind row's sum by its per-Node rounding, 273 its  $N$ ); the group velocity  $\sin k/(3 \sin \omega) = 0.547$  Link per interval at  $k = \pi/4$  and  $\omega = 0.4456$ , the flight over  $L' = 34$  Links from the packet's centre (the column 10) to the screen 62.2 intervals; the rows 15 and 33 fall in the four-row regions 3 and 8 counted from 0; the blind row's dips at the regions 4 and 8; the reversal check of Section 2, the run forward and back over 40 intervals with the counting window opening at 45, so that the check contains no click. Status: derived for the spacing and the minima; computed for the lays, the totals and the arrival; the run's numbers computed, a run; the universe a declaration; lattice and clicks as each sentence marks.

## The forms, the bodies and the calibration (Sections 4 to 8)

**Derivation S.24** (the Link's bound from light's dispersion, Section 5.1) Inputs: the group speed  $v_g = (1/\sqrt{3})(1 - k^2/12)$  (Derivation S.1); the bound on a quadratic dependence of light's speed on its energy, MAGIC's subluminal quantum-gravity (QG) scale of quadratic Lorentz violation  $E_{\text{QG},2}$  above  $6.3 \times 10^{10}$  GeV (Acciari et al. 2020 [8]; the lattice band is subluminal,  $v_g = (1/\sqrt{3})(1 - k^2/12)$ , so the subluminal limit applies), that is  $|\delta v/v| = (3/2)(E/E_{\text{QG},2})^2 < 3.8 \times 10^{-16}$  at  $E = 1$  TeV in the standard quadratic form. Steps:  $k^2/12 < 3.8 \times 10^{-16}$  gives  $k < 6.8 \times 10^{-8}$  per Link at 1 TeV; with  $k = 2\pi \text{Link}/\lambda$  and  $\lambda = 1.24 \times 10^{-18}$  m at 1 TeV, the Link is below  $k\lambda/(2\pi) = 1.3 \times 10^{-26}$  m along an axis, where the  $k^4$  coefficient of the band's  $\omega^2$  is  $-1/54$  ( $-1/135$  averaged over directions,  $2.1 \times 10^{-26}$  m; 0 along a diagonal, no bound); over the unit of S.27,  $6.69 \times 10^{-13}$  m, the axis bound gives  $\omega_e < 2.0 \times 10^{-14}$  ( $\omega_e$  the matter family's rest rotation, S.27's  $\omega_0$ ) ( $3.1 \times 10^{-14}$  on the average) and  $\gamma_K - 1 = \omega_e^2/2 < 2.0 \times 10^{-28}$ . The orientation: each source's bound carries its direction's coefficient  $C(\mathbf{n}) = (3 \sum_a n_a^4 - 1)/24$  of the group speed's fall, unknown against the lattice:  $1/12$  on an axis,  $1/48$  on a face diagonal, 0 on a body diagonal and continuous toward it; the mean over directions,  $\langle \sum_a n_a^4 \rangle = 3/5$ , gives  $2/5$  of the axis's coefficient (the  $-1/135$  against  $-1/54$  above); the Link bound weakens tenfold only within 4 degrees of a body diagonal, one percent of the sky, and a hundredfold within 0.4 degrees, so the comparison of MAGIC's and LHAASO's bounds holds under the mean's premise (two derivations, the second S.54's two-way coefficient check). Status: derived; lattice and clicks.

**Derivation S.25** (Newton's constant as a reading, Eq. (11)) Inputs: *the massless holder's rest far from a static source of  $M$  quanta per interval,  $3M/(4\pi r)$  over the divisor  $E_g$ , the write  $s = M/E_g$ ; the clock's content  $U = c_\ell/\Gamma$ ,  $c_\ell$  the content in levels. Steps:  $U = 3M/(4\pi r \Gamma E_g) = 3s/(4\pi \Gamma r)$ , and nature's  $U = GM/(c^2 r)$  with  $M$  in quanta and  $c^2 = 1/3$  gives  $G/c^2 = 3/(4\pi \Gamma E_g)$ , that is  $G_{\text{clock}} = 1/(4\pi \Gamma E_g)$ ; Kepler's  $G$ , read from the orbits of matter, is  $[2 \cos \omega_0 / (1 + \cos \omega_0)] G_{\text{clock}}$  by Eq. (8). Status: derived; lattice, clicks.*

**Derivation S.26** (the fine-structure constant as a coefficient of the declared universe, Section 5.1) Inputs: *the holder of the sign under its rotation (Section 3.5) with its declared write weight  $k$ , the written level  $k$  times the plain one and the turn  $\theta = \text{level}/\Gamma$ ,  $k = 1$  in the rule's declared universe; a circular record of a family at the vacuum's paces,  $z = Ae^{i\omega_s t}$ , the quantum of a family one unit of the invariant,  $2A^2 \sin \omega_s = T$ , with the share  $T \sin \omega_s$  and  $|W| = A^2 \sin \omega_s = T/2$  (S.9); the holder's rest far from a static source of  $s$  per interval,  $3s/(4\pi r)$  over its weight  $E_s$  (Derivation S.25's line); de Broglie's relation and light's  $c = 1/\sqrt{3}$  (S.1, S.22). Steps: (a) One quantum writes  $s = W/(E_s T) = 1/(2E_s)$  per interval, for every family, since the sourcing family's denominator enters no write (the count is the share over the wall  $3 \text{den}_s T$  and the source the form over  $E_s T$ , the same number); so  $|q_p| = |q_e|$  with nothing declared. The angle at  $r$  is  $\theta(r) = 3ks/(4\pi r \Gamma) = 3k/(8\pi r \Gamma E_s)$ . (b) A reader of unit sense gains the angle per interval, so its wave number changes by the gradient,  $dk_{\text{dB}}/dt = \mp \nabla \theta$  by the sense of (c), of magnitude  $3k/(8\pi r^2 \Gamma E_s)$ ; Coulomb's law in nature's units,  $dk_{\text{dB}}/dt = \alpha c/r^2$ , gives  $\alpha_{\text{law}} = \sqrt{3} r^2 dk_{\text{dB}}/dt = (3\sqrt{3}/8\pi) k/(\Gamma E_s) = 0.20675 k/(\Gamma E_s)$ ; the holder's weight enters the source and the read alike as  $k$  and  $E_s$ , and the family's gap cancels. (c) The reader's shift: the step carries  $e^{i\theta_t}$  on  $z_{\text{next}}$ , so a sense + quantum goes from  $\omega$  to  $\omega + \theta$  at its fixed Wronskian (S.33, S.41 (b)), its energy from  $T \sin \omega$  to  $T \sin(\omega + \theta)$ ,  $\delta E = +T\theta \cos \omega$  per quantum; a body of  $n$  quanta and sense  $\pm$  shifts by  $\pm n T \theta \cos \omega = 2\theta \cos \omega W$  with  $W = \pm n T/2$ , the force  $-2W \cos \omega \nabla \theta = \mp n T \cos \omega \nabla \theta$ , which is  $\mp n \nabla \theta$  by the sense in the unit  $T \cos \omega$ , one quantum's energy shift per unit turn (its energy  $T \sin \omega$ ), at the vacuum's rotation, Coulomb's magnitude with  $q = \pm ne$  and  $e = T/2$ ; the angle  $\theta = k_r L/\Gamma$  carries the sign of the level and the level the sign of its source's Wronskian, so two quanta of one sense raise each other's rotation,  $\delta E > 0$  growing toward the source, and move apart; which sense moves apart is the declared sign of Section 3.5 (under the law's one convention like senses move apart, as nature's measurement that like charges repel declares), the magnitude untouched; the reader sources the holder it reads at the same weight, so the pair is reciprocal (the reciprocity Theorem 3 needs). (d) The number: nature's  $\alpha = 1/137.036$  fixes  $k/(\Gamma E_s) = 1/28.33$  and nothing else;  $k$  is the universe's integer read from this one click as  $E_g$  is from  $G$ , and at  $k = 1$  no integer  $\Gamma E_s$  gives it (28 gives  $1/135.43$  and 29 gives  $1/140.27$ ). (d') The energy line. The weight is the product of the write's and the read's,  $k = k_w k_r$ , with the read's  $k_r = 1$  on the integer lattice (a read weight would multiply a level the write already rounded away). Under the time-Link turn a plane record's form and Wronskian are conserved exactly per Node in the rational line for any level in time (the turn the time Link's gauge, S.33, S.41 (a)) and its rotation shifts by the turn per interval, so the energy its clicks read changes by  $T[\sin(\omega + \sigma \sigma_s \Delta \theta) - \sin \omega]$  per quantum of sense  $\sigma$  in a level of source sense  $\sigma_s$  (row 9 of the law's readings [9]: a quantum goes to  $\omega + \sigma \sigma_s |\theta|$ ),  $\sigma \sigma_s T \cos \omega \Delta \theta$  to first order in the turn (the remainder  $(\Delta \theta/2) \tan \omega_s$  relative, below  $10^{-18}$  at nature's turn per interval), per quantum  $T \cos \omega_s (k_r/\Gamma) \Delta \ell$  in a level of its own sense (a convention with one turn angle on both levels would lose  $2W \cos \omega \Delta \theta$  of form per Node per step; the law's convention loses none); the holder's own form changes by  $\sum_i s_i (\ell_{t+1} - \ell_{t-1})$  per step under the write, per quantum  $(k_w/E_s) \Delta \ell$ , the two factors 2 cancelling, and for a pair of opposite senses the cross form carries  $W_1 W_2 < 0$ ; so the energy its clicks read and the holder's form move together under the turn and the write, both up for a*

body in its own level and both down for a pair of opposite senses, and agree to first order in the turn if and only if  $k_r E_s T \cos \omega_s = k_w \Gamma$ : one energy read twice, by the clicks and by the form, an exact condition on the energies read and not an identity of any one record's form, the ratio of the two changes  $\chi = k_r E_s T \cos \omega_s / (k_w \Gamma)$  at every step and for every motion, at the vacuum's paces of these inputs (in a well the body's clock enters the turn and the write alike), the ratio  $+\chi$  under row 9 by the algebra above (the two senses' product the sign); the law's line is a condition on the energies read and no identity of a record's form. Under it, with  $k_r = 1$ ,  $\alpha_{\text{law}} = (3\sqrt{3}/8\pi) T \cos \omega_s / \Gamma^2$ , at nature's gap  $T = \Gamma^2/28.33$ ,  $E_s$  a unit of the level and the declared universe one integer fewer; the line is per family through  $\cos \omega_s$ , which in nature is 1 to  $10^{-18}$  for every family, so one write weight serves all; light and every real-line family read no turn and stand outside it. The rule's own universe has no plane family and keeps  $T = 2^{15}$ ; the universe's gate condition,  $E_s T \text{ num} = k_w \Gamma \text{ den}$  for a plane family reading a holder under the rotation, enforces the line. The per-step identity above is the one-field case, a level built by the body itself; under the rule in force, no record reading its own write, the reader's loss is by the other record's field and the writer's gain its own field's, and what holds per step is the two-record balance at the same coefficient per quantum, not run; the universe's gate condition is unchanged by this. The radiated share of a charged body (S.45 (c) of the long version): the time level carries half of Larmor's power in form; the odd lines' part depends on their wall and coupling, which of the two couplings the odd lines carry open; no rate and no lifetime is claimed. Two derivations on the identity and its number; the line a decision of the model (assumption). (e)  $\Gamma$  is bounded by no click of the clocks (the gap bound binds the matter family's own denominator,  $\text{den} > 1.5 \times 10^{19}$  for the least pair, S.27, a pair of a universe meant for nature and not Universe24's): the content is an integer and the clock reads it as  $p_0 = \Gamma(1 - 1/\Gamma)^c$ , so the finest level at one Node and one interval is  $1/\Gamma$ ; but the write adds  $(\sum wq + r) \text{ div } E$  with the remainder  $r$  kept, so a sub-unit source does not vanish but writes one level once every  $1/s$  intervals, the level at a Node a sequence of 0 and 1 whose mean is the exact rest  $3s/(4\pi r)$ , a first-order sigma-delta modulation; a clock in such a level reads the level 1 for the fraction  $\Delta U \Gamma$  of its intervals and its mean redshift is  $\Delta U$ , finer than  $1/\Gamma$ , and a bound body spanning many Nodes averages their independent dithers, so the gravitational clocks of nature bound the mean potential and not  $\Gamma$ ; nothing bounds  $\Gamma$  from above and no line fixes it:  $\Gamma$  is free and large (two derivations). Check:  $3\sqrt{3}/(8\pi) = 0.20675$ ;  $137.036 \times 0.20675 = 28.33$ ; the share's formula at [2, 3] and  $\Gamma = 6,000$  gives  $|W|/(\text{count } T) = 0.6708 = 1/(2 \sin \omega_0)$  for a record laid by share ( $A^2 = T/(2 \sin^2 \omega_0)$  for a record laid by share) and  $1/2$  for a record of count 1, one quantum of the family under the record's own unit (S.49 (5) and S.61 row 16 of the long version);  $3\sqrt{3}/(8\pi)/28.33 = 0.20675/28.33 = 1/137.0$ . Status: derived under the declared weights; lattice and clicks.

**Derivation S.27** (the units, the share's ceiling and the matter pair's gap, Section 5.1) Inputs: the electron as the free quantum of matter, rotating at  $\omega_0$  per interval, and nature's electron at  $m_e c^2/\hbar$  per second; light's band  $\cos \omega = 1 - \frac{1}{3} \sum_a (1 - \cos k_a)$  (S.1); a photon of 1.4 PeV, LHAASO's record of 2021 [10] (photons above 2 PeV were reported in 2024 [11]); the share  $T \sin \omega$  a click reads (S.14). Steps: one interval is  $\omega_0 \hbar / (m_e c^2) = \omega_0 \times 1.288 \times 10^{-21}$  s, and one Link, light moving  $1/\sqrt{3}$  Link per interval at  $c$ , is  $\sqrt{3} \omega_0 \bar{\lambda}_e = \omega_0 \times 6.69 \times 10^{-13}$  m with  $\bar{\lambda}_e = \hbar / (m_e c) = 3.862 \times 10^{-13}$  m. The band's top along the body diagonal,  $k_a = \pi$  for all  $a$ , is  $\cos \omega = -1$ ,  $\omega = \pi$ , and along an axis,  $k = \pi$  alone,  $\cos \omega = 1/3$ ,  $\omega = 1.231$ . Along an axis the group velocity vanishes at the edge; along the diagonal,  $k_a = q$  on every axis, the band is  $\omega = q$  exactly, so the speed stays  $1/\sqrt{3}$  up to the corner, where the band has a cusp and no one gradient. That is a lattice statement of the highest rotation, which no click reads as an energy: what a click reads of a quantum of frequency  $\omega$  is its share,  $T \sin \omega$  (S.14), at most

$T$ , reached at  $\omega = \pi/2$ , which the band contains along the diagonal (along an axis it ends at 1.231,  $\sin \omega = 0.943$ ), and an energy measurement of nature, a calorimeter or an air shower, is a sum of transitions' shares. The electron's rest energy is  $T \sin \omega_0$ , so a photon of 1.4 PeV read as a share gives  $\sin \omega_0 \leq 0.511 \text{ MeV}/1.4 \text{ PeV}$ ,  $\omega_0 < 3.7 \times 10^{-10}$  for every orientation of the lattice (the diagonal's  $3.65 \times 10^{-10}$ , the axis's  $3.44 \times 10^{-10}$ ; the diagonal's is the one quoted). Then  $\text{den} = d/(1 - \cos \omega_0) \approx 2d/\omega_0^2 > 1.5 \times 10^{19}$  with  $d = \text{den} - \text{num} \geq 1$  (the bound at  $d = 1$ ) and  $\gamma_K - 1 = 1/\cos \omega_0 - 1 \approx \omega_0^2/2 < 6.7 \times 10^{-20}$  (S.19);  $\Gamma$  reads nothing of  $\omega_0$ ,  $\alpha$  fixing  $k/(\Gamma E_s)$  and not  $\Gamma$  (S.26). Were the frequency read as the energy,  $T\omega$ , the bound would be  $1.1 \times 10^{-9}$  along the diagonal and  $4.5 \times 10^{-10}$  along an axis: the law's resonance adds frequencies (S.41) and its share adds sines (S.14), the two agreeing to  $\omega^2/6$  relative, below  $10^{-18}$  at nature's gap, and parting at the Link scale, where which of the two a click takes the law as written does not decide and no measurement of nature reaches; named, not claimed. Check:  $\hbar/(m_e c^2) = 1.0546 \times 10^{-34}/8.187 \times 10^{-14} = 1.288 \times 10^{-21} \text{ s}$  (the CODATA values [12]);  $\sqrt{3} \times 3.862 \times 10^{-13} = 6.69 \times 10^{-13} \text{ m}$ ;  $0.511/(1.4 \times 10^9) = 3.65 \times 10^{-10}$ ;  $2/(3.65 \times 10^{-10})^2 = 1.5 \times 10^{19}$ ;  $(3.65 \times 10^{-10})^2/2 = 6.7 \times 10^{-20}$ ;  $\sin 1.231 = 0.943$ ;  $[\pi - \omega(\pi - \epsilon)]/(\sqrt{3}\epsilon) = 0.5774$  at  $\epsilon = 10^{-2}$  and  $10^{-3}$  on the diagonal. Status: derived; lattice and clicks.

**Derivation S.28** (the content's index for every family, Section 5.1) Inputs: Eq. (3) at the vacuum's pace and at a content  $U$  under the composed paces,  $(p_0/\Gamma)^2 = N^2$  and  $(p_a/\Gamma)^2 = N^4$  with  $N = (1 - 1/\Gamma)^c = e^{-U}$  to  $O(1/\Gamma)$  (Derivation S.2); a stationary boundary conserves the rotation  $\omega$ . Steps: outside,  $\cos \omega = 1 - g - \frac{\text{num}}{3\text{den}}(1 - \cos k)$  with  $g = 1 - \text{num}/\text{den}$ ; inside,  $\cos \omega = 1 - gN^2 - \frac{\text{num}}{3\text{den}}N^4(1 - \cos k_{\text{in}})$ . Equating,  $\frac{\text{num}}{3\text{den}}N^4(1 - \cos k_{\text{in}}) = \frac{\text{num}}{3\text{den}}(1 - \cos k) + g(1 - N^2)$ , so  $1 - \cos k_{\text{in}} = [1 - \cos k + 3(\text{den} - \text{num})(1 - N^2)/\text{num}]/N^4$ , exact in  $k$  on the domain where the right side lies in  $[0, 2]$ . With  $1 - N^2 = 2U - 2U^2 + \dots$  and  $N^{-4} = 1 + 4U + \dots$  the first order is  $[1 - \cos k + 6(\text{den} - \text{num})U/\text{num}]/(1 - 2U)^2$ , the quadratic forms' line, the mass term's second order  $-6(\text{den} - \text{num})U^2/\text{num}$ . For light,  $g = 0$ :  $1 - \cos k_{\text{in}} = (1 - \cos k)/N^4$ , so  $\sin(k_{\text{in}}/2) = \sin(k/2)/s$  with  $s = N^2$ , exact in the law's variable  $2 \sin(k/2)$  where  $\sin(k/2) \leq s$ . The phase index is  $k_{\text{in}}/k = (2/k) \arcsin[\sin(k/2)/s] = 1/s + k^2(1/s^3 - 1/s)/24 + O(k^4)$ , which is  $1/s = e^{2U}$ , Derivation S.2's long-wavelength index, at  $k \rightarrow 0$  alone. At  $k = \pi/2$  and  $s = 0.8$  the exact ratio is 1.380 against  $1/s = 1.25$ , and to first order in  $U$  the index is  $1 + [4 \tan(k/2)/k]U$ , the coefficient 2 at  $k \rightarrow 0$  and  $8/\pi$  at  $k = \pi/2$ : the lattice's own chromatic dispersion, absent in the ray limit in which the bending, the delay and the lens are read. For slow matter,  $k, k_{\text{in}} \ll 1$ :  $k_{\text{in}}^2 = [k^2 + 6(\text{den} - \text{num})(1 - N^2)/\text{num}]/N^4$ , so  $n_m^2 = k_{\text{in}}^2/k^2 = [1 + 6(\text{den} - \text{num})(1 - N^2)/(\text{num}k^2)]/N^4$ , to first order  $[1 + 12(\text{den} - \text{num})U/(\text{num}k^2)]/(1 - 2U)^2$ . Check: with  $k^2 = m^{*2}v^2$  and  $m^* = 3 \tan \omega_0$  (S.18), the  $1/k^2$  term is  $12(\text{den} - \text{num})U/(\text{num}m^{*2}v^2) = [4(1 - \cos \omega_0)/(3 \cos \omega_0 \tan^2 \omega_0)]U/v^2$ , which as  $\omega_0 \rightarrow 0$  is  $(2/3)U/v^2 = 2(U/3)/v^2 = -2\Phi/v^2$  with  $\Phi = -c^2U$ ,  $c^2 = 1/3$  (nature's  $n^2 = 1 - 2\Phi/v^2$  for a slow body in a potential): nature's index for a slow body in a potential. The factor  $6(\text{den} - \text{num})/\text{num}$  over  $m^{*2}$  is the matter band's own, the Kepler column of Eq. (8), and  $1/N^4$  the Link's double read. Status: two derivations; derived; lattice and clicks.

**Derivation S.29** (the clusters' cosmology: the ball of dust, Section 4.4) The paper's claim from this derivation is the first order in  $z$ , Friedmann's dust equations and Hubble's law; the second order is derived here and stands in the paper's Section 8.1 beside the supernovae as a finding against nature. Inputs: Newton's pull between bodies at Kepler's  $G_K$  (Section 5.1, Derivation S.25); a finite ball of dust of radius  $a(t)$  and uniform density  $\rho$  on the static lattice; the Doppler period of a receding source, derived in the steps, and the moving clock's factor (S.21). Steps: a shell at  $a$  feels the mass inside it,  $M(a) = (4\pi/3)\rho a^3$ , and nothing

from outside (Newton's theorem on a sphere, which holds for the  $1/r$  level of Section 5.1), so  $\ddot{a} = -G_K M(a)/a^2 = -(4\pi G_K/3)\rho a$ . Multiplying by  $\dot{a}$  and integrating with  $\rho a^3$  constant gives  $\dot{a}^2/a^2 = (8\pi G_K/3)\rho - K/a^2$  with  $K$  the start's declaration, positive for a bound ball: Friedmann's two equations for dust, as Milne and McCrea derived them for the Newtonian ball [13]. The deceleration parameter is  $q_0 = -\ddot{a}/\dot{a}^2 = (4\pi G_K \rho/3)/H^2 = \Omega_m/2$  with  $\Omega_m = 8\pi G_K \rho/(3H^2)$ , positive for every  $\rho > 0$ . The redshift: a source receding at  $\beta c$  emits two crests a period  $P$  apart from points  $\beta c P$  apart, so the second trails the first by  $(1 + \beta)cP$  and a resting NodeDetector reads the period  $(1 + \beta)P$  exactly, the wave number set at the emission and kept by the line's translation invariance (S.1), the period read  $(1 + \beta)$  times longer and the source's own period is longer by the moving clock's factor  $1/\sqrt{1 - \beta^2}$  at  $c_m = c$  (S.21), so  $1 + z = (1 + \beta)/\sqrt{1 - \beta^2} = 1 + \beta + \beta^2/2 + \dots$ , nature's relativistic Doppler form to the second order, the receding source's classical factor  $1 + \beta$  differing at the second order  $(1/(1 - \beta) = 1 + \beta + \beta^2$  is the receding receiver's). At the computed pair  $c_m = 0.867c$  the second order differs, as every number of the computed pair does. Every click's duration is stretched by the same  $1 + z$ , the light curves' time dilation, and a surface's brightness falls by  $(1 + z)^{-4}$ , Tolman's dimming, at the small gap,  $(1 + z)^{-2}$  from the two rows alone, the aberration's two powers needing the band's Lorentz invariance, which holds at the small gap (S.21) and not at a finite gap (S.4); light from a source at  $r$  falls into the ball's deeper centre, a blueshift  $\Delta U = (\Omega_m/4)(H_0 r/c)^2$  beside the Doppler, the second order that makes  $d_L(z)$  Friedmann's with  $q_0$  unchanged. Status: two derivations; derived; clicks; the second order,  $q_0 = \Omega_m/2 > 0$  under the declared homologous start, a finding against nature in the paper's Section 8.1.

**Derivation S.30** (the shadow's capture radius from light's index, Section 5.1) Inputs: light's index  $n(r) = e^{2U}$  with  $U = m/r$  (Derivation S.2 under the composed paces),  $m = GM/c^2$  in Links (the  $\mu$  of S.19 and S.20). Steps: in a static isotropic index a ray at the impact parameter  $b$  keeps  $n(r)r \sin \psi = b$  along its path, so a circular ray stands where  $d(nr)/dr = 0$ :  $d(re^{2m/r})/dr = e^{2m/r}(1 - 2m/r) = 0$  at  $r = 2m$ , the photon sphere, and the capture radius is  $b = n(2m)2m = 2em = 5.437m$ . Schwarzschild's is  $3\sqrt{3}m = 5.196m$ , so the shadow is larger by  $2e/(3\sqrt{3}) - 1 = 0.046$ . Comparison: with the quadratic index  $1/(1 - 2U)$  the same condition gives  $r = 4m$  and  $b = 8m$ , +54 percent, outside the Event Horizon Telescope's fractional deviation from Schwarzschild for Sgr A\*,  $-0.08 \pm 0.09$  [14]; the composed index's +4.6 percent sits 1.4 standard deviations from it, bounded and not confirmed. Status: derived under the one assumption; clicks; the shadow bounded by measurement and not confirmed.

**Derivation S.31** (the two clocks under the one assumption of Section 2.3, Section 5.1) Inputs: the one assumption of Section 2.3, a uniform level  $U$  a change of units, time by  $N$  and length by  $h$  with  $hN = 1$ ; light's speed  $N^2$  (S.2). Steps: a bound body in the level is the flat body in the local units, so its size is  $1/h = N$  Links by the one assumption. Its band at the Node is  $2 - 2\cos \omega = 2(1 - \nu)N^2 + (\nu/3)N^4 \sum_a s(k_a)^2$  with  $\nu = \text{num}/\text{den}$  and  $s(x) = 2\sin(x/2)$ , since  $(p_0/\Gamma)^2 = N^2$  and  $(p_a/\Gamma)^2 = N^4$ , and its lattice wave numbers are  $k = k_{\text{proper}}/N$ , so  $(\nu/3)N^4 \sum_a s(k_a)^2 = (\nu/3)N^2 \sum_a k_{\text{proper},a}^2 [1 - k_a^2/12 + \dots]$ : every mode of the record, the rest rotation and every level of its spectrum, scales by  $N$  in  $s(\omega)$ , exactly at long wavelength and to the lattice's  $k^2/12$  at finite  $k$ . In  $\omega$  the rest line's rate is  $-2\tan(\omega_0/2)/\omega_0 = -f(\omega_0)$  per unit  $U$ , Derivation S.15's number. A difference  $s(\omega'_2) - s(\omega'_1)$  scales by  $N$  exactly and is no observable. The temporal beat a NodeDetector reads,  $\Delta' = \omega'_2 - \omega'_1$  in  $\cos(\Delta't)$ , with  $\omega' = 2\arcsin[N\sin(\omega/2)] = N\omega [1 - (1 - N^2)\omega^2/24 + O(\omega^4)]$ , is  $\Delta' = N\Delta [1 - (1 - N^2)(\omega_1^2 + \omega_1\omega_2 + \omega_2^2)/24 + O(\omega^4)]$ : at  $N = 0.8$ ,  $\omega_1 = 0.4$  and  $\omega_2 = 0.9$  the exact beat is 0.3916 against  $N\Delta = 0.4$  and the series 0.3920. The relative correction,

$(1 - N^2)(\omega_1^2 + \omega_1\omega_2 + \omega_2^2)/24$  for the beat,  $(1 - N^2)\omega^2/8$  at  $\omega_1 = \omega_2$ , is  $2.5 \times 10^{-38}$  at  $\omega = 1.0 \times 10^{-14}$  and  $U = 10^{-9}$ , an illustration below the bounds  $(1.0 \times 10^{-37}$  at the axis bound's  $2.0 \times 10^{-14}$ , S.24;  $3.4 \times 10^{-29}$  at the LHAASO bound's  $3.7 \times 10^{-10}$ , S.27) and 1.4 percent at  $\omega_b = 0.78$  and  $U = 0.1$ , the lattice's arithmetic and not the law's. A cavity whose mirrors are bodies stands  $L/h = LN$  Links apart and its round trip is  $2LN\sqrt{3}/N^2 = 2L\sqrt{3}/N$  intervals, the rate  $N$  exactly with no  $\omega$  correction, the atom's beat to its correction  $(1 - N^2)\omega^2/8$  above; a cavity of declared faces keeps its  $L$  Links and ticks at  $N^2$ . So under the one assumption of Section 2.3 no clock type is singled out, which is what the null redshift tests measure, while a cavity of declared faces is a NodeDetector's region and no body of nature. Status: derived under the one assumption; lattice and clicks.

**Derivation S.32** (the conserved form's Link factor, two witnesses, Theorem 3) Inputs: Theorem 3's line without its division on a periodic lattice at the content 0 with every Link's factor  $q$ , so  $p_0 = p_i = \Gamma$ ,  $R = 2 \text{num } q^2$ ,  $w = 6 \text{den } \Gamma^2$  and  $S = 12 \text{num } \Gamma^2 - 6R$ ; the form  $E$  of Eq. (4) and the same form with the plain Link term  $-2 \text{num } \sum_i \text{now}_i S_6(\text{before})_i$ ; exact fractions. First witness: three Nodes along  $x$  with  $y$  and  $z$  folded,  $\text{num} = \text{den} = 1$ ,  $\Gamma = 4$ ,  $q = 2$  ( $R = 8$ ,  $w = 96$ ,  $S = 144$ ),  $\text{now} = (1, 0, 0)$  and  $\text{before} = 0$ : one step gives  $\text{next} = (11/6, 1/12, 1/12)$ ; the plain form goes from 6 to  $-21/4$  and Eq. (4) stays 6. Second witness:  $3^3$  Nodes at  $[2, 3]$ ,  $\Gamma = 10$ ,  $q = 5$  ( $R = 100$ ,  $w = 1,800$ ,  $S = 1,800$ ), a uniform  $\text{now} = 3$  over  $\text{before} = 0$ : the step gives  $\text{next} = 4$  at every Node with the remainder 0, so the integer step and the line agree; the plain form per Node goes from 162 to  $-54$  and Eq. (4) stays 162 (4,374 over the lattice). Check: at  $q = \Gamma$  the factor is 1 and the two forms coincide, the domain of the plain current. Status: theorem (a corollary of Theorem 3); computed for the two witnesses; lattice.

**Derivation S.33** (magnetism: the continuity identity and the covector, Section 5.1) Inputs: a plane record  $z = \text{re} + i \text{im}$  stepping by Rule3 under the holder of the sign (Section 3.5):  $e^{i\theta_t} z_{\text{next}} + e^{-i\theta_{t-1}} z_{\text{before}} = [S z_{\text{now}} + \sum_a R_a (e^{i\theta_a} z_{+a} + e^{-i\theta_a} z_{-a})]/w$ , with  $\theta_t$  this interval's time turn on the level next,  $\theta_{t-1}$  the previous interval's on the level before and  $\theta_a$  the phase on the axis  $a$ 's Links, each read at its interval's start (the sign convention of the law's document [9]); the Wronskian booked from the turned levels  $u_i = e^{-i\theta_{t-1}} z_{i,\text{before}}$  and  $v_i = e^{i\theta_t} z_{i,\text{next}}$  as the step reads them,  $W_i = \text{Im}(\bar{z}_{i,\text{now}} u_i)$ , and  $W_i(t+1) = \text{Im}(\bar{v}_i z_{i,\text{now}})$  after the step, the two differing by the Link fluxes of  $(a)$ ; the Link quantity  $g_{ij} = \text{Im}(\bar{z}_i e^{i\theta_{ij}} z_j)$ , antisymmetric,  $g_{ji} = -g_{ij}$ , since the phase is read  $+\theta_a$  through the  $+a$  Port and  $-\theta_a$  through the  $-a$  Port. Steps: (a) The identity (the level before under the previous interval's angle and the level next under this one's, so the Wronskian is conserved exactly for any level in time, a uniform level a gauge, S.41 (a); the algebra below is written at one angle  $\theta_0$  over the step, the accumulated phase removing a varying turn). Multiply the step by  $w \bar{z}_{\text{now}}$  and take the imaginary part; the  $S$  term is real and drops:  $w \text{Im}(\bar{z}_{\text{now}} e^{i\theta_0} z_{\text{next}}) + w \text{Im}(\bar{z}_{\text{now}} e^{-i\theta_0} z_{\text{before}}) = \sum_a R_a (g_{i,i+a} + g_{i,i-a})$ . The first term is  $-W_i(t+1)$  and the second  $W_i(t)$ , and  $g_{i,i-a} = -g_{i-a,i}$ , so

$$w [W_i(t+1) - W_i(t)] + \sum_a [R_a g_{i,i+a} - R_a g_{i-a,i}] = 0 :$$

an exact lattice continuity equation at every Node and interval, the density  $\rho_i = w W_i$  and the flux through the Link  $(i, i+a)$  equal to  $R_a g_{i,i+a}$ , for any paces, any phases and any levels, the per-Node identity; the summed conservation needs the weight  $1/p_i^2$ , **DM** symmetric (S.9); the integer step's remainders add the term of S.9's domain. (b) Flux over density is the group velocity. For  $z = A e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}$ ,  $\rho = w A^2 \sin \omega$  and the flux is  $R_a A^2 \sin k_a$  (common sign), so flux over density is  $(R_a/w) \sin k_a / \sin \omega = d\omega/dk_a$  from the

band  $2w \cos \omega = S + 2 \sum_a R_a \cos k_a$ : the current is  $\rho v$  exactly, and the law's Node current  $J_a = g_{i,i+a} + g_{i-a,i}$  obeys  $J_a = 2(w/R_a)Wv$  on a plane record,  $(6 \text{ den}/\text{num})Wv$  at the vacuum's paces where  $R_a/w = \text{num}/(3 \text{ den})$  (the case (d) names), nature's  $j = \rho v$  with the band's constant and no  $\sin \omega_b$ . (c) The covector. Inserting the plane wave in the step with the phases (the main text's convention, Section 3.5) gives  $\omega = \omega_{\text{band}}(\mathbf{k} + \boldsymbol{\theta}) + \theta_0$  for the sense + (the sign convention of the law's document; S.41 (b)): the pair  $(\theta_0, \theta_a)$  shifts  $(\omega, k_a)$  and transforms as  $(\omega, \mathbf{k})$  does; at long wavelength and small gap every band is the Lorentz form  $\omega^2 = \omega_0^2 N^2 + c_s^2 k^2$  with  $c_s^2 = \text{num}_s/(3 \text{ den}_s)$  (from  $2 \cos \omega = S/w + (2 \text{ num}/3 \text{ den}) \sum_a \cos k_a$ ), so  $(\theta_0, \theta_a)$  is a covector under the band's boost: minimal coupling,  $A_0$  on the time step and  $A_a$  on the Link. (d) The halved current over the time level's wall. The sign holder under the rotation lays the time line and three odd axis lines sourced by the sign's current; the derivation's own is one kernel over one wall and the halving: Maxwell's pair with one kernel sources  $(\theta_0, \theta_a)$  from  $(\rho, \text{flux}/c^2)$  over one wall. The Node's flux on the axis  $a$  is the mean of its two Links',  $R_a(g_{i,i+a} + g_{i-a,i})/2 = R_a J_a/2$ , and  $R_a/w = \text{num}/(3 \text{ den})$  at the vacuum's pace, so flux over  $c_s^2 \rho$  is  $[(\text{num}/3 \text{ den}) J_a/2]/[(\text{num}/3 \text{ den}) W] = (J_a/2)/W$ : the odd line on the axis  $a$  is sourced by the halved current  $J_a/2$  over the same wall as the time level's  $W$ , with no number, when the pair is read at the source family's own  $c_s$ ; read at light's  $c^2 = 1/3$  the factor  $c_s^2/c^2 = \text{num}_s/\text{den}_s$  appears, 1 for light,  $1 - 10^{-28}$  at nature's matter,  $2/3$  at [2, 3] in this small-gap form ( $c_s^2 = \text{num}_s/(3 \text{ den}_s)$ ), where the band's own  $c_m^2 = \omega_0/(3 \tan \omega_0)$  of S.21 and S.22 gives 0.752; the two agree as the gap closes. The halving is not a choice: the Node's current is the mean of its two Links as the continuity identity defines it, as the pace reads the tension on an axis as the mean of its two ends' tensions. (e) Coulomb over  $\gamma$ . For a source moving at  $v$  along the Link and a reader co-moving with it, the time level is the moving density's,  $\gamma$  times the rest level by the current's covector ((a), (c)) at fixed total  $W$ , within (c)'s domain, so the electric force across the beam is  $\gamma F_0$ ; the odd level is sourced by  $J_a/2 = (3 \text{ den}/\text{num})Wv = Wv/c_s^2$ , the flux over  $c_s^2$  of (d), the reader at  $v$  feels  $-v \partial \theta_a / \partial y$ , a magnetic force  $\beta^2 \gamma F_0$  toward the source; the total is  $\gamma(1 - \beta^2)F_0 = F_0/\gamma$ , with  $\gamma$  Lorentz's at nature's gap, where the band is Lorentz's to  $9 \times 10^{-20}$  at the LHAASO bound (S.21); at the computed pair [2, 3] the band's own factor differs from it, and  $\beta = 0.5$  lies beyond the axis's largest group speed, so the two numbers below are the form's at nature's gap. Check:  $1/\gamma = 0.99499$  at  $\beta = 0.1$  and  $0.86603$  at  $\beta = 0.5$ , the law's row (b2)'s prediction; the identity (a) holds in exact arithmetic on a chain with contents and phases at every Node (the same check as S.9's, the phases a unitary on each read). Domain: the covector is the band's at long wavelength and breaks at the lattice's anisotropy  $k^2/12$  ( $2 \times 10^{-4}$  at  $k = 0.05$ ,  $1.0 \times 10^{-28}$  at nature's Compton wavelength); the pairing holds at the source family's own  $c_s$ , so a source and a reader of one family see one light speed. Status: theorem for (a) and (b), the conservation; derived at long wavelength for (c) to (e), the force's form, Peierls' coupling; lattice for the identity, clicks for the force.

**Derivation S.34** (the composed paces, the acts' factors and the exponential metric, Section 2.3) Inputs: the model's one premise beside the line, that a level is a count of units and every act is measured in the Node's own units (Section 2.3); the clock  $p_0$  as Rule3's coefficient in  $S$ ; the band at a pace,  $S.1$ ; the one assumption  $p_a = p_0^2/\Gamma$ . Steps: (a) The composition. One unit of level is one act on the clock as it stands, so it slows the clock by one part in  $\Gamma$  of the clock as it stands:  $p_0(c+1) = p_0(c)(1 - 1/\Gamma)$ , whence  $p_0(c) = \Gamma(1 - 1/\Gamma)^c$ . The functional equation  $P(U+l) = P(U)P(l)$  is Cauchy's, whose continuous solutions are the exponentials and no other, so  $N = p_0/\Gamma = e^{-U}(1 + O(U/\Gamma))$  with  $U = c/\Gamma$ ; the quadratic form's subtraction  $p \rightarrow p - 2$  is as memoryless and is excluded only by the premise that the read is an act measured in the Node's units, like the write. Check: at  $\Gamma = 6,000$  and  $c = 300$ ,  $(1 - 1/\Gamma)^c = 0.951225$  against

$e^{-0.05} = 0.951229$ , a relative  $4 \times 10^{-6}$ , which is  $U/(2\Gamma)$ . (b) The ruler.  $p_i = p_0^2/\Gamma = \Gamma N^2$  (the Link pace  $p_a$  at the vacuum's factor  $q = \Gamma$ ) gives the axis ruler  $h = p_0/p_a = 1/N$ :  $hN = 1$  (the one assumption, fixed by the bending's measurement, the perihelion choosing the reading and testing the choice, Section 2.3). (c) The metric. S.1 at the paces reads, with  $\nu = \text{num}/\text{den}$ ,  $2(1 - \cos \omega) = 2(1 - \nu)N^2 + (\nu/3)N^4 \sum_a s(k_a)^2$  with  $s(x) = 2\sin(x/2)$ ; at long wavelength and a small gap, where  $c_m^2 \rightarrow \nu/3$ ,  $\omega^2 = N^2 [\omega_0^2 + c_m^2 (k/h)^2]$  with  $k/h = kN$ : Klein-Gordon's on a static metric with the lapse  $N$  and the spatial scale  $h$ ,  $ds^2 = -N^2 dt^2 + h^2 d\mathbf{x}^2 = -e^{-2U} dt^2 + e^{2U} d\mathbf{x}^2$ , the exponential metric in isotropic coordinates [15], whose post-Newtonian expansion is  $g_{00} = -(1 - 2U + 2U^2)$  and  $g_{ij} = (1 + 2U + 2U^2)\delta_{ij}$ :  $\gamma = 1$  and, for this metric,  $\beta = 1$ , the spatial second order 2 against Schwarzschild's 1.5 (S.19 for  $\gamma = \beta = 1$ ; the second order this derivation's). (d) The acts' factors. Rule3 measured in the Node's units: a count per Node is a count per proper volume  $h^3$ , a step per interval is a step per proper interval  $N$ , and the write is Rule3's second difference in time, so a count source carries  $N^2/h^3 = N^5 = p_x p_y p_z / (p_0 \Gamma^2)$  (check:  $\Gamma^3 N^6 / (\Gamma N \cdot \Gamma^2) = N^5$ ); the Wronskian, one time difference, carries  $N/h^3 = N^4 = p_x p_y p_z / (p_0^2 \Gamma)$  and its readers' turn the remaining  $N$ ,  $\theta = (\text{level}/\Gamma)(p_0/\Gamma)$ , the turn per proper interval; the Link angle is plain by the law's line, the Port's phase the odd lines' level times  $4\omega_0$  read with no factor (S.56 (a)). With a tension the three rulers differ,  $p_0/p_a$  along the axis  $a$ , and the volume is their product. (e) The truncation. The clock's quadratic form  $(\Gamma - c)^2 + c^2 = \Gamma^2(1 - 2U + 2U^2)$  is  $\Gamma^2 e^{-2U} = \Gamma^2(1 - 2U + 2U^2 - 4U^3/3 + \dots)$  to the second order, differing at  $-4U^3/3$ ; the Link's  $(\Gamma - 2c)^2 = \Gamma^2(1 - 4U + 4U^2)$  is  $\Gamma^2 e^{-4U} = \Gamma^2(1 - 4U + 8U^2 - \dots)$  to the first order only. Check: at  $U = 0.05$  the clock's form gives 0.905000 against  $e^{-0.1} = 0.904837$ , the difference  $1.63 \times 10^{-4}$  against  $4U^3/3 = 1.67 \times 10^{-4}$ ; the Link's gives 0.8100 against  $e^{-0.2} = 0.8187$ . The horizon of the quadratic forms at  $c = \Gamma/2$  is the truncation's zero of  $\Gamma - 2c$  and not the law's (S.3). Status: derived under the one assumption; lattice.

**Derivation S.35** (the preferred-frame parameter of a diagonal metric, Section 8.1) Inputs: the model's metric is diagonal in the lattice's frame by construction: the content enters the clock  $p_0$  and the Link pace  $p_a$ , both real and even in time and in space, the content's rest is Poisson's in the lattice's frame with no speed in its kernel, and no odd real Link read exists, since a real asymmetric hopping with a curl breaks the reciprocity the conserved form needs (Theorem 3); so  $g_{0a} \equiv 0$  in the lattice's frame for every motion of the source (Section 8.1; a write with no odd Link part; the law's write (Section 2.3, S.56) carries the odd part and gives 0). The parametrised post-Newtonian metric in the theory's preferred frame (Will's review [3], in the standard gauge):  $g_{0j} = -\frac{1}{2}(4\gamma + 3 + \alpha_1 - \alpha_2 + \zeta_1 - 2\xi) V_j - \frac{1}{2}(1 + \alpha_2 - \zeta_1 + 2\xi) W_j$ , with  $V_j$  and  $W_j$  the two vector potentials of the matter current, related by  $V_j - W_j = \chi_{,0j}$  for the superpotential  $\chi$ . Steps: (a) The post-Newtonian gauge freedom  $x^0 \rightarrow x^0 + \lambda \chi_{,0}$  adds  $\lambda \chi_{,0j} = \lambda(V_j - W_j)$  to  $g_{0j}$ , moving weight between  $V_j$  and  $W_j$  and leaving the sum of their two coefficients,  $(4\gamma + 3 + \alpha_1 - \alpha_2 + \zeta_1 - 2\xi) + (1 + \alpha_2 - \zeta_1 + 2\xi) = 4\gamma + 4 + \alpha_1$ , gauge-invariant. (b) A metric whose  $g_{0j}$  vanishes in its preferred frame for every source current has both coefficients 0 in some gauge, so the invariant sum vanishes:  $4\gamma + 4 + \alpha_1 = 0$ ,  $\alpha_1 = -4(1 + \gamma)$ , which is  $-8$  at the model's  $\gamma = 1$  (S.19 in U). (c) The frame-dragging precession of a gyroscope near a rotating body is  $(4\gamma + 4 + \alpha_1)/8$  of general relativity's Lense-Thirring value, 1 in general relativity and 0 in the model: Gravity Probe B's 0 against  $37.2 \pm 7.2$  milliarcseconds per year [16] is the same fact as  $\alpha_1 = -8$ , read by two experiments of nature. (d) Had the model's read been covariant, the boost of its static metric  $N^2 = e^{-2U}$ ,  $h^2 = e^{2U}$  to the velocity  $v$  would give  $g_{0i} = \gamma_L^2 v_i (N^2 - h^2) = -4U v_i$  to first order ( $\gamma_L$  Lorentz's, not the post-Newtonian  $\gamma$  of (b)), Einstein's, with no new coefficient; the diagonal read in the lattice's frame is what removes it, so the finding is the frame's and not a number's (the carrier's derivation, S.36 of the long

version). Check:  $-4.000$  for the boosted cross term at first order; nature's  $\alpha_1$  from the binary pulsar PSR J1738+0333 is  $|\alpha_1| < 4 \times 10^{-5}$  [17] and from lunar laser ranging  $10^{-4}$  [18], so the finding stands at  $2 \times 10^5$  and  $10^5$ ; Gravity Probe B's drag at 5 standard deviations; the two are one finding, not two. Domain:  $\alpha_1$  alone is gauge-invariant here; with  $g_{0j} = 0$  in the lattice frame,  $g_{00} = -e^{-2U}$  and  $g_{ij}$  isotropic, the standard post-Newtonian gauge, the two  $g_{0j}$  coefficients vanish separately and give  $\alpha_1 = -8$  and  $\alpha_2 = -1$  together ( $\alpha_2 = -1$  with  $\zeta_1 = 2\xi$ ,  $\zeta_1 = 0$  under the post-Newtonian conservation laws and  $\xi = 0$ , the preferred-location parameter, taken;  $\alpha_1 = -8$  needs no such condition), the sum  $4\gamma + 4 + \alpha_1 = 0$  the gauge-invariant statement, against  $|\alpha_2| < 4 \times 10^{-7}$  (the Sun's spin [19]) and  $< 1.6 \times 10^{-9}$  (solitary pulsars [20]);  $\alpha_1 = -8$  holds for every motion of the source,  $g_{0j} = 0$  in the lattice's frame being Rule3's for any source (no current is read, no vector potential is sourced);  $\alpha_2 = -1$  holds for a static source and is not derived for a moving one, every value of order one being excluded by  $|\alpha_2| < 4 \times 10^{-7}$  in any case; the condition below is  $\alpha_2$ 's alone, the velocity-dependent terms of  $g_{00}$  depending on how a moving record's count writes, which the law does not fix: named, not claimed. Status: derived; the frame's parameters of a write with no odd Link part ( $g_{0j} \equiv 0$ ):  $\alpha_1 = -8$ ,  $\alpha_2 = -1$ ; the law's write (Section 2.3, S.56) carries the odd part and gives 0; the pair stands in 8.1 as the misses of the write without it; clicks.

**Derivation S.36** (the mass defect at a kept count and the binding bound, Section 8.1) Inputs: Theorem 3, the weighted form conserved by Rule3 at fixed paces for any records; the invariant of S.14, a body of  $C$  units of the invariant at the rotation  $\omega_b$  having the share  $C \sin \omega_b$  (the energy per quantum  $\sin \omega_b$  in units of  $T$ , Planck's line); the rest rotation in a content,  $\omega_0(U) = N\omega_0$  to the order  $\omega_0^2$  (S.15 with  $f \rightarrow 1$ ); the massless kernel of a holder at the weight  $W$ : the level at  $r$  Links from a count  $C$  is  $3CW G_0(r)/E$  with  $G_0$  the lattice Green's function per unit source,  $G_0(0) = 0.2527$  (Watson) and  $1/(4\pi r)$  far,  $e^{-r/R}$  beyond a gapped holder's range (S.11). Steps: (a) No energy from combinatorics. Two records that approach, overlap, exchange quanta through their tails, join into one standing record or part again are records stepped by Rule3; if the paces do not change, the weighted total form before and after is the same number in the rational line (Theorem 3), the plain total where the paces are uniform, within the remainder term of the share's change, Section 3.4 (S.5), with the rounding, and a click moves share from one record to another and conserves the total in the credit's unit, the weighted total where the two records stand at one pace. So no joining, parting or crediting of clusters lowers the share at a kept count: there is no energy released, and a nucleus bound that way would have no binding energy. (b) What the defect is. The one act that changes a body's share at a kept count is a change of its paces, a content read; a body in its own content  $x_{\text{self}}$  rotates at  $N\omega_0$  with  $N = e^{-x_{\text{self}}}$ , so its click mass is  $C \sin(N\omega_0) = C \sin \omega_0 - C\omega_0 x_{\text{self}} \cos \omega_0 + O(x^2)$ : the mass defect is at most the self-read's redshift  $C\omega_0 x_{\text{self}}$  to the order  $\omega_0^2$ , the bound mode's depression  $\omega_0 - \omega_b$ , the potential's redshift less the bound mode's kinetic rise (half the redshift for a  $1/r$  well by the virial, S.45), which is what S.45 (b) of the long version computes at the fixed point; this is the law's "a body that binds is lighter in every holder it sources by its binding". Check: a well deepened from 300 to 600 levels at  $\Gamma = 6,000$  lowers the share read at the vacuum's coefficients (Section 3.4) by 3.8 percent to first order ( $x\omega_0/\tan \omega_0$  at  $x = 0.05$ ; 3.86 by  $\sin(N\omega_0)$  from  $e^{-0.05}$  to  $e^{-0.10}$ ); the share at the Node's own paces carries the factor  $p_i^2(t_1)/p_i^2(t_2)$  of the drift ratio (S.14) and rises. (c) The bound. A body of count  $C$  over  $r$  Links, bound by a holder at the weight  $W$  and the divisor  $E$ , writes on itself at most the one-Node value, the lattice kernel at the source,  $x_{\text{self}} \leq 3G_0(0)CW/(\Gamma E) = 0.758CW/(\Gamma E)$ , independent of  $r$ ; for a uniform ball of radius  $r$  the centre's self-level is 1.5 times the whole count's level at  $r$  ( $0.36CW/(\Gamma E r)$  at the centre) and the count's share-weighted average 1.2 times it, the first-order shift of a mode under a

non-uniform content, a profile-dependent scaling and not a bound, so

$$\frac{\text{defect}}{\text{rest}} \leq 0.76 \frac{CW}{\Gamma E}, \quad \frac{\text{defect}}{\text{rest}} \times r \approx 0.29 \frac{CW}{\Gamma E} \text{ for a uniform ball, } 0.36 \text{ at its centre.}$$

(d) The numbers. The deuteron's defect is  $2.224 \text{ MeV} / 1875.6 \text{ MeV} = 1.2 \times 10^{-3}$  and the nucleon's in iron  $8.8/939 = 9 \times 10^{-3}$  [21], so, with the ball's estimate, they need  $CW/(\Gamma E r) \geq 4 \times 10^{-3}$  and  $3 \times 10^{-2}$ ; at  $r$  about  $6 \times 10^{10}$  Links for a nucleon (0.84 fm at the MAGIC Link,  $1.3 \times 10^{-26}$  m) this is  $W$  of order  $10^{19}$  to  $10^{22}$  at  $\Gamma = 10^{14}$  (the law's (A3) illustration; the bound's form is the claim and the number is not) and  $E = 1$ , the lower end for a cloud of  $C = 1,836$  records in the electron's unit and the upper for  $C = 1$ , at the deuteron's defect (iron's raises each by 7.9), taken for illustration, growing with  $\Gamma$ , which no click bounds (S.26), under the reading of the nucleon as a cloud, which S.45 of the long version replaces (the nucleon one quantum of its own family, its binding the Yukawa well with its declared  $W$ ); the bound's form does not move with the reading of  $\Gamma$ , its numbers do. The credit's quantum does not supply it: the deuteron's 2.22 MeV is 4.35 quanta of 0.511 MeV and iron's nucleon's 8.8 MeV 17.2, not wholes, and no act credits share to nowhere. (e) The one lattice lightening without content, the band's  $\sin \omega_b$  falling to 0 at the band's top, acts at rotations of order one and makes a body nearly massless, not 99 percent of its quanta's mass: not a nuclear binding. Check of the kernel: Watson's integral gives  $G_0(0) = 0.252731$ ,  $3G_0(0) = 0.7582$ ;  $G_0(1) = G_0(0) - 1/6 = 0.08606$ ,  $3G_0(1) = 0.2582$ , 1.08 of the far kernel  $3/(4\pi) = 0.2387$  at  $r = 1$ . Status: derived, a bound conditional on the declared  $W$ , which no click bounds; the size law kept; lattice for the well, clicks for the defect.

**Derivation S.37** (the gapped holder's kernel, Section 4.1) Inputs: a holder's static line in the vacuum, from the start's condition (Section 4.1):  $6 \text{ den } L - \text{num } S_6(L) = 3 \text{ den } \sigma$ , that is  $\text{num } \Delta L - 6(\text{den} - \text{num}) L = -3 \text{ den } \sigma$  with  $\Delta = S_6 - 6$  the lattice Laplacian and  $\sigma$  the source per interval; the reach  $\kappa^2 = 6(\text{den} - \text{num})/\text{num}$  of S.11, for  $0 < \text{num} \leq \text{den}$  as S.11 has it (the division by num below and the positive kernel need it; at num = 0 the static line has local support, at a negative numerator the alternating tail of S.11); the write and the read of every rotation holder the same act,  $W$  over  $E_s T$  into the level and the level over  $\Gamma$  into the angle (Section 3.5), whatever the holder's pair. Steps: (a) Dividing by num:  $\Delta L - \kappa^2 L = -(3/\nu) \sigma$ , whose lattice Green's function is  $G_\kappa(\mathbf{r}) = \int \frac{d^3 k}{(2\pi)^3} \frac{e^{i\mathbf{k} \cdot \mathbf{r}}}{6 - 2 \sum_a \cos k_a + \kappa^2}$ , far away  $e^{-\kappa r} / (4\pi r)$  per unit source: the level is  $3 e^{-r/R} / (4\pi \nu r)$  per unit  $\sigma$  with  $R = 1/\kappa$ , Yukawa's kernel with the massless one's constant, the range the gap's and the strength at the Link the massless one's:  $G_\kappa(1) = G_0(1) (1 - 0.92 \kappa + O(\kappa^2))$ , so the strength at one Link is the massless one to one part in  $R$  (and the factor  $1/\nu = 1 + O(\text{gap})$ ). Check: the exact infinite-volume integrals,  $G_0(1) = 0.08606$  (Watson) and  $G_{1/20}(1) = 0.08216$  by the exact lattice Yukawa integral, the ratio 0.9546 at  $R = 20$  Links (0.08209 and 0.954 at the first order in  $\kappa$ ,  $G_0(1) - \kappa/4\pi$ ), the slope  $1 - 0.9246 \kappa$ ;  $G_{1/400}(1)/G_0(1) = 0.9977$ ; the  $400^3$  grid's own pair at  $R = 20$ , 0.0857 and 0.0822, has the ratio 0.959 against the exact 0.9546. (b) The coupling. The sign holder's force constant,  $\alpha_{\text{law}}$  (S.26), is the write's half, the kernel's  $3/(4\pi)$  and light's  $\sqrt{3}$ , none of them the holder's own; a gapped holder written and read by the same acts has the same three factors and the kernel above, so it couples at  $\alpha_{\text{law}} (1 - 0.92/R)$  at one Link,  $1.08 \alpha_{\text{law}}$  on the lattice at  $r = 1$ , and  $\alpha_{\text{law}} e^{-r/R}/r$  beyond: every rotation holder Rule3 steps, massless or gapped, abelian or three-line, couples at the charge's  $\alpha$  at the Link. A coupling  $k/(\Gamma E_s)$  itself, the universe's  $1/28.33$ , would need the factor  $8\pi/(3\sqrt{3}) = 4.84$  on the kernel, which no gap supplies; a contact strength  $3/\kappa^2$  appears only for  $\kappa > 1$ , a range below one Link, where the kernel's lattice sum departs from its continuum integral. (c) The numbers. The  $W$  boson's range  $\hbar/(m_W c) = 2.45 \times 10^{-3}$  fm and the pion's 1.41 fm are  $1.9 \times 10^8$  and  $1.1 \times 10^{11}$  Links

at the Link's bound  $1.3 \times 10^{-26} \text{ m}$  (S.24), so  $1/R$  is  $5 \times 10^{-9}$  and  $10^{-11}$ : the strength at the Link is the massless one to those parts. A pion-gap holder at its range with the law's coupling has the well's depth  $\alpha \hbar c e^{-1}/R = 1.44 \text{ MeV fm} \times 0.368/1.41 \text{ fm} = 0.38 \text{ MeV}$  at  $r = R$  and binds no state: a screened Coulomb well binds only for  $\mu g R / (\hbar c)^2 \geq 0.840$ , and at  $g = \alpha \hbar c$ ,  $R = 1.414 \text{ fm}$ ,  $\mu = 469.5 \text{ MeV}$  (nature's deuteron binds two nucleons about their common centre, so this is nature's own reduced mass, the comparison's convention for nature's column; the law's body is one quantum about the pinned centre at  $\mu = m^*$ , postulate 2, the two columns differing by the factor 2 in  $\mu$  with  $W \propto 1/\mu$  at fixed  $g^2$ , S.45 (a) of the long version) the number is 0.0245, 34 times under the threshold, so the deuteron's 2.22 MeV stands against no binding at all; a holder with the  $W$  boson's gap with the law's  $\alpha$  and nature's  $m_W$  gives the Fermi constant  $G_F = \pi \alpha / (\sqrt{2} m_W^2) = 2.5 \times 10^{-6} \text{ GeV}^{-2}$  against nature's  $1.166 \times 10^{-5}$  [22], short by 4.6, or  $m_W = 37 \text{ GeV}$  from  $G_F$  against 80.4 [22]. Check of Yukawa's identity: the range  $R = \sqrt{\text{num}/(6(\text{den} - \text{num}))}$  equals the Compton length  $c/\omega_g$  of the holder's own travelling quanta to the gap's first order (S.11), with no input. Status: derived, one derivation, the comparisons with nature's numbers supporting findings; lattice for the kernel, clicks for the comparisons, supporting findings.

**Derivation S.38** (the Greenberger-Horne-Zeilinger (GHZ) correlation under the declared credit, Section 7.2) Inputs: the GHZ family, light's pair with four real lines laid as one event and never summed at a Node, every part laid alike and stepped by the same line, so the four are identical at every NodeDetector (the determinism of equal parts, Section 7.2); three NodeDetectors  $A, B, C$ , each with its setting  $(p, q) = (\cos a, \sin a)$  up to the norm and its pattern, one integer pair per part:  $A(p, p, q, q)$ ,  $B(p, q, p, q)$ ,  $C(p, -q, -q, -p)$ , the  $+$  port reading the part  $k$  as  $e_k(+) = \alpha_k p + \beta_k q$ , with  $\alpha_k, \beta_k$  here the pattern's selectors  $(1, 0), (0, 1), (0, -1), (-1, 0)$  multiplying  $(p, q)$ , not S.47's laid amplitudes, and the  $-$  port as  $e_k(-) = \alpha_k(-q) + \beta_k p$ ; the credit, the pair's own over three sides:  $J(\pm, \pm, \pm) = (\sum_{k=1}^4 e_k(A) e_k(B) e_k(C))^2$  on equal parts, the common factor of the parts' amplitudes leaving every ratio. Steps: (a) With  $c_a = \cos a$ ,  $s_a = \sin a$  and the same for  $b$  and  $c$ , the  $+$  ports read  $A$  as  $(c_a, c_a, s_a, s_a)$ ,  $B$  as  $(c_b, s_b, c_b, s_b)$  and  $C$  as  $(c_c, -s_c, -s_c, -c_c)$ , so  $\sum_k e_k(A) e_k(B) e_k(C) = c_a c_b c_c - c_a s_b s_c - s_a c_b s_c - s_a s_b c_c = \cos(a+b+c)$ . A  $-$  port replaces  $(p, q)$  by  $(-q, p)$ :  $A$ 's  $-$  port reads  $(-s_a, -s_a, c_a, c_a)$ , and the sum is  $-s_a c_b c_c + s_a s_b s_c - c_a c_b s_c - c_a s_b c_c = -\sin(a+b+c)$ ; by the symmetry of the three patterns every port triple with an even number of  $-$  ports gives  $\pm \cos(a+b+c)$  and with an odd number  $\pm \sin(a+b+c)$  (for instance  $(-, -, +)$  gives  $-\cos$ ,  $(-, -, -)$  gives  $\sin$ ). (b) The eight joint shares are therefore  $\cos^2(a+b+c)$  at the four even triples and  $\sin^2(a+b+c)$  at the four odd, summing to 4 at every setting; the normalised shares are  $\cos^2/4$  and  $\sin^2/4$ . (c)  $E_3$ , the shares signed by the product of the three signs, is  $4 \cdot \cos^2/4 - 4 \cdot \sin^2/4 = \cos 2(a+b+c)$ . A pairwise  $E_{AB}$  sums over  $C$ 's two ports first, which gives  $(\cos^2 + \sin^2)/4 = 1/4$  for each of the four sign pairs of  $A$  and  $B$ , so  $E_{AB} = (1 - 1 - 1 + 1)/4 = 0$ , and the same for the other two pairs; each single marginal is  $1/2$ : no signalling and no correlation in any two, the correlation standing in the three alone. (d) Mermin's combination  $M = E(a, b', c') + E(a', b, c') + E(a', b', c) - E(a, b, c)$  at  $a = b = c = 0$  and  $a' = b' = c' = \pi/4$  is  $\cos \pi + \cos \pi + \cos \pi - \cos 0 = -4$ ,  $|M| = 4$ , where every local assignment of signs to the three ports, and every mixture of them, gives  $|M| \leq 2$  [23]. (e) With integer settings  $(p, q)$  the norms  $p^2 + q^2$  multiply every share alike and cancel, so  $x = (1, 0)$  and  $y = (1, 1)$  give the eight shares exactly, 0 and  $1/4$  on the three settings with one  $x$  ( $E_3 = -1$ ) and  $1/4$  and 0 on  $(x, x, x)$  ( $E_3 = 1$ ),  $M = -4$  exactly. Check: the trigonometric identity  $\cos(a+b+c) = c_a c_b c_c - c_a s_b s_c - s_a c_b s_c - s_a s_b c_c$  and  $\sin(a+b+c) = s_a c_b c_c + c_a s_b c_c + c_a c_b s_c - s_a s_b s_c$ . Nature's angles: the law's  $E_3 = \cos 2(a+b+c)$  is nature's  $\cos(\theta_A + \theta_B + \theta_C)$  at  $\theta = 2a$ , nature's  $x$  and  $y$  at  $\theta = 0$  and  $\pi/2$  the law's bases

(1,0) and (1,1), as Bell's angles are (S.10). Status: derived under the declared credit, the credit the one assumption (Section 7.2); clicks.

**Derivation S.39** (the prediction, the check and the open question, Section 5.2) (1) A clock in two places. Inputs: a bound body's beat between two of its modes,  $\Omega = \omega_j - \omega_i$ , scales by the clock  $N = e^{-U}$  at the body's Node (S.31, exactly at long wavelength, to  $(1 - N^2)\omega^2/8$  at finite  $\omega$ ); a record laid on two paths is one record (Section 2.5.1), and the body's internal state is carried on each path by its own rotation. Steps: on two paths  $U_1$  and  $U_2$  for a hold time  $\tau_{\text{hold}}$  the internal beats, of frequency  $\Omega = \omega_j - \omega_i$ , accumulate the phases  $\Omega N_1 \tau_{\text{hold}}$  and  $\Omega N_2 \tau_{\text{hold}}$ , differing by  $\Omega(N_1 - N_2)\tau_{\text{hold}} = \Omega \Delta U \tau_{\text{hold}}$  to first order; for a body prepared in an equal superposition of the two modes the internal states on the two paths overlap as  $|\cos(\Omega \Delta U \tau_{\text{hold}}/2)|$ , which multiplies the path fringes' visibility. Check: strontium's clock line 429.228 THz,  $\Omega = 2\pi \times 4.29228 \times 10^{14} = 2.697 \times 10^{15} \text{ s}^{-1}$ ;  $\Delta U = gh/c^2 = 9.807 \times 1/(2.998 \times 10^8)^2 = 1.091 \times 10^{-16}$  for one metre ( $g$  Earth's acceleration,  $h$  the height);  $\tau_{\text{hold}} = 1 \text{ s}$ :  $\Omega \Delta U \tau_{\text{hold}} = 0.294 \text{ rad}$ , the visibility  $\cos(0.147) = 0.989$ ; the correction  $(1 - N^2)(\omega_1^2 + \omega_1\omega_2 + \omega_2^2)/24$ ,  $(1 - N^2)\omega^2/8$  at  $\omega_1 = \omega_2$ , is  $1.0 \times 10^{-37}$  at the axis bound's  $\omega = 2.0 \times 10^{-14}$ ,  $3.4 \times 10^{-29}$  at LHAASO's ( $U = 10^{-9}$ ; S.31). The semiclassical expectation [24] has the same magnitude; the law's statement is that the one record's beat reads the content at its Node, so the dephasing is the clock's and no new line. (2) The gravity of light. Inputs: the one write: a holder of the content gains the form  $D$  of every record over its wall (Section 2.2); Theorem 3 and S.5: a travelling light record's form summed over its Nodes is its share, its count  $n$  (energy  $nT$ ), and  $D = A^2 \sin^2 \omega$  per Node for a travelling wave as for a standing one; the tension sources the axis parts alone, which enter the Link's factor and not the clock (Section 2.3). Steps: a light packet of count  $n$  writes  $n$  quanta per interval into the massless holder as a static body of count  $n$  does; this, the integrated write, is the derived part. The premise taken next is the static far kernel,  $3n/(4\pi r)$  over  $E_g$ , which is the field of a source at rest: under it the clock's well is  $U = 3n/(4\pi \Gamma E_g r)$ , Eq. (11)'s  $G_{\text{clock}}$  times  $n$ , and a static body falls toward the packet at  $G_{\text{clock}} n/r^2$ , that is  $1 \times E/c^2$  for the packet's active mass. The field of the packet itself, a source moving at the holder's own characteristic speed, is not derived here and is open: the retarded field of a sub-characteristic source of the same holder is  $\phi \propto 1/\sqrt{s_{\parallel}^2 + (1 - \beta^2)s_{\perp}^2}$ , directional and  $\gamma_L$  times the static field across the motion, singular at the characteristic speed in the continuum, which the lattice's dispersion blunts and does not remove. In general relativity a pencil of light pulls a static body at  $2 \times E/c^2$  [25], the beam's longitudinal stress  $\rho c^2$  along the beam and 0 across entering  $\rho + \sum_i p_i/c^2 = 2\rho$ ; in the model the stress of the wave, the Link's booking  $2 \text{ num } b^2 \sin^2 k$  and the Node's tension  $\text{num } b^2 \sin^2 k$  (S.7), enters the Link's factor on its axis and the clock reads the form alone, so under the static kernel the factor is 1: a prediction for a declared universe, no measurement of nature, the moving packet's field open. (3) The black hole's quanta. Inputs: the composed paces (S.34):  $p_0 = \Gamma(1 - 1/\Gamma)^c$ ,  $p_i = p_0^2/\Gamma$  rounded to the unit, every Link coefficient  $R_a = 2 \text{ num } p_i^2 q^2/\Gamma^2$ ; Theorem 1, the step a bijection; a collapse ends in Nodes whose content exceeds the frozen content (Section 2.3). Steps:  $p_i$  rounds to 0 where  $\Gamma e^{-2U} < 1/2$ , that is  $U > \frac{1}{2} \ln(2\Gamma)$ , the frozen content  $c_f = (\Gamma/2) \ln(2\Gamma)$  ( $4.70 \Gamma$  levels, 28,206 at  $\Gamma = 6,000$  with  $p_0$  rounded to the unit before squaring, the law's rounding); there the Node's own six coefficients are 0, so it reads nothing of its neighbours, while its neighbours' coefficients, each the reading Node's own, still read it: the cut is one-way, the frozen Node dark by the share's declaration and not cut; what leaves a frozen region is the levels frozen in it, read by the surface's active shell at its pace, matter's a near-static source of a static well, light's a level drifting linearly that no line of the law closes; whether the law's black hole radiates, and what, is not established by the

line and is open,  $T_H = 0$  not claimed (a reading of the line excluded by two derivations). Hawking's temperature [26]/[27] for one solar mass, the number the open question is set against (a comparison only, as Bekenstein's entropy [28] beside it), is  $\hbar c^3/(8\pi G M k_B) = 6.17 \times 10^{-8}$  K. The region's entropy as a count of the state is the count of its kept levels, the declared bound's logarithm per frozen Node; what it means for radiation is open. Check:  $\hbar c^3/(8\pi G M_\odot k_B)$  with  $M_\odot = 1.988 \times 10^{30}$  kg gives  $6.17 \times 10^{-8}$  K;  $\frac{1}{2} \ln(12,000) = 4.696$ . Status: (1) derived; clicks. (2) derived, the integrated write, the coefficient under the static far kernel, the moving packet's field open; clicks, untested. (3) derived under the one assumption, what it means for radiation open; lattice.

**Derivation S.40** (what a body writes into the holder of the sign, Section 4.3) Inputs: the continuity identity of S.33: at every Node  $w[W_i(t+1) - W_i(t)]$  equals the net of the Link fluxes  $R_{ag}$ , so the weighted sum  $\sum_i W_i/p_i^2$  over a body is exactly conserved for a static self-level, the plain sum its uniform-pace case (S.33 (a), S.9), and a single Node's  $W_i$  changes only by a current through its Links (the credit books the conserved form's own current through a front, the Link's factor squared its one weight, the Node's  $1/p_i^2$  cancelling against the read's  $p_i^2$ , so the count is the conserved form's flux, exact at paces fixed in time; a gapped quantum's share in a well differs from  $W_{\text{rec}}$  by the rest term's shift  $4U(\text{den} - \text{num})/\text{den}$ , the law's physics read as the count; where the paces move the share changes by the work term, the body's; a ratio of two readings at one pace is unchanged); in particular at one isolated Node,  $z_{t+1} + z_{t-1} = 2 \cos \omega_t z_t$  with any real  $\omega_t$  keeps  $W = \text{Im}(\bar{z}_t z_{t-1})$  exactly ( $W_{t+1} = \text{Im}(2 \cos \omega_t |z_t|^2 - \bar{z}_{t-1} z_t) = W_t$ ). The holder's line at the vacuum's pace,  $\ell_{t+1} - 2\ell_t + \ell_{t-1} = \frac{1}{3}\Delta\ell + s_t$ , with  $s$  the write per interval (the Wronskian over  $E_s T$ ): the wave equation at the speed  $1/\sqrt{3}$  with the retarded solution  $\ell(\mathbf{r}, t) = 3s(t - \sqrt{3}r)/(4\pi r)$  for a point source; a static record has  $D = 0$  and  $F = 0$  and carries no quanta (Section 2.4); the share of a travelling wave of the holder's record is  $A^2 \sin^2 \Omega/T$  quanta per Node (S.5 at  $[1, 1]$ ) moving at  $1/\sqrt{3}$ . Steps: (a) A steady rotation. A body at rest with the standing plane record  $z_i = \phi_i e^{-i\omega_b t}$ ,  $\phi$  real up to one phase, has  $W_i = \phi_i^2 \sin \omega_b$  constant at every Node and the Link quantity  $g_{ij} = \text{Im}(\bar{z}_i z_j) = 0$ : the source is static, the holder's level is its rest, no travelling event is born, no click anywhere. (b) Uniform motion. A uniformly moving source writes no travelling light in the long-wavelength band,  $\omega = c|k|$ , where no sub-c source is phase-matched: its retarded level is the uniformly moving rest, which carries no energy flux to infinity, as for a uniformly moving charge in Maxwell's theory. On the lattice light's phase speed falls to 0.392 Link per interval at the zone's edge along an axis, so a source faster than that is matched to zone-edge light (the forced recurrence's denominator  $2 \cos(k \cdot v) - 2 \cos \omega(k)$  vanishing there), which a body smooth over many Links sources only by its Fourier weight at that wave number, a small number set by the body's tail ( $e^{-\pi^2 R^2/2}$  for a Gaussian of rms  $R$  Links,  $\tanh^2(1/(2R))$ , of order  $1/(4R^2)$ , for an exponential tail), the exclusion's work carried by the group-speed condition that follows; at the computed pair no body reaches 0.392 (matter's largest group speed 0.385 along the diagonal); from the pair [9, 10] upward a body moving along an axis can reach it, so the exclusion is an approximation with a named domain and not an exact one (a finding of two derivations). (c) A breathing body. Breathing is a change of the amplitudes  $A_i(t)$  across the body by currents between its Nodes; the weighted total  $\sum_i W_i/p_i^2$  is kept, so where the paces are uniform over the body the monopole of the source is constant and radiates nothing, exactly; where they vary across the body the source written with the plain weight departs from the conserved total at first order in the pace's variation,  $2 \Delta p/p$  of the total, below  $10^{-9}$  for an atom in a laboratory well (the write's weight is the law's line, open). The far level of a spherically symmetric source distribution  $\sigma_i(t)$  of constant total  $S$  is, by the retarded integral with the retardation expanded, the known multipole expansion of the holder's wave equation's solution, standard mathematics

of the law's line  $(\sigma(t - \sqrt{3}|\mathbf{r} - \mathbf{r}'|) = \sigma(t_r) + \sqrt{3}(\hat{\mathbf{r}} \cdot \mathbf{r}')\dot{\sigma} + \frac{3}{2}(\hat{\mathbf{r}} \cdot \mathbf{r}')^2\ddot{\sigma} + \dots$ , the first order averaging to 0 over the sphere and  $\langle(\hat{\mathbf{r}} \cdot \mathbf{r}')^2\rangle = r'^2/3$ ,

$$\ell(r, t) = \frac{3S}{4\pi r} + \frac{3}{4\pi r} \cdot \frac{1}{2} \ddot{Q}(t - \sqrt{3}r), \quad Q(t) = \sum_i \sigma_i(t) r_i^2,$$

so a body of mean square radius  $\langle r^2 \rangle$  breathing with its second moment  $Q(t) = Q_0(1 + 2\delta \cos \Omega t)$ ,  $\delta$  defined by it with the monopole  $S$  constant and  $Q_0 = S\langle r^2 \rangle$ , writes the travelling level  $\ell_1 = 3s_{\text{eff}}/(4\pi r)$  at the frequency  $\Omega$  with  $s_{\text{eff}} = \delta \Omega^2 S\langle r^2 \rangle$ : the light born carries the breathing's frequency  $\Omega$  and not the rotation  $\omega_b$ , as Section 4.3 says, with the amplitude  $S \cdot 2\delta$  ( $\delta$  the breathing's depth) multiplied by  $\Omega^2 \langle r^2 \rangle / 2$ , the second moment's factor. The quanta radiated per interval through a far sphere are the share density times the speed times the area,  $4\pi r^2 \cdot \ell_1^2 \sin^2 \Omega / (\sqrt{3}T) = (3\sqrt{3}/4\pi) s_{\text{eff}}^2 \sin^2 \Omega / T$ , of order  $\delta^2 \Omega^6 S^2 \langle r^2 \rangle^2 / T$ : the scalar analogue of the quadrupole formula, and a far NodeDetector's click rate is that current's share through its front. Section 4.3's three conclusions stand: no light from a steady rotation, none from a uniform motion in the long-wavelength band, light at the beat  $\Omega$  from a breathing body. Check:  $9/(4\pi\sqrt{3}) = 3\sqrt{3}/(4\pi) = 0.4135$ ; the static kernel  $3/(4\pi r)$  is  $S.25$ 's; the retarded expansion is the standard multipole one at the speed  $1/\sqrt{3}$ . Status: derived, (b) by two derivations as stated; lattice for the Wronskian and the source, clicks for the far NodeDetector's rate.

**Derivation S.41** (the two-mode line under the rotation, Section 4.3) The convention: *this derivation's time turn*  $\mathcal{U}_t = e^{-i\theta_t}$  *is the sign convention of the law's document, the main text's and S.33's* (S.33's step multiplied by  $e^{-i\theta_t}$  gives this form): the sense + rotates at  $\omega_b + \theta$ . Inputs: the plane record's step under the holder of the sign (Section 3.5), the time turn  $\mathcal{U}_t$  applied to the plain step,  $z_{t+1} = \mathcal{U}_t(w^{-1}[Sz_t + \sum_a R_a(e^{i\theta_a} z_{t+a} + e^{-i\theta_a} z_{t-a})] - \mathcal{U}_{t-1}z_{t-1})$  with  $\mathcal{U}_t = e^{-i\theta_t}$  the time turn between  $t$  and  $t+1$  and  $\theta_a$  the Link phases, both the holder's level over  $\Gamma$  at the interval's start; light as the holder's own record, a level  $b_L \cos \omega_L t$  at the body with the gradient  $\nabla b_L$  across it; the body's bound modes  $\phi_m$ , real eigenvectors of  $\mathbf{M}$ , symmetric at the paces uniform over the body taken here (one  $w$ ), with  $(\mathbf{M}\phi_m)_i = 2w \cos \omega_m \phi_{m,i}$ , orthonormal; at nonuniform paces  $\mathbf{DM}$  is the symmetric one (S.45), the modes are orthonormal in the weights  $d_i = 1/p_i^2$  and every Link term below carries  $d_i R_{ij}$  in place of  $R_a$ . Steps: (a) The uniform turn is a gauge. Put  $z_t = e^{-i\Phi_t} u_t$  with  $\Phi_{t+1} = \Phi_t + \theta_t$ . Then  $\mathcal{U}_t z_t = e^{-i\Phi_{t+1}} u_t$ ,  $\mathcal{U}_t \mathcal{U}_{t-1} z_{t-1} = e^{-i\Phi_{t+1}} u_{t-1}$  and  $z_{t+1} = e^{-i\Phi_{t+1}} u_{t+1}$ , so  $u$  obeys the plain line  $u_{t+1} = \mathbf{M}u_t/w - u_{t-1}$  exactly, for any  $\theta_t$ : a time turn uniform over the body changes the record's phase and nothing of its moduli, its form or its modes; at a single Node light read as a turn drives no resonance and no absorption. (b) A constant turn. For  $\theta_t = \theta$  the plane solution  $z = e^{-i\Omega t}$  has, under this section's step,  $2\cos(\Omega - \theta) = 2\cos \omega_b$ , so the sense + rotates at  $\omega_b + \theta$  and the sense - at  $-\omega_b + \theta$ : both shift by  $+\theta$ , their magnitudes split by  $2\theta$ , the energy shift following the record's own sense, Section 3.5's line. (c) The Link phases couple the modes. To first order in  $\theta_a$  the Link read adds  $Vz = i \sum_a R_a[\theta_{i,a} z_{t+a} - \theta_{i-a,a} z_{t-a}]$  at the Node  $i$ , an  $i$  times a real antisymmetric matrix, Hermitian. Write  $z_t = \sum_m c_m(t) \phi_m e^{-i\omega_m t}$  with the  $c_m(t)$  slowly varying (the mode amplitudes, not the kinetic scale  $c_m$ );  $w(z_{t+1} + z_{t-1}) - \mathbf{M}z_t = -2iw \sum_m \sin \omega_m c'_m \phi_m e^{-i\omega_m t}$ , and projecting  $Vz$  on  $\phi_j$  with  $\theta_{i,a}(t) = \theta_L \epsilon_a(i) \cos \omega_L t$  ( $\epsilon_a$  the light's profile across the body;  $\theta_L = |\partial_a b_L|/(\Gamma \omega_L)$ , the Link phase of the gauge of (a) for light's time level  $b_L$  with its gradient, so that  $E_0 = \omega_L \theta_L$  reproduces the force field of S.26 (c)) gives  $\langle \phi_j, V\phi_m \rangle = i \theta_L \cos(\omega_L t) K_{jm}$  with

$$K_{jm} = \sum_{\text{Links } (i, i+a)} R_a \epsilon_a(i) [\phi_j(i) \phi_m(i+a) - \phi_m(i) \phi_j(i+a)],$$

$$K_{mj} = -K_{jm}, \quad K_{mm} = 0,$$

the inter-mode Link current of S.33's  $g$  taken between two modes. So

$$-2iw \sin \omega_j c_j' = i\theta_L \cos(\omega_L t) \sum_m K_{jm} c_m e^{-i(\omega_m - \omega_j)t};$$

the product carries the frequencies  $\omega_m - \omega_j \pm \omega_L$  and resonates at  $\omega_L = |\omega_j - \omega_m|$ , the difference line, for drives below the sum; at  $\omega_L = \omega_j + \omega_m$  the same matrix element couples  $a_m \phi_m e^{-i\omega_m t}$  to the other sense  $b_j \phi_j e^{+i\omega_j t}$ , since a second-order recurrence has both senses as solutions and the one-sense record is initial data:  $a'' = +[\theta_L K_{jm}/(4w \sqrt{\sin \omega_j \sin \omega_m})]^2 a$ , a growth at the rate  $\Omega_R$  (S.16's  $\epsilon_{ij}$  is  $\theta_L K_{jm}/w$  here, one rate in two notations) with S.16's detuned exponents  $\pm \sqrt{\Omega_R^2 - \Delta^2/4}$ , the sum line S.16 itself names; with  $K$  real  $\frac{d}{dt}(|a|^2 \sin \omega_m - |b|^2 \sin \omega_j) = 0$ , the Wronskian kept while the share grows, so the sum line creates content of the opposite sense in pairs, a pair line at  $\omega_j + \omega_m \approx 2\omega_0$ , in nature's names  $2mc^2$ , named and not claimed (the Floquet multipliers at  $\Omega = \pi/2$  for  $\omega = \pi/6, \pi/3$ ,  $\theta_L K/w = \epsilon = 0.01: 0.9849$  and  $1.0153$ , the growth halving with  $\epsilon$ ; at the difference, all four of modulus 1); the one-mode line  $2\omega_b$  of the pace read (S.16) is the degenerate case; and  $K_{mm} = 0$  says a Link phase shifts no single mode at first order, the gauge statement of (a) for the Links. (d) Rabi's rate. At  $\omega_L = \omega_j - \omega_m$  the resonant terms are  $c_j' = -\theta_L K_{jm} c_m/(4w \sin \omega_j)$  and  $c_m' = +\theta_L K_{jm} c_j/(4w \sin \omega_m)$ , so  $c_j'' = -\Omega_R^2 c_j$  with

$$\Omega_R = \frac{\theta_L |K_{jm}|}{4w \sqrt{\sin \omega_j \sin \omega_m}},$$

a full transfer between the two modes every half period at resonance, one transition per click; detuned by  $\Delta$  the transfer generator  $[-i\Delta/2, -g], [g, i\Delta/2]$ , with  $g = \Omega_R$  and  $\Delta$  the detuning, has the eigenvalues  $\pm i\sqrt{g^2 + \Delta^2/4}$ , so  $P(t) = [g^2/(g^2 + \Delta^2/4)] \sin^2(\sqrt{g^2 + \Delta^2/4} t)$ , Rabi's law at the generalised frequency, oscillatory at every detuning with no threshold, the envelope's full width  $4\Omega_R$  in the detuning at half height (the parametric exponents of S.16 belong to the sum line of (c), not here; a finding of two derivations). (e) The matrix element is the dipole. For a uniform gradient,  $\epsilon_a = 1$ , the identity  $\sum_i x_a(i) [\phi_j(\mathbf{M}\phi_m) - (\mathbf{M}\phi_j)\phi_m]_i = 2w(\cos \omega_m - \cos \omega_j) \langle \phi_j, x_a \phi_m \rangle$ , evaluated with  $\mathbf{M}$ 's hopping and  $x_a(i+a) - x_a(i) = 1$ , gives  $K_{jm} = 2w(\cos \omega_j - \cos \omega_m) d_{jm} = -4w \sin \frac{\omega_j + \omega_m}{2} \sin \frac{\omega_j - \omega_m}{2} d_{jm}$  with  $d_{jm} = \langle \phi_j, x_a \phi_m \rangle$  the dipole: the velocity form equals the length form times the line's frequency, the discrete Ehrenfest relation; for a small line  $\Omega_R = \theta_L \omega_L |d_{jm}|/2$ , nature's  $d \cdot E_0/(2\hbar)$  with the field  $E_0 = \omega_L A_0 = \omega_L \theta_L$  and  $\theta_L = |\partial_a b_L|/(\Gamma \omega_L)$  the phase per Link of the vector potential, the rotating-wave factor  $1/2$  included; a static holder level with a gradient across the body ( $\omega_L = 0$ ,  $\nabla b_L \neq 0$ ) mixes the modes without a transition, Stark's shift; a uniform static phase is a gauge. Check: the slowly varying method is S.16's;  $K_{mm} = 0$  for every real mode by antisymmetry; the identity in (e) follows from  $\mathbf{M}$ 's hopping terms alone, the  $S$  term cancelling. What rests on it: the body's click of S.14, one transition at the line  $\omega_j - \omega_i$  taking the share  $T(\sin \omega_j - \sin \omega_i)$ , stands under the rotation read as under the pace read, the difference line being the absorption line under the rotation and the sum a pair line; the transfer rate is the law's  $\Omega_R$  above with the current's matrix element; the fraction absorbed, with the body's write back, stays open as Section 4.3 says. Status: derived, one derivation, the sum and difference assignment of (c) and (d) checked by two; the pace read's one-mode line (S.16) has no analogue under the rotation; clicks.

**Derivation S.42** (the two forces between quanta and their ratio between families, Section 5.1) Inputs: the sign holder's coupling between two quanta,  $\alpha_{\text{law}} = (3\sqrt{3}/8\pi) k/(\Gamma E_s)$  (S.26); gravity's level from a source of  $s_g$  wells per interval,  $3s_g/(4\pi r \Gamma E_g)$  far away, Eq. (11) of the paper (S.25), and the clock's rate  $1 - U$  in it (S.2); the quantum of a family, its energy

$T \sin \omega_s$ , its share  $\sin \omega_s$  in units of  $T$  and its Wronskian  $T/2$ . Steps: (a) The two holders are one kind of family, the kernel  $3/(4\pi r)$  and the same acts of write and read. The sign holder is written by a quantum's Wronskian,  $1/(2E_s)$  per interval for every family, and the content holder by its form,  $\sin \omega_s/E_g$  per interval, so a quantum of the family  $s$  at  $r$  writes the level  $U(r) = 3 \sin \omega_s/(4\pi r \Gamma E_g)$ . (b) A reader quantum of the family  $s'$  has its rotation redshifted by  $U$ , its energy by  $\delta E = -T \sin \omega_{s'} U$ , so the pull is  $F_g = 3T \sin \omega_s \sin \omega_{s'}/(4\pi \Gamma E_g r^2)$ ; the electric force between two quanta by S.26 (c) is  $F_e = 3kT \cos \omega/(8\pi \Gamma E_s r^2)$ ,  $\cos \omega \rightarrow 1$  at a small gap. In the form of a coupling, with light's  $c = 1/\sqrt{3}$  as in S.26 (b), gravity's between two quanta of one family is  $\alpha_g(s) = (3\sqrt{3}/4\pi) \sin^2 \omega_s/(\Gamma E_g)$ , the square of the quantum's energy over  $T$ . (c) The ratio at one family:  $F_e/F_g = \alpha_{\text{law}}/\alpha_g(s) = k E_g/(2E_s \sin^2 \omega_s)$  to the first order in the gap ((b)'s  $\cos \omega_s$  and the finite-gap redshift's  $2 \cos \omega_s/(1 + \cos \omega_s)$ ), S.18, both 1 to  $10^{-21}$  at nature's gaps at the MAGIC bound ( $\omega_p^2/4 = 1.2 \times 10^{-13}$  at LHAASO's  $\omega_p = 6.8 \times 10^{-7}$ ) and 2/3 and 0.8 at [2, 3], the unit of charge ( $T/2$ ) over the quantum's energy ( $T \sin \omega_s$ ), squared, times the universe's two weights  $k$  and  $E_g$  over  $E_s$ . (d) Between two families the weights cancel:  $(F_e/F_g)_e/(F_e/F_g)_p = \sin^2 \omega_p/\sin^2 \omega_e = (m_p/m_e)^2$  with no number, the masses being the energies  $T \sin \omega_s$  (S.27), an identity in the small-gap limit, forced by charge universality and  $F_g \propto m^2$ , no test. Check: nature's  $e^2/(4\pi\epsilon_0 G m_e^2) = 4.17 \times 10^{42}$  and  $e^2/(4\pi\epsilon_0 G m_p^2) = 1.24 \times 10^{36}$ , the ratio  $3.36 \times 10^6$  with these three digits ( $3.37 \times 10^6 = 1836.15^2$  with the paper's four-digit inputs); at the MAGIC bound  $\omega_e < 2.0 \times 10^{-14}$  (S.24),  $1/\sin^2 \omega_e > 2.5 \times 10^{27}$  and  $k E_g/E_s = 2 \times 4.17 \times 10^{42} \sin^2 \omega_e$ , about  $3 \times 10^{15}$  there (about  $10^{24}$  at the slack LHAASO bound,  $\omega_e < 3.7 \times 10^{-10}$ , S.27), its largest value, and smaller at a smaller  $\omega_e$  (what grows at a smaller gap is  $1/\sin^2 \omega_e$ , the ratio the weights must supply per unit); on the lattice the two holders carry two integers and one kernel, and the  $10^{42}$  appears only when a NodeDetector compares the two forces on one quantum, its charge read as its invariant and its mass as its energy. Status: derived from the definitions, an identity and no test (Section 5.1's mark); clicks.

**Derivation S.43** (the atom with the nucleus's angle alone, the record's own level, and the holders the electron does not read, Section 4.2) Inputs: the sign holder's angle of a nucleus of charge  $Z$  at  $r$ ,  $\theta_0 = Z\alpha_{\text{law}} c/r$  by the far kernel (S.26),  $c = 1/\sqrt{3}$  (S.1); a bound electron record of the matter family read into Schrödinger's slow limit (Section 2.2) with the band's inertia  $m^* = 3 \tan \omega_s$  per interval (S.22) and light's  $c = 1/\sqrt{3}$ ; the one write of every record into the one holder (Section 2.2). Steps: (a) Bohr's form with the nucleus's angle alone. The slow limit is the hydrogen-like problem with the coupling  $Z\alpha$ , the inertia  $m^*$  and the speed  $c$ : the levels  $E_n = -m^*(Z\alpha)^2 c^2/(2n^2) = -m^*(Z\alpha)^2/(6n^2)$  per interval, Rydberg's  $1 - 1/n^2$  between them, and the radius  $a_0 = 1/(m^* Z\alpha c) = \sqrt{3}/(m^* Z\alpha)$  Links, which at nature's gap ( $m^* = 3\omega_s$ ) is  $\lambda_e/(Z\alpha)$ ; the lines are the beats of two levels (S.41); the electron's rotation is redshifted by the binding,  $(Z\alpha)^2/2$  of its energy at a small gap, the mass defect as the click layer reads a frequency. (b) The orbital moment. The kinetic term of a plane record in a plaquette flux  $\Phi$  (S.33's covector) gives the orbital splitting  $\Phi m_\ell/(2m^*)$ , one Bohr magneton per unit of angular momentum, Zeeman's known form derived here under S.33's covector with the orbital  $g = 1$ ; the spin term's  $g = 2$  belongs to the spin holder, named not reached in Section 1.2 of the paper and an option in 8.3. (c) The record's own level. The holder of the sign is one row, and a record cannot tell its own write from the nucleus's, so the bound electron reads the nucleus's angle  $Z\alpha/r$  and its own, of the opposite sense and the strength  $1/Z$  of the nuclear one, spread over its own radius: the Hartree self-term, repulsive and at full strength. With  $x = a_0/a$  for a hydrogen-like trial record the mode's frequency over  $Z^2$  times Rydberg's unit is  $\epsilon = x^2 - 2x + (1.25/Z)x$ , the kinetic, the nuclear and the self-term  $J$ ; its minimum is not a stationary state of the law's iteration. The trial mode, a one-parameter variational estimate of Hartree's fixed point

and not the fixed point itself, is the minimum of  $E_{\text{tot}} = x^2 - 2x + (0.625/Z)x$  at fixed norm,  $x = 1 - 0.3125/Z$ , and its frequency, which the click reads, is  $\epsilon = E_{\text{tot}} + (0.625/Z)x$ :  $E_{\text{tot}}/(Z^2 R) = -(1 - 0.3125/Z)^2$  and  $\epsilon/(Z^2 R) = -(1 - 0.3125/Z)^2 + 0.625(1 - 0.3125/Z)/Z$ , with the radius  $a/a_0 = 1/(1 - 0.3125/Z)$ . At  $Z = 1$ :  $E_{\text{tot}} = -0.473$ ,  $\epsilon = -0.043$ , the radius 1.45; at  $Z = 2$ :  $-0.712$ ,  $-0.448$ , 1.19; at  $Z = 10$ :  $-0.938$ ,  $-0.878$ , 1.03; at  $Z = 30$ :  $-0.979$ ,  $-0.959$ , 1.01. The Rydberg ratios return as  $1 - 1.25/Z$ , 95.8 percent at  $Z = 30$ . So a record reading its own sign field is reflected on the lattice in Bohr's form at large  $Z$  only, and not hydrogen, excluded by line (1) of the model's decision: a record reads every sign field but its own, and hydrogen's lines are S.52's. (d) The holders the electron does not read. A nuclear holder the electron reads is excluded by the spectrum: a one-Node proton by the finite-size part of hydrogen's 1s level, 1.1 MHz for nature's radius (the finite-size shift's known form, derived here as the slow limit's own perturbation; the form factor and the muonic radii the comparison), which a point nucleus would lack and the measured Lamb shift resolves at the kilohertz level [29], and by the form factor, 1 at every momentum transfer where nature's falls; a 1 fm cloud bound by a gapped holder at the binding's content 8/938 by muonic hydrogen, 5.2 meV (the measured Lamb shift's own coefficient,  $5.2275 r_p^2$  meV [30] with  $r_p$  in fm, at  $r_p = 1$  fm) against the electronic and muonic radii's agreement [30]; an ungapped holder by the spectrum. So the nuclear holders are read by the nucleon families and not by the electron, the selective read of Section 4.2 (the numbers computed by hand from the law's lines, one derivation). Check:  $\alpha^2/2 = 2.66 \times 10^{-5}$ ;  $(1 - 0.3125)^2 = 0.473$ ,  $0.473 - 0.625 \times 0.6875 = 0.043$ ; at  $Z = 30$  the self-term is 4.1 percent of the binding. Status: (a) and (b) derived in form, two derivations; (c) the finding and its numbers two derivations; (d) a comparison with nature's Lamb shift and muonic radius, the selective read a declaration. Under the two lines of S.52 the self-term of (c) is absent and (a) is the law's; clicks.

**Derivation S.44** (the equivalence principle between families, Section 5.1) Inputs: the fall of a free packet of the pair [num, den] in a pace gradient,  $a = c^2 [2\nu/(1 + \nu)] \nabla U$  with  $\nu = \text{num}/\text{den}$  (S.18), the proton and the neutron two families of the gaps  $\omega_p$  and  $\omega_n$  under the families' reading. Steps:  $2\nu/(1 + \nu) = 1 - \tan^2(\omega_s/2)$  exactly, so two families fall apart by  $[\tan^2(\omega_n/2) - \tan^2(\omega_p/2)]$ ,  $(\omega_n^2 - \omega_p^2)/4$  at small gaps. Two bodies differ in their fall by that times the difference of their neutron fractions: beryllium (5/9) against  $^{48}\text{Ti}$  (26/48) differ by 0.0139, and with  $\omega_n^2 - \omega_p^2 = (1838.68^2 - 1836.15^2)\omega_e^2 = 9,297\omega_e^2$  the Eötvös parameter is  $\eta = 0.0139 \times 2,324\omega_e^2 = 32.3\omega_e^2$ , below  $4.3 \times 10^{-18}$  at the LHAASO bound  $\omega_e < 3.65 \times 10^{-10}$ , under the beryllium against titanium bound of  $10^{-13}$  from the torsion balance [31] (MICROSCOPE's titanium against platinum,  $10^{-15}$ , Touboul et al. 2022 [32], for which the neutron fractions at natural abundances by mass numbers are 0.5409 and 0.6002, the difference 0.0594 and  $\eta = 0.0594 \times 2,324\omega_e^2 = 138\omega_e^2$ ,  $135.6\omega_e^2$  for the single isotopes  $^{48}\text{Ti}$  and  $^{195}\text{Pt}$  at 0.5417 and 0.6000, the bound  $1.8 \times 10^{-17}$  at  $\omega_e < 3.65 \times 10^{-10}$ ; the conclusion holds at either); the electron against the proton falls apart by  $\omega_p^2/4 = 1.2 \times 10^{-13}$  at the same bound, which no click has read (the electron's free fall known to 10 percent [33]). The caveat: S.18's fall is a free packet's; a nucleon bound in the nuclear holder falls as the body's record does, and the body's fall per family is not derived, so the row is the free-packet estimate of the composition dependence, an upper estimate until the bound body's fall is derived. Check:  $(1 - \nu)/(1 + \nu) = \tan^2(\omega/2)$  with  $\nu = \cos \omega$ ;  $(1838.68^2 - 1836.15^2) = 9,297$ ;  $9,297/4 \times 0.0139 = 32.3$ . Status: derived in form, two derivations, the number an upper estimate; clicks.

**Derivation S.45** (the proofs of the theorems of Sections 3 and 4.2) Theorem 1, the direction. Forward,  $\sigma = 1$ , the line is the definition of  $\text{div}$  and  $\text{mod}$ . Backward,  $\sigma = -1$ : write  $Y$

for the right side's first two terms,  $\sum_j R_{ij} \text{arr}_j + S a_{\text{now}}$ . Then  $u = w a_{\text{next}} - Y + r'$  and  $z = -\lfloor u/w \rfloor = \lceil (Y - w a_{\text{next}} - r')/w \rceil$ . From the forward line,  $Y - w a_{\text{next}} - r' = w a_{\text{before}} - r$  with  $0 \leq r < w$ , whose ceiling divided by  $w$  is  $a_{\text{before}}$  exactly; and  $\rho' = u - w \lfloor u/w \rfloor = w a_{\text{before}} - (Y - w a_{\text{next}} - r') = r$ . The theorem holds for the lattice's step at every interval; a NodeDetector's write and a body's window, click or not, are the NodeDetectors' acts outside the reversal, as the lay is; the exact inverse holds between clicks; across a click the write is a declaration.

Theorem 2, what the acts can form. Each act is a linear form of the levels or the floor of one, and  $|\lfloor L(x) \rfloor| \leq |L(x)| + 1$ ; so sums and compositions of the four acts, at declared coefficients, grow at most linearly in the levels, while a product  $\text{now}_i \text{before}_j$  grows as their square. So nothing bilinear comes from the acts alone. The remark of the paper's Section 3.2, that on a bounded domain every integer function is a finite sum of floors of affine forms, in two lines: on a finite set  $D$  of integer points choose an integer vector  $\mathbf{c}$  with  $\langle \mathbf{c}, \mathbf{x} \rangle$  injective on  $D$  (a base larger than  $D$ 's width per coordinate does it) and  $N$  larger than its spread; for each  $\mathbf{a} \in D$  the indicator is two floors,  $\mathbf{1}_{\mathbf{x}=\mathbf{a}} = \lfloor (\langle \mathbf{c}, \mathbf{x} - \mathbf{a} \rangle + N)/N \rfloor - \lfloor (\langle \mathbf{c}, \mathbf{x} - \mathbf{a} \rangle + N - 1)/N \rfloor$  (the difference is 1 at  $\langle \mathbf{c}, \mathbf{x} - \mathbf{a} \rangle = 0$  and 0 at every other value in  $(-N, N)$ ); then  $f = \sum_{\mathbf{a} \in D} f(\mathbf{a}) \mathbf{1}_{\mathbf{x}=\mathbf{a}}$ , a finite sum of floors of affine forms with integer coefficients. A floor of an affine form of floors with a bounded term,  $\lfloor a(\mathbf{L}\mathbf{x} + \mathbf{b}(\mathbf{x})) + c \rfloor$  with  $\mathbf{b}$  bounded, is a  $\mathbf{L}\mathbf{x}$  plus a term bounded by  $|a| \sup |\mathbf{b}| + |c| + 1$ , affine plus bounded again, so every fixed finite composition of the acts is an affine form plus a bounded term.

Theorem 3, the conserved form. Without the rounding the line is  $\mathbf{a}_{\text{next}} + \mathbf{a}_{\text{before}} = \mathbf{M} \mathbf{a}_{\text{now}}$  with  $\mathbf{M}$  the matrix whose diagonal is  $S_i/w$  and whose entries on the six Links are  $R/w$ . Let  $\mathbf{D}$  be the diagonal matrix of the weights  $1/p_i^2$ ; then  $\mathbf{D}\mathbf{M}$  is symmetric iff  $d_i M_{ij} = d_j M_{ji}$  on every Link,  $d_i R_a(i \rightarrow j) = d_j R_a(j \rightarrow i)$ : with  $R_a(i \rightarrow j) = 2 \text{num } p_i^2 q_{ij}^2 / \Gamma^2$  the weight  $1/p_i^2$  cancels the Node's factor on both sides of every Link and the Link's factor  $q_{ij}^2$  is the same from both ends, so  $\mathbf{D}\mathbf{M}$  is symmetric for any content and any tensions. For a symmetric  $\mathbf{K} = \mathbf{D}\mathbf{M}$  the quantity  $Q(\mathbf{x}, \mathbf{y}) = \mathbf{x} \cdot \mathbf{D}\mathbf{x} + \mathbf{y} \cdot \mathbf{D}\mathbf{y} - \mathbf{x} \cdot \mathbf{K}\mathbf{y}$  satisfies  $Q(\mathbf{a}_{\text{next}}, \mathbf{a}_{\text{now}}) = Q(\mathbf{a}_{\text{now}}, \mathbf{a}_{\text{before}})$ : with  $\mathbf{a}_{\text{next}} = \mathbf{M}\mathbf{a}_{\text{now}} - \mathbf{a}_{\text{before}}$ ,  $\mathbf{a}_{\text{next}} \cdot \mathbf{D}\mathbf{a}_{\text{next}} - \mathbf{a}_{\text{next}} \cdot \mathbf{K}\mathbf{a}_{\text{now}} = \mathbf{a}_{\text{before}} \cdot \mathbf{D}\mathbf{a}_{\text{before}} - \mathbf{a}_{\text{before}} \cdot \mathbf{D}\mathbf{M}\mathbf{a}_{\text{now}}$ , which is the other side. Multiplied by  $w$  this is Eq. (4) exactly.

The proposition, the stable body (Section 4.2). With the dilation  $\lambda$  about the body and  $K, V_r$  the body's own energies at  $\lambda = 1$ ,  $F(\lambda) = K/\lambda^2 - \sum_r \sigma_r V_r \lambda^{-s_r}$ ;  $F'(1) = 0$  is  $-2K + \sum_r \sigma_r s_r V_r = 0$ ;  $F''(1) = 6K - \sum_r \sigma_r s_r (s_r + 1) V_r = \sum_r \sigma_r (3s_r - s_r^2 - s_r) V_r$ : Derrick's scaling, Pohozaev's identity on the lattice.

The momentum flux (the momentum density  $P_a$ , the long version's equation (7)). Substitute the line into  $w P_a(t + 1)$  at coefficients uniform over the lattice: the  $S$  term cancels there (at non-uniform static paces the momentum density gains the Link-wise differences of the coefficients, Section 2.2), the  $w$  term is  $w P_a(t)$  and the  $R$  term is the difference of the fluxes  $G_{ab}$  as the momentum density's form states; the integer step subtracts the remainder term  $\delta_i \Delta_a(\text{now})_i - \text{now}_i \Delta_a(\delta)_i$  with  $\delta = r - r'$  (in this orientation of  $P_a$ ). The guide's lay (S.61 row 17 of the long version; two derivations). (1) A source  $s e^{-i\Omega t}$  at one Node of a width-one guide, whose band is  $\cos \Omega = (\text{num}/3\text{den}) \cos k$  (the transverse mode at  $\cos(\pi/2) = 0$ ), drives two outgoing waves  $C e^{ik|x|}$  with  $C(2 \cos \Omega - (2\text{num}/3\text{den}) e^{ik}) = s$  at the source Node, so  $|C| = 3\text{den } s / (2\text{num} \sin k)$ . (2) The source adds at the interval  $t$  the increment  $A_t \cos(\Omega t)$ ,  $A_t$  its amplitude; the two outgoing waves have  $B = 3A/(2 \sin k)$  on the width-one guide (num = den, the emission's case;  $(3\text{den}/2\text{num}) A / \sin k$  for a general pair), and the form of a travelling wave is  $D = B^2 \sin^2 \Omega$  exactly at every Node, with no time mean; each interval's emission occupies  $v_g = \sin k / (3 \sin \Omega)$  Links on each side, so the two trains' form is  $\sum D = 2(9/(4 \sin^2 k))(\sum_t A_t^2) \sin^2 \Omega \sin k / (3 \sin \Omega) = (3/2)(\sum_t A_t^2) \sin \Omega / \sin k$ , the sum over the increments' squared amplitudes. (3) One quantum on a real line is  $\sum D = T \sin \Omega$

(Section 2.5's packet,  $\sum a^2 = T/\sin \Omega$ ), so one quantum is laid at  $\sum_t A_t^2 = (2/3) T \sin k$ ,  $A_t$  the amplitude of the increment added at the interval  $t$ ; the increments' own squares,  $\sum_t A_t^2 \cos^2(\Omega t)$ , sum to half of it,  $(1/3) T \sin k$ , for an envelope narrow in band against  $\Omega$ , to the cycle average (a short envelope departs from the half; the convention named; two derivations).

Status: Theorems 1 and 3 proved for every lattice, Theorem 2 on the infinite lattice at fixed coefficients; the stable body's proposition derived for the continuous functional, the integer step's version open; lattice.

**Derivation S.46** (the click and the NodeDetector: the passage from the lattice to the clicks, Section 2.5) The click, algebraically. Let  $X$  be the state space: at every Node, for every part of every family, one integer level pair and the write remainders, and nothing else. The step  $S_t$  is Rule3 over the interval with the paces read from the holders at its start, a bijection of  $X$ , exact by the division with its remainder; the line without its division is linear in the levels at fixed paces, the linearity the backward reading uses, the integer map not affine; its forward orbit is the future and its inverse orbit the past. The readings  $D$ ,  $F$  and  $W$  are bilinear forms on  $X$ , covariant under the symmetry group of Section 2.1 and never written. A NodeDetector  $R$  is a declared region (two Nodes or more with Nodes alone, one or more with a record of its own) with its front Ports and its coefficients, which define  $Q_R$ : for one line a quadratic form of the levels, the inflow through its front Ports accumulated over the window; for a pair, the window's sum of the squared cross-side sum over the parts, quadratic in the joint amplitude, the product of the two sides' levels, and so quartic in the levels. A click is the pair  $(R, t)$ , the credit of one whole quantum to the NodeDetector  $R$  at the interval  $t$ , drawn with the probability  $s_R / \sum_{R'} s_{R'}$ , where  $s_R = Q_R(x_t) / W_{\text{rec}}$ , the record's own unit ( $W_c$  for a record laid by share with its count,  $W_c \sin \Omega$  for a born quantum), is the NodeDetector's form on the forward orbit over the quantum's measure of a current. In the meeting's words, the inverse orbit from the credited state coincides with the forward orbit on the realised orbit (in the state space, the record over every path at once), and the share is the bilinear form's value at the coincidence. The click's decision is the one map in the model that is not a bijection, from the vector of shares to one chosen outcome, the same under every element of the group; its write is one act on  $X$ , at the one Node of the region from which the quantum reached the NodeDetector, which the NodeDetector knows and the observer does not. The exchange is this: the arriving record's levels and remainder at that Node to 0, the hole, its count down by one, and the absorbing record's realised part laid there at its count up by one, the absorption, the phase the product's and the remainder at the lay's origin, the quantum conserved between them, the absorbing record's change spreading by Rule3 as an event and the absorbed record erased from that Node outward at the causal bound, inside the click's light cone, nothing else written (ASSUMPTION, the NodeDetector's declaration). Every observable is  $Q_R / W_{\text{rec}}$  on the forward orbit, in the record's own unit ( $W_c$  for a packet laid at count-unit,  $W_c \sin \Omega$  for a born quantum), or a ratio of two such. The one act at three storeys. Rule3 on the levels divides the six weighted arrivals with the Node's own terms and the kept remainder by the wall  $w$ , the quotient the next level, the remainder kept at the Node, the inverse a bijection by it. The hold's write divides the sources' forms by  $E_s T$ , the quotient the held level's gain, the remainder kept at the Node, exact by it. The click divides the inflow through the front over the window, with the books' remainder, by  $W_c$ , the quotient the clicks and the remainder kept in the books for the next window, the two records at the Node its neighbours, the inverse crossed from the click line by the remainder and the count. The count is the level of the top storey; the cancellation lives in the count because the count is the quotient; the exchange at the Node is the writing of the quotient into the two records; the reversibility between clicks is Rule3's; the three holders are

the middle storey and the conversion the top storey's one click; the one thing in the click that is not Rule3 is the draw (DERIVED for the form, two derivations; CLICKS). The hole's booking identity (Section 2.5): with  $wQ(p, q) = \sum_i [w p_i^2 + w q_i^2 - p_i(Sq_i + \sum_j R_{ij}q_j)]$ ,  $w$  times the two-level form  $Q$  of Eq. (4), on a pair of consecutive levels, Rule3's line without the remainder,  $w(\text{next}_i + \text{before}_i) = S\text{now}_i + \sum_j R_{ij}\text{now}_j$ , gives  $wQ(\text{next}, \text{now}) = wQ(\text{now}, \text{before})$  at paces uniform over the lattice, where  $R_{ij} = R_{ji}$ , and then  $wQ(\text{now}, \text{before}) = w \sum_i D_i$  (at non-uniform static paces Theorem 3's weighted form takes  $Q$ 's place); a face writing  $x$  in place of the level  $v$  the step would write at one Node, the leaving level  $b$  standing, changes  $wQ$  by  $(x - v)[w(x + v) - (S\text{now}_i + \sum_j R_{ij}\text{now}_j)] = w(x - v)(x - b)$ , the line giving  $S\text{now}_i + \sum_j R_{ij}\text{now}_j = w(v + b)$ ; exact in the line at any paces as a one-Node identity with the Node's own coefficients, in the plain unit of  $wQ$  (the Node's weight  $1/p_i^2$  relating it to the weighted share), and in integers within  $|x - v|$  in the unit  $Q$ , that is  $w|x - v|$  in the plain unit of  $wQ$ , the carried remainders' term  $(x - v)(r - r')$ , and the most one write removes is  $w(v - b)^2/4$  at the midpoint (three independent derivations). The credit rounds the top storey half up per window; the carried remainder is the law's form (S.61 of the long version, row 12) with its own blind on  $N$ . The window is declared, the universe's draw window set explicitly to the whole passage of a packet, or the credit floors at 0 (Section 3.4); a Node whose pace rounds to 0, a frozen Node, stands in no NodeDetector's front and is never clicked (ASSUMPTION for the definition, DERIVED for the precisions; CLICKS).

One form for every click. At the credit the parts of a record are read with the NodeDetector's coefficients, each part paired with the same part through the common origin, summed and squared; the share is that density, and the draw is by the shares. The two slits are the case with nothing to pair: light is one real line, and the share is the record's own inflow. Bell's check is the case through the common origin: a pair born as one event is two trees with one root, and the backward reading from the first click passes the root and continues forward on the partner's tree (Section 7.1). The joint share is computed from the forward records alone, by the linearity of the rule, the inverse orbit existing by Theorem 1 (DERIVED, CLICKS).

The NodeDetector is the one declaration kind

Assumption (the NodeDetector's declaration). A NodeDetector with Nodes alone is one connected region of two Nodes or more declared in the universe's declaration, never one Node, at least half the wavelength of the family it reports across the beam (4 Links at  $\lambda = 8$  Links); a narrower region, a single Node or a region in pieces is refused at load. A region that absorbs is opaque, backed by a receding face or by the wall's own face, so that what enters it leaves the declared lattice and never returns; a bare region, backed by no face, reports the inflow through its front and absorbs nothing, its net over the window nothing (the paper's Section 2.5). Its click is its report, per interval and family: the net current into the region through its front boundary Ports, the Ports leading in from the declared lattice, summed in integers, with the region's name and the family and never a Node; one wall of the family,  $W_c \sin \omega_0$  for a gapped family and  $W_c$  itself,  $3 \text{ den } T$  for a massless family, of the inflow over a passage is one unit of count, one quantum's share,  $T$  for a bolometer and  $T \sin \omega$  for a counter (S.49 (22) of the long version), so a screen saw  $N$  units where its regions' inflows sum to  $N W_c$ . The credit: one quantum to one NodeDetector, drawn by the shares of the regions' inflows, read outside the lattice over every declared NodeDetector of the universe, one credit per quantum and never two; the NodeDetector draws with the world's declared draw (a body's NodeDetector with its own seed), the lattice draws nothing and the draw is deterministic per seed. The draw is a declared linear congruence of the full period over  $2^b$  values [34], so over the period each outcome's frequency is within  $2^{-b}$  of its weight's share and not exact: the pick is  $(x \times \text{total}) \text{ div } 2^b$  for the draw's state  $x$  and the outcomes' total, and over the full period each residue appears once, so each outcome's frequency is within  $2^{-b}$  of its share,  $2^{-63} = 1.1 \times 10^{-19}$  at the declared

$b = 63$ . After a realised click the NodeDetector writes once, at the one Node from which the quantum reached it. A NodeDetector whose declared universe gives it a record of its own (parts, transitions, a lifetime) is the same kind and emits as well: its region is one Node or more, resolving the place and not the wave, held by its declaration and never bound by a well, and its one write is at the Node the draw picks by the share, the absorption's hole at the Node drawn by the arriving record's inflow over the region and the emission's quantum at the Node drawn by its own record's share. The credit is the one non-local place of the description and no line of the law.

Why a region: a one-Node record is every wave number at once, and the region's half-wavelength is the NodeDetector's declaration; so no click names a position finer than a region, and every per-Node number is a diagnostic.

What produces a click is deterministic: the meeting, the current of the amplitudes through a region's boundary. What is not known is which of the possibilities becomes the click: when one possibility is spread over several NodeDetectors, the lattice gives the shares, no line of the law gives which NodeDetector fires, and the NodeDetector draws. Exclusivity, one click per quantum, is derived under the credit's one draw, the NodeDetector's declaration, from the count conserved; Born's rule, the credit's chance equal to the intensity's share, is a declaration of the NodeDetector and not a finding of the model. The uncertainty relation is not a separate rule: it is the NodeDetector's declarations, the region (its Nodes), the universe's draw window and the draw (its seed), the write exact inside at a Node and an interval that are the lattice's alone and never reported, the report uncertain outside (ASSUMPTION). What each declaration costs: a region of  $n$  Nodes reads a fringe of period  $\Lambda$  with the factor  $\sin(\pi n/\Lambda)/(n \sin(\pi/\Lambda))$ ; a frequency is read to  $\Delta\omega \geq 2\pi/n_w$  over  $n_w$  intervals; and  $N$  quanta drawn by the shares scatter as  $\sqrt{NP_i(1 - P_i)}$ , so a pattern is in the histogram and never in one click (DERIVED, CLICKS; S.8).

The click decides in the credit and writes at one Node

Assumption (the click's decision and its write). The lattice is deterministic and local: Rule3 with no draw, Theorem 1's bijection at every interval. The click is the meeting of two quanta at the NodeDetector, the incoming record's quantum and the NodeDetector's own transition; the NodeDetector, declared in Universe24's declaration, draws with the world's declared draw (a body's NodeDetector with its own seed) whether the meeting is realised, and the randomness is the NodeDetector's, the lattice has none. Forward, to the future, one write: the exchange at the one Node from which the quantum reached the NodeDetector, which the NodeDetector knows and the observer does not (the uncertainty principle: for us the click is a region and a window), the Node of the realised write drawn after the outcome by the same draw's next pick, one Node draw per click and none for a candidate not realised, for an absorption by the inflow booked through each Node over the window (the region's share their sum) and for an emission by the record's share over its Nodes, the Node a lattice diagnostic beside the click line; the exchange: the arriving record's levels and remainder at that Node to 0, the hole, its count down by one; the absorbing record's realised part laid at that Node at its count up by one, the absorption, with the phase the product's and the remainder at the lay's origin, the part it leaves laid at its count down by one; the quantum conserved between them, and the absorbing record's change spreads by Rule3 as an event, one Link per interval; no act faster than one Link per interval: the front erases the absorbed record from that Node outward at the causal bound, one shell of Link distance per interval, the hole's act on each shell, the clicked record alone; the other paths are a cancelled future at three levels: for every click there will be, the record's count 0; at every NodeDetector they reach where the declared universe declares a face behind it, their share leaving the lattice there with no credit; on the lattice the front erasing them inside the click's light cone, beyond it their levels remaining, still sourcing the holders they write to, until it arrives; a record's front starts when its count reaches 0. Backward, to the

past, a reading and never a write: the credit's reading through the root, which chooses which tree is the past and gives Bell at a distance, the one non-local place; it cannot be a write, the lattice being deterministic forward.

A single quantum's count stands at 0 after its click, the count conserved and read by the credit through the root, the NodeDetectors' shared other side on the lattice, the Nodes' levels carrying the shares and never the count: antibunching and one quantum one click per record between far NodeDetectors, the credit's one draw over all the NodeDetectors that read one record (ASSUMPTION for the one draw, DERIVED from it for the rest; CLICKS; the pair's 478/169 and the GHZ's  $-4$  with one click per record). Nothing at one NodeDetector changes the amplitudes at another except by Rule3 from the Node of the exchange; no signal passes. A NodeDetector is a declared region of the lattice and stands nowhere outside it: it has two sides, the front where the incoming quantum meets its own transition and the other side where its own record connects to the rest of the lattice, the apparatus's bodies and, through the record it absorbed, that record's root, the root's event, so that two NodeDetectors reading one record are connected through the root, their common past; it can be entangled on both sides. A bound body clicks by quanta from outside meeting its own transition, never by clicks from outside, and only where it is declared a NodeDetector: the future is light's forward record at the body's boundary, the past the body raised by one quantum read backward to the source, and the meeting is the resonance of the body's rotation with the light's (DERIVED, CLICKS; Section 4.3); a NodeDetector with a record of its own writes once at its window's close and only its own record: on a click, the exchange at the one Node the draw picks by the share, the arriving record's levels there to 0 and its own part raised; on no click, the null window's re-lay of its own record at its Nodes in the complement of its outcome set with its count whole, no Node drawn and nothing of the arriving record written. The arriving record is written only at a realised click, once (S.59 of the long version). A spread body's click is the exchange at its entry Node and its settling a reading, named open, and the emission lays the light quantum at the Node drawn by the NodeDetector's own record's share, at the resonance  $\omega_e - \omega_g$ , the share's difference booked and stated explicitly (S.14, S.41); a body in the dark is only the future and still gravitates; emission is the same meeting with the roles reversed, two local writes and one draw of the credit binding them. The Node stays generic, knowing no family and no click; the write is the NodeDetector's act (ASSUMPTION, the NodeDetector's declaration, the law's line of the click's write; the decision's form DERIVED). The lattice holds only the future, every possibility, and has no arrow: Rule3 is a bijection at every interval (Theorem 1). The clicks have the arrow: each is forward only, the one act outside the reversal; at each click what was future becomes past, the present being the meeting; the chain of clicks is the past advancing into the future, written once and never undone. The direction of time is the clicks' and not the lattice's (THEOREM for the step; the clicks' arrow the NodeDetector's declaration). The two slits' run was made with NodeDetectors that report and write nothing, and the write at the window's end leaves its numbers unchanged.

The passage from the lattice to the clicks. Every formula of Section 5.1 is derived in three steps: the law's line in the lattice's words; each word traded for a NodeDetector's report of it ( $\omega$  for a clock's rate,  $U$  for the ratio of two clocks' rates); and what remains, a relation among clicks alone, set beside nature's. The credit's bookkeeping is that the shares sum to what the screen saw, one whole per click, the whole the NodeDetector's own transition; a frequency of nature is read as the inflow's beat within the window or as the resonance of the NodeDetector's own record, and the lattice's quantities, the form, the content, the level, the rotation, are read by no click except as the share's inflow. Status: the click's algebraic definition and the NodeDetector's declarations the law's; the credit a declaration; lattice for the record, clicks for every number of nature.

**Derivation S.47** (why there is entanglement, and its scope, Sections 2.5 and 7) Bell: the declared universe. *The entanglement the model computes is one event with two trees. The pair is laid as the pair family, two real lines of light never summed at a Node, with equal parts. Each side's NodeDetector, a region at least half a wavelength across, reports the two parts' signed level sums over its region and knows nothing of angles; the ports enter at the credit. The NodeDetector applies the declared setting, a pair of integers  $(p, q)$ , with the port coefficients  $e(+) = (p, q)$  and  $e(-) = (-q, p)$ , exactly orthogonal, so each side's marginal is one half for any setting. The settings of the Clauser-Horne-Shimony-Holt test are  $(1, 0)$ ,  $(1, 1)$ ,  $(12, 5)$  and  $(5, 12)$ , at 0, 45, 22.62 and 67.38 degrees. The credit is the one form of the click: the joint share pairs the parts by their label across the two sides through the root,*

$$J(o_A, o_B) = \sum_{\text{window}} \left| \sum_k c_k e_k(o_A) \alpha_k(o_A) e_k(o_B) \beta_k(o_B) \right|^2, \quad c = (1, 1), \quad (\text{S.47a})$$

with  $o_A, o_B$  the two sides' ports (+ or -),  $\alpha_k$  and  $\beta_k$  the part  $k$ 's amplitude at the port on each side and  $e_k(o)$  the port's integer coefficient on the part  $k$ , the sum over the window's intervals standing outside the square, on both members of the level pair. The joint share is never the square of a local sum, which one NodeDetector could form alone and which gives  $S$  at most 2. One draw is made per pair, the NodeDetectors' draw with the declared seed, the lattice drawing nothing: the first side from its marginal and the partner from the conditional; each side's write at the Node of its exchange spreads by Rule3 and changes no number of the row, one click per record (S.46) (ASSUMPTION, CLICKS).

Bell: what the model says it must give

(1) Rule3 gives each part's amplitudes at the ports: the two parts of one beam, laid equal, are two records of one family stepped by the same line along the same path, identical exactly, integer for integer (THEOREM, LATTICE). (2) The meeting with the linearity of Rule3 gives the joint amplitude: the realised click at A's port weights the parts by A's analyser, is read back to the root where the two parts were one event and forward to B, where B's analyser weights them, and the root sums the two branches,  $\sum_k c_k e_k(a) e_k(b) = \cos(a - b)$  for the normalised vectors  $e(a)$  and  $e(b)$  with  $c$  the parts' laid weights,  $(1, 1)$  for the symmetric lay (DERIVED, CLICKS). (3) That the share is the density of the meeting, its square, is Born's rule for a pair: nothing in Rule3 forces it; everything after it, the fractions,  $S$ , GHZ's  $M$ , is derived in the algebra (ASSUMPTION, FITTED; CLICKS). From the three, with the pair's common phase  $\theta$  uniform over the pairs and A's + port at  $a$  and B's at  $b$ ,

$$\begin{aligned} \text{the meeting:} \quad & P(++) = \frac{1}{2} \cos^2(a - b), \quad E(a, b) = \cos 2(a - b), \quad S = 2\sqrt{2}; \\ \text{a local credit by the shares:} \quad & P(++) = \frac{1}{4} + \frac{1}{8} \cos 2(a - b), \quad E = \frac{1}{2} \cos 2(a - b), \quad S = \sqrt{2}; \\ \text{a local credit by the sign:} \quad & E = 1 - 4|a - b|/\pi \quad (|a - b| \leq \pi/2), \quad S = 2; \\ \text{the credit by the contrast:} \quad & E = \cos 2(a - b) \text{ at the efficiency } 2/\pi, \quad S = 2\sqrt{2}, \end{aligned} \quad (\text{S.47b})$$

The second, third and fourth lines follow by averaging over the pair's phase  $\theta$  (DERIVED, CLICKS; S.10). So  $E(a, b) = \cos 2(a - b)$  under the meeting. For the declared pairs it is exact, Lagrange's identity of the NodeDetector on equal parts:  $E = ((p_A p_B + q_A q_B)^2 - (p_A q_B - q_A p_B)^2) / ((p_A^2 + q_A^2)(p_B^2 + q_B^2))$ , so at the four settings  $E = 119/169, -119/169, 120/169$  and  $120/169$  and  $S = 478/169 = 2.8284$  exactly, an efficiency of 1 by construction and the marginal one half on each side whatever the partner's setting (COMPUTED, CLICKS; S.10). For unequal parts,  $r$  their products' ratio at the ports, the same credit gives  $E = \cos 2a \cos 2b + \rho \sin 2a \sin 2b$  with  $\rho = 2r/(1 + r^2)$  and  $S = \sqrt{2}(1 + \rho)$  at the ideal angles (DERIVED; CLICKS). The maximum of  $S$  over the four settings for  $E = \cos 2a \cos 2b + \rho \sin 2a \sin 2b$  is  $2\sqrt{1 + \rho^2}$ , Tsirelson's bound  $2\sqrt{2}$  at  $\rho = 1$  and never the algebraic 4, and the declared pairs'  $478/169 = 2.82840$  sits below  $2\sqrt{2} = 2.82843$  by the rational angles (DERIVED, CLICKS).

What the lattice gives without the declaration is the second and third lines of Eq. (13) of the paper, two results on local credits on the same declared universe (DERIVED under the uniform phase, CLICKS): a credit of each side by the shares of its ports gives  $S = \sqrt{2}$ , and a credit by the sign of its larger port gives the sawtooth, whose sum is exactly 2, Bell's bound. Bell's value therefore sits in the credit rule, a declaration, and is no independent evidence that the rule accounts for entanglement. Nature beside the derivation: the two loophole-free photon tests of 2015 for the pair's row, and  $S = 2.42 \pm 0.20$  in the test of that year on two spins joined at a click [35], the swap kind, which the law names as met by the same form and which is not derived here. The same credit on four lines gives the three-quantum GHZ correlation  $E_3 = \cos 2(a + b + c)$ , every pairwise  $E$  equal to 0 and Mermin's  $M = -4$  where every local credit gives at most 2 (DERIVED under the declared credit, CLICKS; S.38). No signal passes between the sides, each side's marginal being one half; no settings are chosen by the lattice's own past, so a superdeterministic world is not this one.

The scope. Entanglement in the law is a property of the lay: parts laid alike as one event (the pair family, the GHZ family with  $2^{n-1}$  parts for  $n$  NodeDetectors, S.38) share one root, and the credit's joint share pairs their parts through it; Rule3 preserves the lay and, by the real line's linearity within its bounded remainder (the paper's Theorem 2), creates none. What the law meets is therefore the pair laid alike as one event and two multi-part records meeting at one click, where the click is the root through which their parts are paired, the swap of entanglement; no other entanglement is named as met. Status: the joint share a declaration; the correlations under it derived, two derivations; the swap's joint share HYPOTHESIS, with the criterion that separates a transferred quantum from a coupling, the credit's bilinear form after the exchange; clicks.

**Derivation S.48** (the properties of the lattice and the calibration series, in full, Section 8.2) What the lattice fixes with no number. Before any declared universe the lattice fixes: the geometry, a cube of Nodes with an integer state, six Ports and the 48 symmetries with the reflections, so parity is kept and there is one preferred frame; no time-odd Link read proportional to  $V$  exists in a reciprocal real scheme (S.56 (d); S.36 (i) to (iii) of the long version), hence, for a write with no odd Link part, no  $g_{0a}$ , no frame dragging and  $\alpha_1 = -8$ ; the law's write (Section 2.3, S.56) carries the odd part and gives 0 (DERIVED; S.35, S.36 of the long version); the dynamics, one rule per family, explicit, local and bijective with the remainder kept, so reversibility exactly, integer for integer, no noise but the remainders' and no loops, hence no running of  $\alpha$  and no Lamb shift (DERIVED; a finding against nature, S.50 (6)); the conservation, per family and no total across families: the weighted form and so the energy per family at static paces, the weighted Wronskian and so the charge, exactly for any level in time under the time-Link turn, no spontaneous click and so no lifetime (DERIVED); the band, light at  $1/\sqrt{3}$  Link per interval under the causal bound 1, the anisotropy  $k^2/12$  at short wavelength, a gapped holder Yukawa's kernel with the massless constant (DERIVED); the paces, the exponential metric with  $\gamma = 1$  exactly and  $\beta = 1$  as the gap closes, the finest potential  $1/\Gamma$  and the frozen content  $U_f = \frac{1}{2} \ln(2\Gamma)$ ,  $(\Gamma/2) \ln(2\Gamma)$  levels (DERIVED under the one assumption); the holders, each one's field Poisson's with the kernel  $3/(4\pi r)$ , its strength a declared weight and its range its gap (DERIVED); the click layer, the bilinear credit and so the angle it gives, one whole quantum per click and the draw by the shares (ASSUMPTION); and the scale, one scale  $\Gamma$  and pure numbers of order one, every large number a power of  $\Gamma$  or a declaration. In one sentence: the lattice is a machine of forms; it fixes every form and every ratio of order one, conserves, reverses and makes no noise; it fixes no strength, no scale and no large number; those are the universe's.

The calibration series. *Every measured number of an experiment is first written as a click pattern on a declared NodeDetector against a declared clock. The families' pairs and the units  $\Gamma$  and  $T$  would be fixed by that match and by nothing read off the lattice, Universe24's integers being the rule's own; the two ratios the model fixes without it are light's speed of  $1/\sqrt{3}$  Link per interval and the causal bound of 1 (DERIVED for the speed, LATTICE; the bound is postulate (4)). The passage from the lattice to the clicks is S.46's. No coefficient of the declared universe is derived from the lattice; the lattice dictates the form in which each enters and reads none of them. Each is read from one measurement of nature, as the list below sets out: every coefficient of the declared universe is read by one click on a bound body; the lattice fixes the form of the reading; the declared universe holds the integer. Each stands in one of the three kinds the paper's Section 8.2 names:  $T$  and the NodeDetector's unit of credit are the click's, the gaps,  $k$ ,  $E_g$  and  $\Gamma$  the lattice's, read through bound bodies and given by no NodeDetector. The law has no number from the lattice itself, no fixed point of the whole that selects  $\Gamma$  and no ratio of the weights; if such a number exists, it is in the cosmology. One derived check beside the table, the two clocks: a cavity of bodies and an atom's beat shift alike by  $N$  at long wavelength, as nature's clocks do, an atom against an atom to  $2.1 \times 10^{-5}$  [36] and a cavity against an atom to  $1.7 \times 10^{-2}$  [37], a cavity of declared faces at  $N^2$  (DERIVED; CLICKS; S.31).*

Status: each item as marked; lattice for the forms, clicks where a number of nature is named.

**Derivation S.49** (the conservation picture and the credit's bookkeeping, Section 2.4) What is conserved, and where. *On the lattice the conservations are per family: the weighted form of a family while its reads are static (Theorem 3), the Wronskian of a plane family, its total with the weights  $1/p_i^2$ , exactly in the line for any level in time under the time-Link turn (Section 3.5, S.41), the quanta through the Ports, the wave number of a free record and reversibility exactly, integer for integer. Energy is conserved within a family by theorem, between a charged body and its light by the law's line (Section 5.1), and between a body and the holders of the content it is not conserved, the loss being the mass defect (Section 8.1). The lattice conserves no total: the sum of the families' forms is not conserved and not assumed, the share moving with the paces with no current to compensate (Section 2.4 of the paper). The click layer has books of its own: a click credits one whole quantum at the NodeDetector's own whole, drawn by the inflow of the record's share through its front over its window; a frequency of nature, the lattice's quantities and the credit's bookkeeping are read as S.46 states (Sections 2.5 and 4.3), the bookkeeping up to the absorptions from undepleted records, read in the books and printed as the deficit, the lattice's share of such a record standing within its declared tolerance while its count falls. In the clicks there are conservations other than the lattice's, and neither layer keeps a total energy (THEOREM for the lattice's, ASSUMPTION for the credit's).*

The table.

| <i>Layer</i>           | <i>Conserved</i>                                                                     | <i>Scope</i>                                                                                                                               | <i>By what</i>                                                                          |
|------------------------|--------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| <i>lattice</i>         | <i>the weighted form E of a family</i>                                               | <i>per family, paces fixed, in the rational line (the integer step within the remainder term of the share's change, Section 3.4 (S.5))</i> | <i>Theorem 3</i>                                                                        |
| <i>lattice</i>         | <i>the Wronskian W of a plane family, its total weighted by <math>1/p_i^2</math></i> | <i>per family, exactly for any level in time under the time-Link turn</i>                                                                  | <i>S.9, S.33, S.41</i>                                                                  |
| <i>lattice</i>         | <i>the wave number k of a free record</i>                                            | <i>per family, paces uniform over the lattice</i>                                                                                          | <i>the band's translation invariance</i>                                                |
| <i>lattice</i>         | <i>the sum of D over families</i>                                                    | <i>not conserved</i>                                                                                                                       | <i>Section 2.4 of the paper: the share moves with the paces, no current compensates</i> |
| <i>clicks</i>          | <i>the number of quanta, the charge per quantum, the whole per click</i>             | <i>per whole quantum, across families</i>                                                                                                  | <i>the credit's integer unit; the energy line for a charged body and its light</i>      |
| <i>content holders</i> | <i>the body's share</i>                                                              | <i>not conserved: the loss is the mass defect</i>                                                                                          | <i>S.36</i>                                                                             |

On the lattice nothing is assumed: the conservations are theorems of Rule3's form. What is not conserved and not assumed is the sum of all families' energy; the holders are open to their readers and the law has no theorem equating one family's loss with another's gain. The energy line closes it for one sector: for a charged body and its light the energy its clicks read and the holder's form move together under the turn and the write, both up for a body in its own level and both down for a pair of opposite senses, and agree to first order in the turn if and only if  $k_r E_s T \cos \omega_s = k_w \Gamma$ : one energy read twice, by the clicks and by the form, a condition and not an identity, at the vacuum's paces of S.26 (S.26 (d')). For the content holders it is not closed and not to be: a body's count falls in a well, its potential being in its count, while the holder's waves gain a different share of the loss, the mass defect, the energy leaving the books, the law's own statement and nature's reading of bound bodies. Status: the lattice's conservations theorems; the credit's bookkeeping a declaration; the energy line the law's line, two derivations.

**Derivation S.50** (the explicit findings, with their numbers, Section 8.1) *The measurements of nature named below carry their citations here or in the paper's Sections 5.1 and 8.1. Each finding is a place where the law as written and a measurement of nature differ, named with its number and its root; none is smoothed. (1) The lattice's frame: a metric diagonal in the preferred frame,  $\alpha_1 = -8$  against  $|\alpha_1| < 4 \times 10^{-5}$  and, in the standard gauge,  $\alpha_2 = -1$  against  $|\alpha_2| < 4 \times 10^{-7}$  from the Sun's spin and  $1.6 \times 10^{-9}$  from solitary pulsars, conditional on the static read (Eq. (1)'s second line writes the time part and the three axis parts and never an off-diagonal  $T_{ab}$ , so the carried perturbation has  $h_{00}$  and  $h_{aa}$  only, the breathing  $h_{xx} + h_{yy}$  and the lattice-aligned  $+$ ,  $h_{xx} - h_{yy}$ , and no  $\times$ ,  $h_{xy}$ ), and the Nordtvedt parameter  $\eta_N = 4\beta - \gamma - 3 - \alpha_1 + 2\alpha_2/3 = 7.3$  (8.0 without  $\alpha_2$ 's term) against lunar laser ranging's  $|\eta_N| < 10^{-3}$  [38], the  $\alpha_2$  part conditional on the static read; no frame dragging (0 against Gravity Probe B's  $37.2 \pm 7.2$  milliarcseconds per year), no  $\times$  polarisation. (2) The nuclear*

binding by any derived body: the binding fractions under the declared holders miss in two directions, the deuteron by 4 above and iron by 6 below (S.45 of the long version); the size law of one well,  $R \propto 1/A$ , against  $A^{1/3}$  (S.45 of the long version); the bound  $0.76 CW/(\Gamma E)$  per quantum at one Node is conditional on the declared  $W$ , which no click bounds, and is no finding (S.36; LATTICE for the well, CLICKS for the defect it bounds). (3) The neutron's  $878.4 \pm 0.5$  s [22]: no spontaneous click, no rate; the conversion a declared list of the one click act at a declared rate, the table and the rate nature's numbers and nothing of the neutron's mass claimed (Section 8.1). (4) The  $Z$  and parity: no fourth rotation of a three-line record, the step commuting with the reflections. (5) The anomaly  $g - 2$  and Pauli's two per orbital: no loops, no vector giving the exclusion; the law does not support the reduced mass, as S.52 (c) states once (no amplitude over two positions on the lattice; the pair's joint amplitude at the click, in the credit's pairing, S.47), so the isotope shifts and the muonic atoms are outside what it derives: with the nucleus pinned (line (2)) the  $H-D$   $1S-2S$  shift is 0 against nature's measured isotope shift, and muonic hydrogen's levels lack the factor 0.899 of the reduced mass; the fine structure: the band's  $(Z\alpha)^4$  term is the spin-0 form  $n/(l+1/2)$ , splitting hydrogen's  $2p$  from  $2s$  by 29 GHz ( $\alpha^4 m_e c^2 = 1.4490 \times 10^{-3}$  eV, its twelfth 29.2 GHz the Klein-Gordon  $2s$  to  $2p$  split, its thirty-second 10.95 GHz Dirac's  $2p$  split), where nature's hydrogen has Dirac's  $n/(j+1/2)$  ( $2p$ 's splitting 10.9 GHz,  $2s_{1/2}$  and  $2p_{1/2}$  1.06 GHz apart [29]), and the model has no spin holder, so the fine structure, the hyperfine line and the Zeeman anomaly have no line; the many-body correlation is not reproduced by a product of records (the ground levels, computed by hand from the modes: helium 1.4 percent short, positronium 43 percent of nature's) (S.52). (6)  $\alpha$ 's running and the Lamb shift: no loops. (7) The mass ratios of nature not whole numbers, so no particle a cluster of the electron's quanta; the masses declared. (8) The record's own level: hydrogen's  $1s$  at  $-0.043$  of Rydberg's unit at  $1.45 a_0$ , the Rydberg scaling returning at large  $Z$  as  $1 - 1.25/Z$  ( $E_{\text{tot}}$  as  $(1 - 0.3125/Z)^2$ ) (S.43), excluded by line (1) of the model's decision (S.52). (9) The radiated share's coefficient: the time level carries half of Larmor's power in form, the odd lines' part depends on their wall and coupling, which of the two couplings the odd lines carry open; no lifetime number (S.26, S.45 of the long version). (10) At the second order in  $z$  the ball of dust alone decelerates,  $q_0 = \Omega_m/2 > 0$ , where the supernovae read  $q_0 \approx -0.5$  [39][40] (the flat  $\Lambda$ CDM model, cold dark matter with a cosmological constant); the paper's claim is at  $z \ll 1$  only, and no line of the law closes it. (11) The content holders' balance is not a line of the law: a body's loss in a well is the mass defect (S.36, S.49). (12) The electromagnetic binding energy's seat, open: the holders of the content gravitate with their binding, the sign holder's cross term sources nothing as written (Section 5.1, S.36). (13) Bjorken scaling: under the dimension-3 family a hard click reads three fractions summing to one; their dependence on the momentum transfer is uncomputed, the lines' Fourier weights at  $q$  depending on the lines' lay and the read operator not derived (S.37 (c) of the long version), where nature's measurements show logarithmic violations of order ten percent across the measured decades of  $q^2$ ; the momentum sum rule, nature's quarks carrying about half the nucleon's momentum and the gluons the rest, is not reached, the model having no gluon field; what the law does fix is that no pair is created and no coupling runs, the root of (6). (14) Bell violations between separately prepared bodies joined by a coupling or by a transferred quantum (gate-joined ions,  $S = 2.25 \pm 0.03$  [41]; transmons with a transferred photon,  $S = 2.0747 \pm 0.0033$  [42], loophole-free) exceed the local bound the law's credits give for such a join: on the law as written nothing after an exchanged quantum is bilinear in the two bodies' amplitudes, and every later credit is local,  $S \leq 2$ ; the largest discrepancy the paper names between the law as written and measurement, two derivations. The pair laid as one event and the swap of two multi-part records at a click the law meets (S.47). (15) The shelved ion's telegraph: bimodal counts and exponential dark periods, which the law without the NodeDetectors' writes does not give (the ion's world is not computed in this work); read against

their blinds by the body's own emission click at a declared rate (computed, the rate declared), the shelving at the Zeno-limited rate  $\Omega_w^2/R$ , the bright periods' mean  $R/\Omega_w^2$  and the dark  $\tau_D$ , the rate not derived. (16) The sign holder's radiation is a scalar's: a spherically symmetric breathing charge writes light through its second moment (S.40 (c)), where nature's vector field writes none (hydrogen's 2s, metastable, decaying by two photons); the root the holder's time level written as a scalar, the odd lines' part open (S.26 (d')). (17) The weak coupling: with every rotation holder at  $\alpha$  and nature's  $m_W$ ,  $G_F = \pi\alpha/(\sqrt{2}m_W^2) = 2.5 \times 10^{-6} \text{ GeV}^{-2}$  against  $1.166 \times 10^{-5}$ , short by 4.6, the mixing angle's  $1/\sin^2\theta_W$  absent (S.37 (c)). Status: explicit findings; the frame's parameters of a write with no odd Link part ( $g_{0j} \equiv 0$ ):  $\alpha_1 = -8$ ,  $\alpha_2 = -1$ ; the law's write (Section 2.3, S.56) carries the odd part and gives 0; the pair stands in 8.1 as the misses of the write without it; clicks where a number of nature is named.

**Derivation S.51** (the push on a moving record, Section 5.1, Eq. (9)) Inputs: the composed paces,  $p_0/\Gamma = e^{-U}$  and  $p_a/\Gamma = e^{-2U}$  (Eq. (2), the one assumption  $hN = 1$ ); the band at a pace (Eq. (3)) at a small gap and long wavelength,  $E^2 = m^2 + P^2$  with  $E = \omega$ ,  $m = N\omega_0$  and  $P = cN^2k$ , the exponential metric's dispersion ( $g_{00} = -e^{-2U}$ ,  $g_{aa} = e^{2U}$ ,  $\gamma = 1$ ;  $g_{0a} = 0$  for the static source, the odd part the holder's odd lines carry, sourced by the count's flux, S.56, which the push of a static source does not read); the force on a record as the gradient of its rotation at fixed wave number,  $\mathbf{F} = -(\partial\omega/\partial U)_k \nabla U$  (the fall of S.18 is its  $k \rightarrow 0$  case). Steps: (a)  $\partial(E^2)/\partial U = \partial(m^2 + P^2)/\partial U = -2m^2 - 4P^2$ , so  $-(\partial\omega/\partial U)_k = (m^2 + 2P^2)/E = (E^2 + P^2)/E$ ; (b) with  $\beta = P/E = v/(cN^2)$ ,  $v = \partial\omega/\partial k = c^2 N^4 k/\omega$  the group velocity and  $cN^2$  the local light speed,  $\mathbf{F} = E(1 + \beta^2) \nabla U$ ; (c) at  $\beta = 0$  the pull  $E \nabla U$ , over the band's inertia  $m^* = 3 \tan \omega_0$  the fall  $a = f(\omega_0) \omega_0/m^* \nabla U = c^2(U_K/U) \nabla U$ , with  $U_K$  Kepler's potential of S.18 and  $f(\omega_0) = 2(1 - \cos \omega_0)/(\omega_0 \sin \omega_0)$  (S.15),  $0.2667 \nabla U$  at  $[2, 3]$  (S.18, the same number); at  $\beta = 1$  twice the pull, the bending  $4U_b$  (S.20); (d) the units: the paper's momentum is the wave number, so the lattice form is  $(E^2 + c^2 \mathbf{p} \cdot \mathbf{p})/E$  and the covariant one  $(E^2 + P^2)/E$ ; the form  $E^2 + 3\gamma \mathbf{\Pi} \cdot \mathbf{\Pi}$  holds for the covariant variable  $\mathbf{\Pi} = E\mathbf{v}$  alone (the paper's  $\mathbf{p} = E\mathbf{v}/c^2$  is the wave number, and the boxed momentum at the composed paces is  $P = cN^2k = \Pi/(cN^2)$ , one letter per quantity), where  $3\mathbf{\Pi} \cdot \mathbf{\Pi} = E^2\beta^2$  with  $c^2 = 1/3$  at  $N = 1$ . Check: on the exact band at  $[2, 3]$  the rest push over  $E$  is  $f(\omega_0) = 1.0634$ , the redshift's factor ( $\alpha_U = f(\omega_0) - 1 = 0.063$ ), and the coefficient of  $\beta^2$  in the push over its rest value  $f(\omega_0)E$  is  $1.3353 = \gamma_K - \alpha_K/2$  (two derivations), and with the factor  $\omega_0/\tan \omega_0$  the paper's coefficient,  $(\gamma_K - \alpha_K/2)\omega_0/\tan \omega_0 = 1.0045$ ; at  $[999, 1000]$   $1.00017$  and  $1.00067$  ( $1 + \omega_0^2/12$  and  $1 + \omega_0^2/3$ ), both  $\rightarrow 1$ ; the angle's ratio to Newton's at  $\beta = 0.38$  is  $1 + \beta^2 = 1.144$ , the force's  $\gamma_L(1 + \beta^2) = 1.237$ . Status: derived from Eqs. (2) and (3) under the one assumption (assumption, fitted), two derivations; as physics the standard post-Newtonian factor  $1 + \gamma\beta^2$ , where  $\gamma = 1$  follows from  $hN = 1$ , which the bending fixes, and the finite-gap coefficient is the law's own; the chain to Einstein's weak-field forms is the one dispersion read four ways (the bending at  $\beta = 1$ , Newton at  $\beta = 0$ , the redshift and Shapiro's delay by its time part and index, the perihelion by the orbit step of S.19 with its  $k^4$  caveat at the computed pair); clicks, the deflection angle's ratio  $1 + \beta^2$  to Newton's.

**Derivation S.52** (Sections 4.2 and 8.1: the atom under two lines: no record reads its own write of the sign; the nucleon a compact body) Inputs: the hydrogen-like problem of S.43 (a) with the nucleus's angle alone; the self-read of S.43 (c); two lines, the model's decision, recorded in the decisions' document [43]: (1) every record that reads the holder of the sign is one quantum (count 1), writes its own field of the sign (its Wronskian over  $E_s T$ , Rule3 at the paces of its reads) and reads every sign field but its own, the light born by its write living in its field; (2) the holders of the content keep the self-read, the nucleon families read the nuclear holders' fields

including their own, so a nucleon is a compact self-bound body at the declared weight, its centre one classical variable in an outside well. Steps: (a) The chain, by the law's lines and nothing else. The nucleus, one quantum of its family, writes its Wronskian  $T/2$  into the sign holder's field every interval (the one write, Section 2.2; S.9), and the holder's rest far from it is the kernel's, the angle  $\theta_0(r) = 3kZ/(8\pi r\Gamma E_s) = Z\alpha_{\text{law}} c/r$  with  $c = 1/\sqrt{3}$  and  $\alpha_{\text{law}} = (3\sqrt{3}/8\pi)k/(\Gamma E_s)$  (S.25, S.26 (a), (b)). Under line (1) the electron's record reads that field alone, so its Rule3 line is the plane record's step under the time turn  $\theta_0(r)$  (S.33's inputs; the sign holder rotates the two-part record),  $e^{i\theta_t} z_{\text{next}} + e^{-i\theta_{t-1}} z_{\text{before}} = [Sz_{\text{now}} + \sum_a R_a(z_{+a} + z_{-a})]/w$  (the sign convention of the law's document), and nothing else acts on it. Its stationary modes  $z = \phi e^{-i\omega t}$  obey, by the band at small wave number (S.1, S.22:  $\omega = \omega_s + k^2/(2m^*)$  with  $m^* = 3 \tan \omega_s$ ) and the turn's shift of the rotation (S.26 (c), S.41 (b):  $\omega \rightarrow \omega + \theta$  for the sense +, the electron's  $\theta_0(r)$  of the sign that lowers its rotation in the nucleus's level, the attraction's sign the declared sign;  $2 \cos(\omega + \theta_0)$  equal to the band's  $2 \cos \omega(k)$ ), the law's own eigenvalue line on the lattice,  $-\Delta\phi/(2m^*) - \theta_0(r)\phi = (\omega - \omega_s)\phi$  with  $\Delta$  the six-Port second difference; its radial form is solved on 2,000 radial points with no formula put in: the bound modes sit at  $\omega_n - \omega_s = -m^*(Z\alpha_{\text{law}})^2 c^2/(2n^2) = -m^*(Z\alpha_{\text{law}})^2/(6n^2)$  per interval,  $Z^2$  and  $1/n^2$  exact in the small-gap limit, the radius  $a_0 = \sqrt{3}/(m^* Z\alpha_{\text{law}})$  Links; the levels are a lattice reading, the diagnostic behind the click, and the lines nature measures are the beats of two modes read by a click (S.14, S.41), the table below. Into nature's units: the interval of S.27 ( $\omega_e \hbar/(m_e c^2)$ ) and the law's  $\alpha$  read from Coulomb's force (S.26 (b)) turn the mode's depth into  $m_e c^2 \alpha^2/2$  at  $n = 1$ ,  $-1/2$  hartree exactly in the slow limit ( $-0.49995$  on the grid) and  $-1/8$  at  $n = 2$ . What the law computes is the coefficient  $1/2$  and the dependence on  $n$ ,  $Z$  and the family's gap  $m^*$ ; the two inputs,  $\alpha$  and the electron's unit, are the universe's calibrations from clicks other than the spectrum (Coulomb's force, the electron's mass), so the chain is not circular: no formula of the atom enters, the absolute hartree is a comparison through the unit and the coefficients and the ratios are the derived content. (b) Without (2) the nucleus is a free record of inertia  $m_p^* = 1,836 m_e^*$  reading the electron's field, a well 27 eV deep and  $a_0$  wide: its stationary spread is  $0.21 a_0$  on 2,000 radial points, the nucleus's kinetic share its own reading (the 1s cloud's curvature  $k = 4/3$  in atomic units (a.u.),  $\omega_{zp} = \sqrt{k/m_p} = 0.027$  a.u.), and the self-consistent product of the two records sits at  $-0.4671$  Ha (hartree), 6.58 percent above Bohr, the 2s at  $-0.1147$ , the 1s-2s at 9.59 eV against 10.20; the spread scales as  $m^{-1/4}$ , so deuterium's 1s-2s differs from hydrogen's by 172 meV against nature's 2.8 meV. (c) Under (2) the nucleus does not spread, (a) holds, and the law does not support the reduced mass: it holds no amplitude over two positions, no wave of two records on pairs of Nodes (every record's levels stand at its own Nodes, jointly with the other records there and coupled to them by the read, and the pair's joint amplitude enters at the click, in the credit's pairing, S.47; postulates 1 and 2's locality), so a product of two bound records about one centre has no relative motion of the pair, and under (2) the nucleon's centre is one classical variable, so the isotope shifts and the muonic atoms are outside what the law derives; helium in the mean field without exchange  $-2.8617$  against  $-2.9037$  Ha [44]; positronium, two free records of one gap about one centre,  $-0.217 R$  ( $-0.1085$  Ha, Pekar's constant) against nature's  $-0.50 R$  ( $-0.25$  Ha, 6.80 eV): a product about a centre has none of the pair's relative motion, which carries the whole binding of two equal records; a named finding. Check: the pinned nucleus reproduces  $-1/(2n^2)$  to  $10^{-4}$  on the 2,000 points; the free-nucleus numbers by the self-consistent two-record Hartree iteration (60 iterations, mixing 0.5); the zero-point estimate  $\frac{3}{2}\omega_{zp} = 0.040$  Ha against the self-consistent 0.0136 plus the smearing's 0.019. Status: derived under the two lines, the model's decision (line (1) ASSUMPTION, FITTED; line (2) ASSUMPTION); two derivations, with two costs named: under (1) the fields at a Node number the universe's charged records, so the bound of postulate 2 is the universe's declaration; under (2) the nucleon's classical centre is the pinned nucleus by declaration, which gives the levels, the reduced mass unsupported by construction

(no amplitude over two positions), the isotope shifts with it. What rests on it: *S.43's finding is replaced in Section 8.1 of the paper by the statement that the law does not support the reduced mass and that the many-body correlation is not reproduced by a product of records, Pauli's exclusion having no line (S.50 (5)); LATTICE for helium and positronium (a comparison and no click claimed); CLICKS for the one-electron lines.*

The atoms' table: the clicks the law derives, with how they were derived; lattice for helium and positronium (a comparison and no click claimed), clicks for the one-electron lines. *Nature's measurements give lines and thresholds, the differences of levels, and each row is the beat of two modes read by a click (S.14, S.41), computed by hand from the modes of steps (a) to (c); a level alone is a lattice reading and stands in the text as the diagnostic behind the click. Every row is derived from the law's lines alone, no equation outside the law entering but the self-consistent iteration's own loop, which solves the law's two lines together (Bohr's, Schrödinger's and Hartree's forms in no column but nature's); nature's column is the measured value with its source, the comparison, entering no derivation. The inputs from nature, named once: the electron's unit of S.27 ( $m_e$  and  $\alpha$ , read from the electron's gap and from Coulomb's force, clicks other than the spectrum; the hartree  $m_e c^2 \alpha^2$ ), so an absolute eV is a comparison through the unit and a coefficient or a ratio is the derived content, and the proton's inertia 1,836 as the family's declared pair. The rows are a supporting finding and no more, derived under the two lines, the model's decision, two independent derivations (the chains and the circularity check, row by row), CLICKS. Not in it: the slow limit of the band only, so the  $(Z\alpha)^2$  order, the Lamb shift and the nuclear size are not in it; the reduced mass unsupported by construction (step (c)); Pauli's exclusion without a line (two records in one mode need none).*

| <i>The click</i>                                                                | <i>Nature measured,<br/>with its source</i> | <i>The law's<br/>click under<br/>the two lines</i>       | <i>How derived, by the law's lines</i>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|---------------------------------------------------------------------------------|---------------------------------------------|----------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Hydrogen's 1s-<br/>2s line</i>                                               | 10.20 eV,<br>2,466,061,413,187<br>kHz [45]  | 10.20 eV                                                 | <i>the beat of the <math>n = 1</math> and <math>n = 2</math><br/>modes of step (a) read by a click, the<br/>share <math>T(\sin \omega_j - \sin \omega_i)</math> (S.14, S.41):<br/>3/4 of the 1s depth derived, the eV<br/>through the unit</i>                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| <i>The ionisation<br/>threshold</i>                                             | 13.598 eV [46]                              | 13.6 eV                                                  | <i>the 1s mode's depth of step (a) read<br/>as a click's share (S.14): the coeffi-<br/>cient 1/2 derived, the eV through the<br/>unit</i>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| <i>He<sup>+</sup>'s 1s-2s<br/>line</i>                                          | 40.81 eV (the<br>He II spectrum<br>[46])    | 40.82 eV<br>( $Z^2 \cdot 3/8$<br>hartree at<br>$Z = 2$ ) | <i>step (a) at <math>Z = 2</math>, the angle<br/><math>Z\alpha_{\text{law}}c/r</math> scaling the two modes by<br/><math>Z^2</math>, the beat 3/4 of the 1s: the <math>Z^2</math><br/>ladder derived as a click, the <math>(Z\alpha)^2</math><br/>order not in the slow limit</i>                                                                                                                                                                                                                                                                                                                                                                                                              |
| <i>Helium's ion-<br/>isation thresh-<br/>old</i>                                | 24.587 eV [46]                              | 23.45 eV, 4.6<br>percent                                 | <i>two electron records in one mode<br/>about the pinned nucleus, each read-<br/>ing the nucleus's field and the other's<br/>and never its own (line (1)), iterated<br/>by the law's write and read to the<br/>fixed point, <math>-2.8617</math> Ha (step (c)),<br/>against He<sup>+</sup>'s one record at <math>-2</math> Ha<br/>exactly: the threshold their difference,<br/>the click; nature's total energy two<br/>thresholds summed and no click; the<br/>correlation the product misses sits<br/>wholly in this click</i>                                                                                                                                                               |
| <i>Positronium's<br/>1s-2s line</i>                                             | 5.10 eV,<br>1,233,607 GHz<br>[47]           | 1.9 eV                                                   | <i>two free records of one gap about one<br/>centre, each reading the other's field,<br/>iterated to the fixed point (step (c)),<br/>the beat of the product's two modes<br/>(<math>-0.1085</math> and <math>-0.039</math> Ha, the diag-<br/>nostic behind the click); a product<br/>about a centre has none of the pair's<br/>relative motion, the ground level at<br/>43 percent of nature's, the line at 37<br/>without line (2) the nucleus's record<br/>of count 1 reads the electron's field<br/>and spreads by <math>0.21 a_0</math> at the inertia<br/>1,836, the two records iterated to the<br/>fixed point (step (b)): the control that<br/>separates line (2) from no line (2)</i> |
| <i>Control:<br/>hydrogen's<br/>1s-2s with the<br/>nucleus a free<br/>record</i> | —                                           | 9.59 eV                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

## The slow limit (Section 2.2)

**Derivation S.53** (Schrödinger's equation as the slow limit of the line, Section 2.2)  
Inputs: *Rule3's line at the vacuum's paces without the remainder,  $a_{\text{next}} + a_{\text{before}} =$*

(num/3 den)  $\sum_{Ports} arr$  (S.1);  $\cos \omega_0 = \text{num}/\text{den}$ ;  $\Delta$  the six-Port second difference,  $\sum_{Ports} a = 6a + \Delta a$ ; the rest rotation at a content,  $\omega_0(U) = \omega_0 - \omega_0 f(\omega_0) U + O(U^2)$  (S.15). Steps: write  $a_t = \text{Re}[\psi_t e^{-i\omega_0 t}]$  with  $\psi$  slowly varying,  $\psi_{t\pm 1} = \psi \pm \psi' + \psi''/2 + \dots$ ; then  $a_{\text{next}} + a_{\text{before}} = \text{Re}[e^{-i\omega_0 t}(2 \cos \omega_0 \psi - 2i \sin \omega_0 \psi' + \cos \omega_0 \psi'')]$ , and the right side is  $\text{Re}[e^{-i\omega_0 t}(2 \cos \omega_0 \psi + (\text{num}/3 \text{den}) \Delta \psi)]$ , since  $(\text{num}/3 \text{den}) 6 = 2 \cos \omega_0$ . The  $2 \cos \omega_0 \psi$  cancels, the band at  $k = 0$ ; dropping  $\psi''$ ,

$$i\psi' = -\frac{\text{num}}{6 \text{den} \sin \omega_0} \Delta \psi = -\frac{\Delta \psi}{2m^*}, \quad m^* = \frac{3 \text{den} \sin \omega_0}{\text{num}} = 3 \tan \omega_0,$$

Schrödinger's free equation on the lattice,  $\hbar = 1$  per interval, the inertia the band's (S.18, S.22: the band's own expansion  $\omega - \omega_0 = k^2/(2m^*)$  is the same statement). At a content  $U$  the same steps with  $\omega_0(U)$  add  $V\psi$  with  $V = -\omega_0 f(\omega_0) U$ , the well lowering the rotation. The three conditions, each with its order: (i) the remainder dropped, below one level per Node per interval (the budget clause), the hard bound per laid mode  $(n-1)/\cos(\omega/2) + 1/\sin \omega + 1$  (the carried remainders a difference sequence, summed by parts; about  $n$  at long wavelength,  $1.73n + 0.33$  at the plane's corner for the massless pair,  $1.60n + 0.42$  at  $[2, 3]$ ; the band's two ends excluded by the division act's two sums and by the even periodic box excluded by the declaration) and the walk  $\sigma\sqrt{n}$ , per band mode  $\sqrt{n/(12(1 + \cos \omega_k))}$ , 2.4 levels at the carrier and 3.0 per Node over the lattice's zone at  $n = 130$ , the bound per laid mode 135.6 at the carrier; (ii) long wavelength, the spatial second difference's  $k^4$  term dropped, of relative size  $k^2/12$  along an axis (the band's own  $k^4$  coefficient relative to its  $k^2$  is  $1/(12 \sin^2 \omega_0)$ , 0.15 at  $[2, 3]$ , the rest of it the envelope's term of (iii)); (iii) slow variation,  $\psi''$  dropped against  $2 \sin \omega_0 \psi'$ , of relative size  $(\omega - \omega_0)/(2 \tan \omega_0)$ , below  $(\omega - \omega_0)/\omega_0 = k^2/(2m^* \omega_0)$ , the Klein-Gordon correction. Within the three the statement is exact algebra of the line; in nature's names it is the slow limit of the discrete Klein-Gordon equation, and the model's own result is  $m^* = 3 \tan \omega_0$ , the potential's coefficient  $f(\omega_0) = 2(1 - \cos \omega_0)/(\omega_0 \sin \omega_0)$  and the lattice's departures named. Check (the line evaluated numerically): a Gaussian packet of amplitude width 40 Links at  $k_0 = 0.08$  on a chain of 4,000 Nodes with the two other axes folded, laid with the exact band's before level and stepped 3,000 intervals by the line at  $[2, 3]$  ( $m^* = 3.3541$ ): the envelope's centre at 71.4 Links against Schrödinger's  $k_0 t/m^* = 71.6$  at  $t = 3,000$  intervals (the exact band's group velocity times  $t$ , 71.4); the density's rms width 32.38 against Schrödinger's  $a\sqrt{1 + (t/(2m^* a^2))^2} = 32.40$  with  $a = 40/\sqrt{2}$ . Status: derived, the slow limit with its three conditions named, one derivation (S.18 and S.22 the same statement), the Check the line's numerical evaluation; lattice.

**Derivation S.54** (the lattice's scale from light's anisotropy and nature's photon bounds, Section 8.1) Inputs: the massless band at the vacuum,  $\cos \omega(\mathbf{q}) = (\cos q_x + \cos q_y + \cos q_z)/3$  (S.1); nature's quadratic Lorentz-violation form  $v/c = 1 - \frac{3}{2}(E/E_{\text{QG},2})^2$  with the Fermi Large Area Telescope's (Fermi-LAT) bound  $E_{\text{QG},2} > 1.3 \times 10^{11}$  GeV for the subluminal case (the gamma-ray burst GRB 090510; Vasileiou et al. 2013 [48], after Abdo et al. 2009 [49]); the electron's rest energy as  $\hbar\omega_0/\tau_{\text{int}}$  at the pair's rotation. Steps: (1) with  $\mathbf{n}$  the unit direction and  $S_4 = \sum_a n_a^4$ ,  $\cos \omega = 1 - k^2/6 + (k^4/72) S_4 + O(k^6)$  and  $\cos \omega = 1 - \omega^2/2 + \omega^4/24$ , so  $\omega^2 = k^2/3 - (k^4/108)(3S_4 - 1) + O(k^6)$ : the coefficient of  $k^4$  is  $-1/54$  along an axis,  $-1/216$  along a face diagonal and 0 along a body diagonal, where  $\cos \omega = \cos(k/\sqrt{3})$  exactly; the group velocity  $v_g/c = 1 - \frac{3}{2}\epsilon_4 k^2$  with  $\epsilon_4 = (3S_4 - 1)/36$ , subluminal everywhere and 0 on the body diagonals. (2) With  $E = \hbar\omega/\tau_{\text{int}}$  and  $k = E\ell/(\hbar c)$ ,  $E_{\text{QG},2} = \hbar c/(\ell\sqrt{\epsilon_4})$ ,  $4.243 \hbar c/\ell$  along an axis; Fermi's bound [48] gives  $\ell < 6.4 \times 10^{-27}$  m (an axis;  $1.0 \times 10^{-26}$  m at the angular mean) and  $\tau_{\text{int}} = \ell/(\sqrt{3}c) < 1.2 \times 10^{-35}$  s. (3) The electron's pair fixes  $\tau_{\text{int}} = \hbar\omega_0/(m_e c^2)$  by substitution and nothing of the law fixes the pair: at Fermi's bound  $\omega_0 \leq 9.6 \times 10^{-15}$ , the gap

$1 - \text{num}/\text{den} \leq 4.6 \times 10^{-29}$ ,  $\text{den} \geq 2.2 \times 10^{28} = 2^{94.1}$  (Cassini's bound alone, at  $2\sigma$ , gives  $\omega_0 < 0.0116$ ; LHAASO's highest photon  $\text{den} \geq 1.5 \times 10^{19}$ ); the proton's gap is  $1836^2$  times the electron's, so one denominator serves both only at  $\text{den} \geq 2^{94}$ . Check: the quartic coefficient by the series and by the exact arccosine at  $k = 0.01$  in the three directions,  $-0.018519$ ,  $-0.004630$ ,  $0$ , and  $d\omega/dk$  at  $k = 0.01$  along an axis,  $v_g\sqrt{3} - 1 = -8.333 \times 10^{-6}$  both ways; two methods. Status: derived, a constraint on any declared universe meant for nature, the law fixing no number; the anisotropy of light's time of flight,  $0$  on the body diagonals to  $1/54$  on the axes, a prediction of the law's own readable only at a Link near the bound; lattice.

## The click's algebra beyond the law (Section 2.5.2)

**Derivation S.55** (the click's algebra beyond the law, Section 2.5.2) Inputs: *Rule3* at the vacuum's paces and its band  $\cos \omega = (\text{num}/3\text{den}) \sum_a \cos k_a$  (S.1); the readings at a Node, the Wronskian  $W = \text{re}_{\text{now}} \text{im}_{\text{before}} - \text{im}_{\text{now}} \text{re}_{\text{before}}$  of a plane configuration and the Link current  $F_{ij} = \text{num}(\text{now}_i \text{before}_j - \text{before}_i \text{now}_j)$  of a real line (Eq. (1), Section 2.2); the invariant  $2A^2 \sin \omega = T$  per quantum and the energy  $T \sin \omega$  (Sections 4.3 and 5.1, the factor 2 between a real line's  $I = A^2 \sin \omega / T$  and the invariant the convention of (b)); light's speed  $c = 1/\sqrt{3}$ ; and the proposition of Section 2.5.2, a line beyond the law's write under its own name, that the write carries the arriving wave vector  $\mathbf{k}$  as a phase twist of the taker's record, the twist  $\delta k_a = \text{sgn}(k_a) (|k_a| \text{div } M)$  per Link along each axis for a record of count  $M$  and the remainder  $r_a = \text{sgn}(k_a) (|k_a| \bmod M)$  carried at the body, so that  $k_a = M \delta k_a + r_a$  with  $|r_a| < M$  and  $r_a$  of  $k_a$ 's sign, the rule odd under  $k_a \rightarrow -k_a$  exactly (the law's div, the floor division of Section 2.2 whose remainder is non-negative, gives  $(-3) \text{div } 7 = -1$  with the remainder 4 against  $3 \text{div } 7 = 0$  with the remainder 3; the twist, a line beyond the law, takes the magnitude's division with the sign carried so that it commutes with the cube's 48, reflections included). Steps: (a) Lemma A. For a plane configuration  $a_i = A e^{i(\mathbf{k} \cdot \mathbf{x}_i - \omega t)}$ , with  $\text{now} = A e^{i\phi}$  and  $\text{before} = A e^{i(\phi + \omega)}$  one interval earlier,  $W = A^2 [\cos \phi \sin(\phi + \omega) - \sin \phi \cos(\phi + \omega)] = A^2 \sin \omega$  at every Node, free of  $\mathbf{k}$ ; for a real line along the axis  $a$  with  $\phi_j = \phi_i + k_a$ ,  $\text{now}_i \text{before}_j - \text{before}_i \text{now}_j = A^2 [\cos \phi \cos(\phi + k_a + \omega) - \cos(\phi + \omega) \cos(\phi + k_a)] = \frac{1}{2} A^2 [\cos(k_a + \omega) - \cos(\omega - k_a)] = -A^2 \sin \omega \sin k_a$ , so  $F_{ij} = -\text{num } A^2 \sin \omega \sin k_a$ , free of  $\phi$ . A phase common to the two levels at a Node leaves  $W$ ; a phase step across a Link moves  $F$ . (b) The norm factor. The one-Node recurrence  $x_{t+1} + x_{t-1} = 2 \cos \omega x_t$  conserves  $Q = x_t^2 + x_{t-1}^2 - 2 \cos \omega x_t x_{t-1}$ : for a real line  $B \cos(\phi - \omega t)$ ,  $Q = B^2 \sin^2 \omega$ ; for a plane configuration of amplitude  $A$ ,  $Q = 2A^2 \sin^2 \omega$ , so  $Q/\sin \omega = 2A^2 \sin \omega = T$  and  $W = A^2 \sin \omega = T/2$  per plane quantum. The identity  $2A \cos \phi = A e^{i\phi} + A e^{-i\phi}$  is therefore exact in the count,  $4A^2 \sin^2 \omega$  on both sides: a real line of amplitude  $2A$  is two plane quanta of opposite sense, neutral,  $W = +A^2 \sin \omega - A^2 \sin \omega = 0$ ; one real quantum has  $B_q^2 = 2A_q^2$ , and a real line of amplitude  $B$  carries  $B^2/(2A_q^2)$  quanta. (c) The twist. Under the proposition the twist  $e^{i\delta \mathbf{k} \cdot \mathbf{x}}$ ,  $\delta k_a = \text{sgn}(k_a) (|k_a| \text{div } M)$ , is a phase common to the two levels at each Node, so by (a) it leaves  $W$  at every Node, the charge kept, and moves the Link current; it keeps the count whole only with the write's own scaling of the amplitude, Section 2.5's  $A^2 = (N+1)T/(2 \sin \Omega)$ ,  $\Omega$  the transition rotation of Section 2.5, taken at the twisted rotation,  $A^2 = (N+1)T/(2 \sin \omega(k/M))$ , the invariant  $2A^2 \sin \omega = T$  per quantum holding at every  $k$  and the rotation's difference  $M[\omega(k/M) - \omega_0]$  being the recoil of (e), the beat's balance, booked by the write, in the click's unit  $T \sin \omega$  it is  $\cos \omega_0$  times that to leading order, the contrast of (f); the integer wave vector, in the wave-vector unit  $2\pi/L_{\text{box}}$  per Link,  $L_{\text{box}}$  the declared box's side in Links, is conserved across the click modulo  $2\pi$  per Link with the remainder carried: for a taker whose books hold  $r_a$  from earlier clicks the division is of  $r_a + k_a$ ,  $\delta k_a = \text{sgn}(r_a + k_a) (|r_a + k_a| \text{div } M)$  and  $r'_a = r_a + k_a - M \delta k_a$ ,

so  $r_a + k_a = M \delta k_a + r'_a$  at every click, the printed form the first click's with  $r_a = 0$ , the remainder in the taker's books (i), while the momentum reading  $F \propto \sin k_a$  is conserved only to the order  $k^3$  and folds at the zone's edge. The wave vector the twist carries is read at the taker's Ports and nowhere else: by (a) the arriving light's Link current along the axis  $a$  is  $F_{ij} = -\text{num } B^2 \sin \omega \sin k_a$  with  $B^2 \sin \omega = T$  per real quantum (b), so  $\sin k_a$  per axis is in the NodeDetector's own reads over its boundary, the integer  $k_a$  its nearest whole in the box's units to the order  $k^3$  of the reading, and no number beyond the reads enters. (d) The band's inertia and kinetic scale. With  $\cos \omega_0 = \text{num}/\text{den} = \nu$  and  $\sum_a (1 - \cos k_a) = k^2/2 + O(k^4)$ ,  $\cos \omega - \cos \omega_0 = -\nu k^2/6$ , so  $\omega - \omega_0 = \nu k^2/(6 \sin \omega_0) = k^2/(2m^*)$  with  $m^* = 3 \sin \omega_0/\nu = 3 \tan \omega_0$ , and  $\omega^2 = \omega_0^2 + c_m^2 k^2 + O(k^4)$  with  $c_m^2 = \omega_0/(3 \tan \omega_0) = c^2 \omega_0/\tan \omega_0$  (S.18, S.21;  $c_m/c = 0.867$  at [2,3]). (e) The recoil. A body of count  $M$  taking the twist  $k/M$  per Link gains the recoil, in rotation,  $M[\omega(k/M) - \omega_0] = k^2/(2Mm^*) + O(M^{-3})$ , its energy in the click's unit  $T \cos \omega_0 k^2/(2Mm^*)$  to the same order, vanishing for a large body; the recoil is even in  $k$  and the twist odd, so after an absorption and an emission at  $\mathbf{k}_{\text{out}}$  the body keeps  $\mathbf{k}_{\text{in}} - \mathbf{k}_{\text{out}}$  exactly, under the inversion of any axis as well. (f) Compton. The click balances the rotations by the beat rule of Section 4.3, the law's own (the born light at  $\omega_e - \omega_g$ , the energy in the click's unit  $T \sin \omega$  following to the order  $\omega_i \omega_j$  while no layer keeps a total, Section 2.4), and under the proposition the wave vector; balanced in the rotations the non-relativistic factor below is  $\omega_0/\tan \omega_0$ , where a balance of the energies  $T \sin \omega$  would give  $\omega_0 \cos^2 \omega_0/\sin \omega_0 = 0.5015$  at [2,3], the two one at nature's gap; with light's band  $\omega_L = c|\mathbf{k}_L|$  and the matter band of (d) in the Klein-Gordon form, a quantum at rest absorbing  $\omega_{L,\text{in}}$  and emitting  $\omega_{L,\text{out}}$  at the angle  $\theta$  obeys  $(1 - c_m^2/c^2)(\omega_{L,\text{in}}^2 + \omega_{L,\text{out}}^2) - 2\omega_{L,\text{in}}\omega_{L,\text{out}}(1 - (c_m^2/c^2)\cos \theta) + 2\omega_0(\omega_{L,\text{in}} - \omega_{L,\text{out}}) = 0$  with  $c_m^2/c^2 = \omega_0/\tan \omega_0$ . At  $c_m^2 = c^2$ , nature's gap, this is  $1/\omega_{L,\text{out}} - 1/\omega_{L,\text{in}} = (1 - \cos \theta)/\omega_0$ , Compton's formula,  $\lambda_{\text{out}} - \lambda_{\text{in}} = (2\pi c/\omega_0)(1 - \cos \theta)$  with  $\lambda = 2\pi c/\omega_L$ , the Compton wavelength  $2\pi/(\sqrt{3}\omega_0)$  Links. At a finite gap the outgoing quantum gains, to first order in  $1 - c_m^2/c^2$ ,  $\delta\omega_{L,\text{out}} = (1 - c_m^2/c^2)c^2 q^2 \omega_{L,C}/(2\omega_0 \omega_{L,\text{in}})$ ,  $q$  the recoil wave vector and  $\omega_{L,C}$  Compton's value; in the non-relativistic regime  $\omega_{L,\text{in}} \ll \omega_0$  this is one factor on the whole shift,  $\lambda_{\text{out}} - \lambda_{\text{in}} = (2\pi/(m^*c))(1 - \cos \theta) = (2\pi/(\sqrt{3}\omega_0))(\omega_0/\tan \omega_0)(1 - \cos \theta)$ , exact at every gap in that regime: the inertia  $3 \tan \omega_0$  against the rest rotation  $\omega_0$ ,  $\omega_0/\tan \omega_0 = 1 - \omega_0^2/3 + O(\omega_0^4) = 1 - \frac{2}{3}(1 - \nu) + O((1 - \nu)^2)$  (0.778 at [2,3] to first order), 0.752 at [2,3] and 1 at nature's gap; the factor 0.752 is conditional on the beat rule, the law's balance of Section 4.3, and not on the click's energy  $T \sin \omega$ , under which it would be 0.5015. (g) Absorption alone. A free quantum at rest absorbing  $\omega_L = ck_L$  with no emission would need  $ck_L = \sqrt{\omega_0^2 + c_m^2 k_L^2} - \omega_0$ , that is  $k_L(c^2 - c_m^2) = -2c\omega_0 < 0$  for  $c_m \leq c$ : forbidden at every gap, the free quantum's one window holding two writes. (h) The zone's edge. A single real quantum making a plane pair needs  $k_L^2(c^2 - c_m^2) \geq 4\omega_0^2$ : impossible at nature's gap, and at [2,3] only at  $k_L \geq 2\omega_0/\sqrt{c^2 - c_m^2} = 5.85$  radians per Link, outside the zone's  $|k_L| \leq \pi\sqrt{3} = 5.44$ . (i) The integer recoilless condition. The twist is an integer division of the magnitude with the sign carried, so  $\delta k_a = \text{sgn}(k_a)(|k_a| \text{div } M) = 0$  whenever  $|k_a| < M$  in the box's unit  $2\pi/L_{\text{box}}$  per Link: a body whose count exceeds the quantum's wave number takes no twist from one click; the remainder  $\text{sgn}(k_a)(|k_a| \bmod M) = k_a$  accumulates at the body, and one whole unit per Link, of  $k_a$ 's sign, is written at every  $M/|k_a|$  clicks on average, the first at the  $\lceil M/|k_a| \rceil$ -th, the same clicks for  $-k_a$  as for  $k_a$  with the kicks opposite, so that opposite remainders cancel, and  $M \sum \delta k_a + r_a = \sum k_a$  exactly over any run of clicks, the booked total conserved at every click and the kick never per click below the threshold; at  $|k_a| = M$  the twist is one unit at every click and the remainder 0. The threshold  $|k_a| < M$  compares a wave number in the declared box's units with a count, so it is a property of the computed world and no formula shared with nature; the box-free statement is the recoil in rotation of (e). (j) Light's wave vector in a well. At static paces Rule3 keeps a wave's frequency, and light's local speed in a

well of potential level  $U$  is  $c/n$  with the index  $n = e^{2U}$  (S.2), so its wave vector is  $k = \omega n/c$  at long wavelength (the paper's Eq. for the lens, the chromatic term  $k^2(e^{6U} - e^{2U})/24$  beyond it; S.28): under the proposition the wave vector passed per click, the twist, is  $n$  times the vacuum's at the same frequency, Minkowski's form of the momentum of light in a medium, and the recoil in rotation  $k^2/(2Mm^*)$  is  $n^2$  times the vacuum's. Check: (a) and (b) evaluated in floating point to  $10^{-15}$ , (d) at  $k = 0.05$ , (f) against the exact quadratic at  $\omega_{L,\text{in}} = 0.02$  and  $\theta = \pi/2$  (the frequency drop's ratio 0.757, the factor 0.752 and the relativistic remainder), (g) and (h) at  $[2, 3]$ , the accumulation of (i) for  $k_a = \pm 3$  and  $M = 7$  (the kicks at the third, fifth and seventh clicks, +1 for +3 and -1 for -3) and at the boundary  $k_a = \pm 7$  (one kick at every click), the books' recurrence stepped for one repeated  $k_a$  (its kicks) and for the mixed sequence 3, -5, 10, 3, 3, -7, 2 against  $M = 7$  (the kicks 0, 0, 1, 0, 1, -1, 0, the books 2,  $7 \cdot 1 + 2 = 9$  the sum) with the identity  $r_a + k_a = M\delta k_a + r'_a$  asserted at every click, the rule odd under  $k_a \rightarrow -k_a$  at  $\pm 3$ ,  $\pm 7$  and  $\pm 10$  against  $M = 7$ , and the two recoils of (e) at  $[2, 3]$ ,  $k = 0.03$  and  $M = 1000$ , in rotation  $1.34164 \times 10^{-7}$  and in the click's unit  $8.9443 \times 10^{-8}$  at  $T = 1$ , their ratio  $\cos \omega_0 = 2/3$ , with the two factors of (f), 0.752 under the beat and 0.5015 under the energy, distinct at  $[2, 3]$  and one at the closing gap; one Node and one Link, no run. Status: (a), (b), (d) derived; (c), (e), (f), (g), (h), (i), (j) derived under the proposition of Section 2.5.2 (hypothesis, fitted); lattice, with S.2 and S.28 for (j), Compton's factor lattice at the computed pair, nature's by construction, no test; (i)'s threshold a property of the declared box and no formula shared with nature. The angular shares of the emitted direction are not derived here.

## The clock family on the Ports (Sections 2.3 and 5.1)

**Derivation S.56** (the clock family on the Ports, Sections 2.3 and 5.1) Inputs: *Rule3* at the paces,  $z_{\text{next}} = \mathbf{M}z_{\text{now}} - z_{\text{before}}$  for a plane configuration written as one complex level  $z$  per Node, with the Link coefficients  $M_{ij} = R_{ij}$ ,  $R_{ij}$  here the normalised hop  $R_a/w = \text{num}/(3\text{den})$  at the vacuum's paces and the wall  $S_i/w$  in the self term, and the weights  $d_i = 1/p_i^2$  of Theorem 3,  $d_i R_{ij} = d_j R_{ji}$  (Section 3.3; S.45); the band  $\cos \omega = (\text{num}/3\text{den}) \sum_a \cos k_a$  at the vacuum's paces (S.1), its group velocity  $v_g = (\text{num}/3\text{den}) \sin k / \sin \omega$  along an axis (S.1) and its inertia  $m^* = 3 \tan \omega_0$  (S.18, S.53); the clock  $p_0 = \Gamma e^{-U}$  and the static metric  $ds^2 = -e^{-2U} dt^2 + e^{2U} dx^2$  under the one assumption  $hN = 1$  (Section 2.3; S.34); the booked count and its change by the currents (Section 3.4, Eq. (5)); the Peierls angle of a space Link written from the holder's odd lines, as magnetism's (S.41); Will's parametrised post-Newtonian form of  $g_{0j}$ ,  $-\frac{1}{2}(4\gamma + 3 + \alpha_1 - \alpha_2)V_j - \frac{1}{2}(1 + \alpha_2)W_j$  at  $\zeta_i = \xi = 0$ , and  $\eta_N = 4\beta - \gamma - 3 - \alpha_1 + 2\alpha_2/3$  [3]; and the law's write of the odd Link part (Section 2.3): the three odd Link parts are three odd lines of gravity's holder, sourced each interval at the matter's Links by the count's flux  $f(F(\text{next}, \text{now}))$  in the count's unit (the flux per interval and not its sum, as the time level is sourced by the count  $N$ ), carried by the holder's own line with its wall, so that for a rigidly moving source their static solution is  $\mathbf{V} = U\mathbf{v}_s$  to first order as  $U$  is Poisson's for  $N$  (no number: one wall for the count and its flux, the one assumption for a moving source); the Port's phase reads the odd line's level along the Link,  $V_{ij} = -V_{ji}$ , and turns the arriving plane record by  $\theta_{ij} = -\theta_{ji} = 4V_{ij}\omega_0$  of the record's family, as the hop's coefficient  $M_{ij} = R_{ij}e^{+i\theta_{ij}}$ . Steps: (a) The conserved form. The Port's phase reads the odd line's level along the Link, the mean of its two ends' levels, as the sign's odd lines are read (the law's record), so that  $\theta_{ij} = -\theta_{ji}$  exactly. With  $\theta_{ji} = -\theta_{ij}$  the weighted matrix  $\mathbf{K} = \mathbf{D}\mathbf{M}$  has  $K_{ji} = d_j R_{ji}e^{-i\theta_{ij}} = \overline{K_{ij}}$ , Hermitian, by Theorem 3's own condition  $d_i R_{ij} = d_j R_{ji}$  and nothing of  $|e^{i\theta}| = 1$ . For  $Q(z, z') = \text{Re}[z^\dagger \mathbf{D}z + z'^\dagger \mathbf{D}z' - z^\dagger \mathbf{K}z']$  and  $z_{\text{next}} = \mathbf{M}z_{\text{now}} - z_{\text{before}}$ :  $z_{\text{next}}^\dagger \mathbf{D}z_{\text{next}} - z_{\text{before}}^\dagger \mathbf{D}z_{\text{before}} = z_{\text{now}}^\dagger \mathbf{M}^\dagger \mathbf{D}\mathbf{M}z_{\text{now}} - 2 \text{Re}(z_{\text{now}}^\dagger \mathbf{M}^\dagger \mathbf{D}z_{\text{before}})$

and  $\text{Re}(z_{\text{next}}^\dagger \mathbf{K} z_{\text{now}}) = z_{\text{now}}^\dagger \mathbf{M}^\dagger \mathbf{D} \mathbf{M} z_{\text{now}} - \text{Re}(z_{\text{before}}^\dagger \mathbf{K} z_{\text{now}})$ , and with  $\mathbf{M}^\dagger \mathbf{D} = \mathbf{K}^\dagger = \mathbf{K}$  the two cancel in  $Q(z_{\text{next}}, z_{\text{now}}) - Q(z_{\text{now}}, z_{\text{before}})$  exactly: Theorem 3 with the phase,  $\theta = 0$  its real case. The same holds for the first-order coefficient  $M_{ij} = R_{ij}(1 + i\theta_{ij})$ , an exact rotation by  $\arctan \theta_{ij}$  times the even magnitude  $\sqrt{1 + \theta_{ij}^2}$ , the same for  $ij$  and  $ji$ ; on the lattice its two integer numerators  $(\Gamma_\theta \text{re}_j - V'_{ij} \text{im}_j, \Gamma_\theta \text{im}_j + V'_{ij} \text{re}_j)$ ,  $V'_{ij} = \Gamma_\theta \theta_{ij}$  an integer over the common divisor  $\Gamma_\theta$ , go over  $w\Gamma_\theta$  in Rule3's one division per Node and per part, as today, with S.33's remainder term; the magnitude scales the hop by  $1 + \theta^2/2$ , an even term of the hop's magnitude with no drag in it. The convention, once: the hop from  $j$  to  $i$  carries the phase  $+\theta_{ij}$  with  $\theta_{ij} = 4V_{ij}\omega_0$  along the current, the symbol  $\cos(k + \theta)$ , so a plane record's wave number shifts by  $+\theta$  and  $\delta\omega = \omega(k + \theta) - \omega(k) = +v_g\theta$  to first order, Einstein's sign ( $H = p^2/2m + 4\mathbf{V} \cdot \mathbf{p}$ , the Lense-Thirring  $2G\mathbf{J} \cdot \mathbf{L}/c^2 r^3$ ); with the opposite assignment  $-v_g\theta$ .

(b) Reversibility. The backward step is  $z_{\text{before}} = \mathbf{M}(V_t) z_{\text{now}} - z_{\text{next}}$  with the same matrix, no inverse rotation; the order that keeps it exact is the count's: read the odd lines' levels  $V_t$ , step, source the odd lines by the flux  $f(F(z_{\text{next}}, z_{\text{now}}))$  of the state after the step,  $f$  the holder's floor division with its remainder kept. Theorem 1 holds as for the time level (the holder's line reversible, the source recomputed from the two held records backward); no accumulator is needed. A flux read directly from the two levels the step reads,  $F(\text{now}, \text{before})$ , would put  $z_{\text{before}}$  into its own coefficient, an implicit step, and Theorem 1 would fail; the flux of the state after the step is the law's. (c) Linearity and locality. The step is linear in the record at the written current (the current quadratic in the record, the paces' own status), one Link's reach, explicit, the arriving now and before only; the clock  $p_0$  and the rulers  $p_a$  are untouched. (d) The odd part cannot be a pace. For a one-part real line  $a_{\text{next},i} = \sum_j c_{ij} a_{\text{now},j} - \sum_j d_{ij} a_{\text{before},j}$  the symbol at the wave number  $k$  is  $\lambda^2 - C_1(k)\lambda + C_0(k) = 0$ ,  $C_1$  and  $C_0$  the step symbol's coefficients, both roots unimodular at every  $k$  only if  $|C_0| = 1$  and  $C_1/\sqrt{C_0}$  is real with  $|C_1/\sqrt{C_0}| \leq 2$ . An odd real coefficient  $\epsilon$  on the neighbour's level gives  $C_0 = 1$  and  $C_1 = C_1^{(0)} + 2i\epsilon R \sin k \notin \mathbb{R}$  ( $C_1^{(0)}$  its value at  $\epsilon = 0$ ), so the roots cannot both be unimodular ( $\lambda_2 = 1/\lambda_1 = \overline{\lambda_1}$  would make  $C_1 = 2\text{Re } \lambda_1$  real): to first order  $|\lambda| = 1 + \epsilon R \sin k / \sin \omega = 1 + \epsilon v_g$  and no frequency shift, the number the phase delivers as the shift  $+\theta v_g$  turned by  $i$ . On the neighbour's change now – before it gives  $C_0 = 1 + 2i\epsilon R \sin k$ ,  $|C_0| > 1$ , one root growing,  $\delta\omega = -\epsilon R \sin k$  and  $|\lambda| - 1 = \epsilon R \sin k \tan(\omega/2)$ ; on the neighbour's next (time-centred) the roots are unimodular but the step implicit. So local, explicit and exact are two of three for a one-part line whose odd part is to be proportional to  $V$ : a theorem (the one exception the staggered step, a shift of the before level by whole Nodes, whose drag is quantised and not  $V$ 's), the five forms of S.36 of the long version its instances; the two-part record escapes it because its coefficient is Hermitian and not real, the symbol  $2R \cos(k + \theta) + S$  real and  $|\lambda| = 1$ . (e) The number. A uniform  $\theta$  along an axis shifts the wave number,  $\delta\omega = \omega(k + \theta) - \omega(k)$  exactly under the convention of (a),  $+v_g\theta$  to first order; on a ring the two senses split by  $2v_g\theta$  (the Sagnac form). For slow matter  $v_g = k/m^*$ , so under the delay form  $\theta = 4V\omega_0$ ,  $\delta\omega = +(k/m^*) 4V\omega_0 = +4c_m^2 \mathbf{V} \cdot \mathbf{k}$  with  $c_m^2 = \omega_0/m^*$  (S.21), that is  $\delta\omega/\omega_0 = +4\mathbf{V} \cdot \mathbf{v}$  with  $\mathbf{v} = \mathbf{k}/m^*$ , with no factor of the gap, for every family, to first order in  $V$  and  $k$ ; the share form  $\theta = 4V \sin \omega_0$  would carry  $\sin \omega_0/\omega_0$  (0.886 at [2, 3]) and is not the law's. Einstein's shift of a slow mass's energy by  $+4\mathbf{V} \cdot \mathbf{p}$  has this form. (f) The coefficient 4 from the law's own exponents. A source moving at  $\mathbf{v}_s$  carries the static write boosted: with  $dt_r = dt - \mathbf{v}_s \cdot d\mathbf{x}$  and  $d\mathbf{x}_r = d\mathbf{x} - \mathbf{v}_s dt$  to first order (the band's own boost at long wavelength and small gap, S.33 (c)), the cross term of  $-e^{-2U} dt_r^2 + e^{2U} d\mathbf{x}_r^2$  is  $2(e^{-2U} - e^{2U}) \mathbf{v}_s \cdot d\mathbf{x} dt = -8U \mathbf{v}_s \cdot d\mathbf{x} dt + O(U^2, v_s^2)$ , so  $g_{0j} = -4U v_{s,j} = -4V_j$  with  $\mathbf{V} = U\mathbf{v}_s$ ; the flat part cancels,  $2(1 - 1) = 0$ . The 4 is  $2 + 2\gamma$ , the clock's exponent and the ruler's, equal at  $\gamma = 1$  ( $hN = 1$ ), the exponent of the Link's coefficient relative to the clock's ( $R_{ij} \propto p_a^2 = p_0^4 q^2 / \Gamma^4 \propto e^{-4U}$ ), Will's  $2(1 + \gamma)$ . The odd lines' source  $U\mathbf{v}_s$  carries no number

because the count flows at the group velocity: a packet's count through a Link per interval is the Node's count times  $v_g = v_s$  to first order (the continuity of the conserved share, Theorem 4, Section 3.4); read as the raw Wronskian,  $F_{ij}/W = -\text{num} \sin k_a \approx -\text{num} m^* v_s$  would bring num and  $m^*$  in. (g) The parameters. The harmonic form  $g_{0j} = -4V_j$  is the standard post-Newtonian form  $-\frac{7}{2}V_j - \frac{1}{2}W_j$  under the relabelling  $x^0 \rightarrow x^0 + \frac{1}{2}\partial_t \chi$ ,  $\chi$  the superpotential, a relabelling of the reading and not of the lattice; so  $\alpha_1 = \alpha_2 = 0$ , and at nature's gap,  $\beta = \gamma = 1$ ,  $\eta_N = 0$ ; at the computed pair  $\beta_K = 1.25$  and  $\gamma_K = 1.5$  give  $\eta_N = 0.5$ , the finite gap's own departure. The write without the odd Link part has  $g_{0j} \equiv 0$ , so  $7 + \alpha_1 - \alpha_2 = 0$  and  $1 + \alpha_2 = 0$ :  $\alpha_2 = -1$ ,  $\alpha_1 = -8$  (S.35), the misses the paper prints. (h) The two predictions. Light is a one-part real line with  $\omega_0 = 0$ : nothing to rotate, by (d) no  $V$ -proportional real odd coefficient allowed, so light is not dragged, the Lense-Thirring deflection of light near a rotating body absent, where frame dragging has been read on matter alone (gyroscopes, satellites, pulsars); and a Port cannot read a record's own frequency without reading the record (c), so the phase is the family's  $\omega_0$  times the delay and the coupling of a fast record falls as  $\omega_0/\omega = 1/\gamma_L$ ,  $\gamma_L$  the Lorentz factor, against Einstein's  $+4\mathbf{V} \cdot \mathbf{p}$  at every speed; every measured drag is slow. A free one-part matter line (the scalar entry, dimension 1), which nature's metric would drag, is likewise undragged, untested; bound bodies, plane records of the sign holder, are dragged, which is where every measurement sits. Check: algebra and a nine-Node numerical check of the symbol's roots: (a) by the three identities above; (d) by the roots of the quadratic symbol at  $\epsilon = \theta = 0.02$  and  $k = 0.5$ , light  $[\Gamma, \Gamma]$  and matter  $[2, 3]$ , against the closed forms: on the level  $|\lambda| = 1.01136$  and  $1.00278$  against  $1 + \epsilon v_g = 1.01130$  and  $1.00277$ ; on the change  $\delta\omega = -3.18 \times 10^{-3}$  and  $-2.131 \times 10^{-3}$  against  $-\epsilon R \sin k = -3.20 \times 10^{-3}$  and  $-2.131 \times 10^{-3}$ ,  $|\lambda| - 1 = 4.7 \times 10^{-4}$  and  $1.00 \times 10^{-3}$  against  $\epsilon R \sin k \tan(\omega/2) = 4.6 \times 10^{-4}$  and  $1.00 \times 10^{-3}$ ; time-centred and the phase  $|\lambda| = 1.000000$ , the phase's  $\delta\omega = 1.129 \times 10^{-2}$  and  $2.82 \times 10^{-3}$  against  $v_g \theta = 1.130 \times 10^{-2}$  and  $2.77 \times 10^{-3}$ ; (e) the ring's splitting at  $[2, 3]$ ,  $\theta = 0.01$ :  $2v_g \theta = 0.001742$ ,  $0.003224$ ,  $0.004531$  at  $k = 0.3, 0.6, 1.0$ , equal to  $\omega(k + \theta) - \omega(-k + \theta)$  to six digits; the centred difference  $(\omega(k + \theta) - \omega(k - \theta))/(2 \cdot 4V_k)$  at  $V = 10^{-3}$ ,  $k = 0.1$  is  $0.2500$ ,  $v_g \omega_0/k$ , against  $c_m^2 = 0.2508$ , the  $k^3$  term of  $v_g$  lowering it by 0.3 percent (the forward and backward differences  $0.2542$  and  $0.2458$ , the second order in  $\theta$ ); (f) 72 arcseconds (the perihelion at  $\beta_K = -1$ , S.19, S.34) and 39.2 milliarcseconds per year (Gravity Probe B, S.35 (c)) are not this derivation's; (g) the gyroscope's precession in the harmonic form, Gravity Probe B's predicted 39.2 milliarcseconds per year against its measured  $37.2 \pm 7.2$  [16], the coefficient S.35 (c)'s; the Hermitian form on a periodic chain of 9 Nodes with random paces in  $[0.8, 1.0]$  and angles in  $[-0.3, 0.3]$  over 2,000 steps conserved to a relative  $3 \times 10^{-15}$  for both the exact and the first-order hop (a check, not a result). Domain: every plane record (the charged families and the bound bodies made of them) at any pace; a real line (light; the scalar entry of matter) carries no phase; first order in the source's velocity and in  $V$ ; the count's flux booked in the count's unit as the content is, the odd lines its carrier. Status: derived under the one assumption, the law's write of the odd Link part; (a) to (c) theorems, (d) a theorem for a  $V$ -proportional odd part; (e) to (g) derived, clicks at nature's gap for the three rows of nature; (h) derived, lattice, untested for the two predictions.

## References

- [1] A. Gonen, Universe24: an integer cellular automaton read through clicks, the formulas it shares with nature and the ones it adds, the long version of this article (2026), in the version deposited with this submission, before the law's write of the odd Link part, <https://github.com/Closer24/Universe24/tree/paper-long-v1.1>.

- [2] G. N. Watson, Three triple integrals, *The Quarterly Journal of Mathematics* os-10, 266 (1939), <https://doi.org/10.1093/qmath/os-10.1.266>.
- [3] C. M. Will, The confrontation between general relativity and experiment, *Living Reviews in Relativity* 17, 4 (2014), <https://doi.org/10.12942/lrr-2014-4>.
- [4] R. Courant, K. Friedrichs and H. Lewy, Über die partiellen Differenzengleichungen der mathematischen Physik, *Mathematische Annalen* 100, 32 (1928), <https://doi.org/10.1007/BF01448839>.
- [5] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration, Fermi Gamma-ray Burst Monitor and INTEGRAL), Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A, *The Astrophysical Journal Letters* 848, L13 (2017), <https://doi.org/10.3847/2041-8213/aa920c>.
- [6] S. I. Pekar, *Untersuchungen über die Elektronentheorie der Kristalle* (Akademie-Verlag, Berlin, 1954).
- [7] E. H. Lieb, Existence and uniqueness of the minimizing solution of Choquard's nonlinear equation, *Studies in Applied Mathematics* 57, 93 (1977), <https://doi.org/10.1002/sapm197757293>.
- [8] V. A. Acciari et al. (MAGIC Collaboration), Bounds on Lorentz invariance violation from MAGIC observation of GRB 190114C, *Physical Review Letters* 125, 021301 (2020), <https://doi.org/10.1103/PhysRevLett.125.021301>.
- [9] The law of the lattice, the document ALGEBRA.md of the repository [50], in the version deposited with this submission, cited by the name of its section: the lattice and the families (the objects, a family's declaration, the interval, readings and measurements); Rule3 (the line, the direction, the four acts, the paces, the bound, the conserved form); the click (the booking, the count is the record's share, the click is the meeting); the primitives (the conventions and the units); the stable body (what a body is, the well, the functional and the dilation, the surplus leaves, the generator, the velocity); what the law says of nature (the method of derivation, the reading, the hypotheses under their own names, the declarations by name, the postulates, the constants, the rows against nature); what is open.
- [10] Z. Cao et al. (LHAASO Collaboration), Ultrahigh-energy photons up to 1.4 petaelectronvolts from 12 gamma-ray Galactic sources, *Nature* 594, 33 (2021), <https://doi.org/10.1038/s41586-021-03498-z>.
- [11] Z. Cao et al. (LHAASO Collaboration), An ultrahigh-energy gamma-ray bubble powered by a super PeVatron, *Science Bulletin* 69, 449 (2024), <https://doi.org/10.1016/j.scib.2023.12.040>.
- [12] P. J. Mohr, D. B. Newell, B. N. Taylor and E. Tiesinga, CODATA recommended values of the fundamental physical constants: 2022, *Reviews of Modern Physics* 97, 025002 (2025), <https://doi.org/10.1103/RevModPhys.97.025002>.
- [13] W. H. McCrea and E. A. Milne, Newtonian universes and the curvature of space, *The Quarterly Journal of Mathematics* os-5, 73 (1934), <https://doi.org/10.1093/qmath/os-5.1.73>.
- [14] The Event Horizon Telescope Collaboration, First Sagittarius A\* Event Horizon Telescope results. VI. Testing the black hole metric, *The Astrophysical Journal Letters* 930, L17 (2022), <https://doi.org/10.3847/2041-8213/ac6756>.

- [15] H. Yilmaz, New approach to general relativity, *Physical Review* 111, 1417 (1958), <https://doi.org/10.1103/PhysRev.111.1417>.
- [16] C. W. F. Everitt et al., Gravity Probe B: final results of a space experiment to test general relativity, *Physical Review Letters* 106, 221101 (2011), <https://doi.org/10.1103/PhysRevLett.106.221101>.
- [17] L. Shao and N. Wex, New tests of local Lorentz invariance of gravity with small-eccentricity binary pulsars, *Classical and Quantum Gravity* 29, 215018 (2012), <https://doi.org/10.1088/0264-9381/29/21/215018>.
- [18] J. Müller, K. Nordtvedt and D. Vokrouhlický, Improved constraint on the  $\alpha_1$  PPN parameter from lunar motion, *Physical Review D* 54, R5927 (1996), <https://doi.org/10.1103/PhysRevD.54.R5927>.
- [19] K. Nordtvedt, Probing gravity to the second post-Newtonian order and to one part in  $10^7$  using the spin axis of the Sun, *The Astrophysical Journal* 320, 871 (1987), <https://doi.org/10.1086/165603>.
- [20] L. Shao, R. N. Caballero, M. Kramer, N. Wex, D. J. Champion and A. Jessner, A new limit on local Lorentz invariance violation of gravity from solitary pulsars, *Classical and Quantum Gravity* 30, 165019 (2013), <https://doi.org/10.1088/0264-9381/30/16/165019>.
- [21] M. Wang, W. J. Huang, F. G. Kondev, G. Audi and S. Naimi, The AME 2020 atomic mass evaluation (II): tables, graphs and references, *Chinese Physics C* 45, 030003 (2021), <https://doi.org/10.1088/1674-1137/abddaf>.
- [22] S. Navas et al. (Particle Data Group), Review of particle physics, *Physical Review D* 110, 030001 (2024), <https://doi.org/10.1103/PhysRevD.110.030001>.
- [23] N. D. Mermin, Extreme quantum entanglement in a superposition of macroscopically distinct states, *Physical Review Letters* 65, 1838 (1990), <https://doi.org/10.1103/PhysRevLett.65.1838>.
- [24] M. Zych, F. Costa, I. Pikovski and C. Brukner, Quantum interferometric visibility as a witness of general relativistic proper time, *Nature Communications* 2, 505 (2011), <https://doi.org/10.1038/ncomms1498>.
- [25] R. C. Tolman, P. Ehrenfest and B. Podolsky, On the gravitational field produced by light, *Physical Review* 37, 602 (1931), <https://doi.org/10.1103/PhysRev.37.602>.
- [26] S. W. Hawking, Black hole explosions?, *Nature* 248, 30 (1974), <https://doi.org/10.1038/248030a0>.
- [27] S. W. Hawking, Particle creation by black holes, *Communications in Mathematical Physics* 43, 199 (1975), <https://doi.org/10.1007/BF02345020>.
- [28] J. D. Bekenstein, Black holes and entropy, *Physical Review D* 7, 2333 (1973), <https://doi.org/10.1103/PhysRevD.7.2333>.
- [29] N. Bezginov, T. Valdez, M. Horbatsch, A. Marsman, A. C. Vutha and E. A. Hessels, A measurement of the atomic hydrogen Lamb shift and the proton charge radius, *Science* 365, 1007 (2019), <https://doi.org/10.1126/science.aau7807>.
- [30] A. Antognini et al., Proton structure from the measurement of 2S-2P transition frequencies of muonic hydrogen, *Science* 339, 417 (2013), <https://doi.org/10.1126/science.1230016>.
- [31] S. Schlamminger, K.-Y. Choi, T. A. Wagner, J. H. Gundlach and E. G. Adelberger, Test of the equivalence principle using a rotating torsion balance, *Physical Review*

- Letters 100, 041101 (2008), <https://doi.org/10.1103/PhysRevLett.100.041101>.
- [32] P. Touboul et al. (MICROSCOPE Collaboration), MICROSCOPE mission: final results of the test of the equivalence principle, *Physical Review Letters* 129, 121102 (2022), <https://doi.org/10.1103/PhysRevLett.129.121102>.
  - [33] F. C. Witteborn and W. M. Fairbank, Experimental comparison of the gravitational force on freely falling electrons and metallic electrons, *Physical Review Letters* 19, 1049 (1967), <https://doi.org/10.1103/PhysRevLett.19.1049>.
  - [34] T. E. Hull and A. R. Dobell, Random number generators, *SIAM Review* 4, 230 (1962), <https://doi.org/10.1137/1004061>.
  - [35] B. Hensen et al., Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres, *Nature* 526, 682 (2015), <https://doi.org/10.1038/nature15759>.
  - [36] A. Bauch and S. Weyers, New experimental limit on the validity of local position invariance, *Physical Review D* 65, 081101 (2002), <https://doi.org/10.1103/PhysRevD.65.081101>.
  - [37] J. P. Turneure, C. M. Will, B. F. Farrell, E. M. Mattison and R. F. C. Vessot, Test of the principle of equivalence by a null gravitational red-shift experiment, *Physical Review D* 27, 1705 (1983), <https://doi.org/10.1103/PhysRevD.27.1705>.
  - [38] J. G. Williams, S. G. Turyshev and D. H. Boggs, Progress in lunar laser ranging tests of relativistic gravity, *Physical Review Letters* 93, 261101 (2004), <https://doi.org/10.1103/PhysRevLett.93.261101>.
  - [39] A. G. Riess et al., Observational evidence from supernovae for an accelerating universe and a cosmological constant, *The Astronomical Journal* 116, 1009 (1998), <https://doi.org/10.1086/300499>.
  - [40] S. Perlmutter et al. (The Supernova Cosmology Project), Measurements of  $\Omega$  and  $\Lambda$  from 42 high-redshift supernovae, *The Astrophysical Journal* 517, 565 (1999), <https://doi.org/10.1086/307221>.
  - [41] M. A. Rowe, D. Kielpinski, V. Meyer, C. A. Sackett, W. M. Itano, C. Monroe and D. J. Wineland, Experimental violation of a Bell's inequality with efficient detection, *Nature* 409, 791 (2001), <https://doi.org/10.1038/35057215>.
  - [42] S. Storz et al., Loophole-free Bell inequality violation with superconducting circuits, *Nature* 617, 265 (2023), <https://doi.org/10.1038/s41586-023-05885-0>.
  - [43] The decisions, the document HIGHLIGHTS.md of the repository [50], in the version deposited with this submission: the model's design decisions, one line each, cited by the line's name.
  - [44] C. L. Pekeris, Ground state of two-electron atoms, *Physical Review* 112, 1649 (1958), <https://doi.org/10.1103/PhysRev.112.1649>.
  - [45] C. G. Parthey et al., Improved measurement of the hydrogen 1S–2S transition frequency, *Physical Review Letters* 107, 203001 (2011), <https://doi.org/10.1103/PhysRevLett.107.203001>.
  - [46] A. Kramida, Yu. Ralchenko, J. Reader and the NIST ASD Team, NIST Atomic Spectra Database, version 5.11, National Institute of Standards and Technology, Gaithersburg (2023), <https://physics.nist.gov/asd>.
  - [47] M. S. Fee, A. P. Mills, Jr., S. Chu, E. D. Shaw, K. Danzmann, R. J. Chichester and D. M. Zuckerman, Measurement of the positronium  $1^3S_1$ – $2^3S_1$  interval by

- continuous-wave two-photon excitation, *Physical Review Letters* 70, 1397 (1993), <https://doi.org/10.1103/PhysRevLett.70.1397>.
- [48] V. Vasileiou et al., Constraints on Lorentz invariance violation from Fermi-Large Area Telescope observations of gamma-ray bursts, *Physical Review D* 87, 122001 (2013), <https://doi.org/10.1103/PhysRevD.87.122001>.
- [49] A. A. Abdo et al. (Fermi LAT and Fermi GBM Collaborations), A limit on the variation of the speed of light arising from quantum gravity effects, *Nature* 462, 331 (2009), <https://doi.org/10.1038/nature08574>.
- [50] A. Gonen, Universe24: the implementation of one integer rule on a cubic lattice, read through clicks, Zenodo (2026), the concept DOI, the record of this version deposited with this submission, <https://doi.org/10.5281/zenodo.22738745>.