

# Universe24: Physical Relations from a Common Discrete Update Rule

Alon Gonen

Independent researcher

alon@defounder.ai ORCID 0009-0008-2701-4599

September 2026

## Abstract

The paper begins with one algebraic object: the integer group ring of a cyclic group, the phase grain, on a cubic lattice, the GameBoard, whose symmetries are the 48 signed permutations of the axes, their rotations the cube's group of order 24; one rule of six integer operations on that ring, bounded, acts at every Node and interval. Outside the lattice there are detectors and their clicks only; a record passes to a detector at one comparison, the click, the one non-local step, the only read-out. Exact on the lattice: the books, Gauss's law of a free family's flux, the pace of every direction, the click's weight as a positive quadratic form under one axiom of the apparatus, a pair's count marginals, and the CHSH sum as an exact rational of the phase grain,  $181/64$  from 512 to 8192. Recovered in a limit:  $c = 1/\sqrt{3}$ , Born's rule, Tsirelson's value, Young's spacing, and under a shell average the inverse square and Poisson's equation. Every reading is compared with nature by kind, the failures stated as the law's own: three pass, twelve fail, two are bounds, one is not compared. Two relations enter as labelled conjectures from the algebra, the bending's coefficient  $2(1 + \gamma)$  and the atom's  $1/j^2$  ladder; neither is established. The law as built recovers no relativistic dynamics, and its sequential wheel signals in the order of a pair's outcomes though not in their counts. One value,  $S = 181/64$ , lies inside the measured CHSH value at 1.05 standard errors.

## 1 Introduction


**The algebra first.** This paper begins with one algebraic object and states what follows from it. The object is the integer group ring of a cyclic group, the phase grain  $N_\phi$ , carried on a cubic lattice whose symmetries are the 48 signed permutations of the axes, their rotations the group of the cube of order 24 (Theorem 1); the law is six integer operations on that ring, and every result the paper calls exact is an identity of it under declared tables and world conditions (Theorems 2 to 6, the conservation books, Gauss's law, the pace of every direction). From the same object, with the hypotheses on how a click reads it, one chain reaches above the board: under (A1) with (A3), locality Outside, the  $r$ -free lines of Lorentz's factors, and under (A2) their scale and Einstein's step; under (A1) and five assumptions of the law as built, with no (A2), the equivalence principle and, in a limit under a shell average, Newton's inverse square (the paragraphs 'Newton from the small step' and 'The road to each'): reached from the algebra, not derived from nothing, and the law as built meets (A1) and not (A2), so what the runtime shows of that chain is read row by row in Table 2.

**The paper in one page.** Inside a cubic GameBoard one update law of bounded integers propagates records as a beam by algebraic formulas at every Node and interval: one map with a rate and a wall per component, its rates and walls made of six integer operations. Outside, the game above the board, there are detectors and their clicks only, and the beam passes to them at one comparison, the click, the one non-local step and the only read-out; the GameBoard is read in no other way (P11). Physics above the board is arithmetic on counts, a theorem of the click's definition, and the law beneath it is modern algebra: the rules put into the cells are a cyclic group, a group ring, integer matrices, an evaluation at the roots of unity and a shift, and every result the paper calls exact is an identity of that algebra under its declared tables, on the board or in a named limit, compared with the register's readings by kind. Exact on the GameBoard: the books, Gauss's law of a free family's flux, the pace of every direction with its Manhattan bound, the click's weight a quadratic form of power 2 under one axiom of the apparatus, a pair's marginals and its CHSH sum, 181/64 from 512 to 8192; recovered in a named limit:  $c = 1/\sqrt{3}$ , Born's rule, Tsirelson's value  $2\sqrt{2}$  (a limit, not a bound of the construction, whose finite-grain  $S$  lies on both sides of it), Young's spacing, and under an average that covers the shell, the condition part of the claim, Newton's and Coulomb's inverse square, the retarded potential and Poisson's equation. What no rule forces is said in the same ledger: the Lorentz factor (how the clicks arrive at it is the frame, under an amplitude split the law as built lacks), the bending of light, the values of the masses, the harmonic constants. Against nature three readings pass (1a, 2b, 9); twelve fail in the table, eleven in a registered run (1b, 1c, 2a, 3, 4b, 6, 7b, 8a, 8b, 13, 14) and one a declared input refuted (8c), with four more failing by a pin whose run is not made and not in the table (4a, 5b, 11a, 11c); two are bounds (5a, 7a); one is not compared (2c); one is history (12, the lattice pin met under the law before the entry, not a pass against nature); the hypotheses beside rows 6, 13 and 14 change no verdict of the law (Table 2), the failures the law's own. Two relations enter as conjectures from the algebra, labelled so and distinguished from what a detector reads: the bending's coefficient  $2(1 + \gamma)$  on the one constant, and the atom's ladder  $1/j^2$  with Balmer's 27/20; the ring under its rule is read and its constant does not close within the grain, and the ladder is not read on the board, no lattice able to hold the atom's scale being run. The law as built meets (A1) and not (A2), so it recovers no relativistic dynamics; and its sequential wheel signals in the order of the pair's outcomes, though not in their counts. One finite-grain value,  $S = 181/64$ , inside Poh et al. at 1.05 standard errors, a prediction once the wheel meets no-signalling in the order of its outcomes. The method is one platform, by the algebra and by the simulator alike: an Inside step on a family's attributes, the conversion by an emitter and a detector, an Outside form as a ratio of counts, and a detector's reading pinned before the run.

**The general formula.** The law is one map  $\Phi$  applied at every Node of the GameBoard at every interval, per component  $s$  of the state, each with its own rate  $r$  and wall  $d$ :

$$\boxed{s \leftarrow s + r, \quad e \leftarrow \text{sign}(s) \min(\lfloor |s|/d \rfloor, a), \quad s \leftarrow s - e d.} \quad (1)$$

Each component of the state is a bounded integer accumulator with a declared rate and a declared wall: it is translated by its rate, counted in walls (capped at  $a = 1$  for the drive, uncapped elsewhere), and reduced by the walls it holds; every wall counted is an event, a Link crossed, a phase step taken, a count completed, a birth, a push, and nothing else happens. The rates and walls are made of six integer operations and nothing else. They are the translation; the multiplication by a declared integer matrix or bilinear form; the sum in the group ring of the phase circle; a permutation; the evaluation of a record at the roots of unity with its norm; and the division with the remainder kept, together with the comparison that is the event. One comparison, the click, is



one Node and its six Ports, the Links to its six neighbours

Figure 1: The GameBoard in space: a cubic array of Nodes, here  $6 \times 6 \times 6$ , each joined to its six neighbours by Links through the Ports  $\pm x$ ,  $\pm y$ ,  $\pm z$  (the faint lines); one Node and its six neighbours marked, the Nodes one interval away from it, whose convex hull is the octahedron of Figure 3. Nothing sits at a Node beyond the rows there. Drawn from the definitions; no run.

the only read-out. Only a detector's reading is a measurement. Section 2 gives the full definition; every derivation of this paper names which of these rules it starts from.

**The words of this paper.** The *GameBoard* is the cubic array of Nodes, each joined to six neighbours by Links through directional Ports; time is a sequence of intervals. *Inside* is the board and what moves on it, where no one measures; *Outside* is the game above the board, the emitters, the detectors and their clicks, the only place a number is read. A *row* is one message on the board (a Node, a direction, an age, a phase, an amount, a multiplicity and its record); a *record* is the rows born together at one emitter, one quantum. The *click* is a detector's one comparison, which reads a record once and deletes it: the only read-out, and the one non-local step. A *rung* is one integer threshold of the click's ladder over the birth phase (Eq. (9)); the cell the phase falls in is the outcome. Every number carries its *kind*: DETECTOR, a count of clicks, the only measurement; GAMEBOARD, a number of the board's state, a diagnostic and never a result. Figure 1 draws the board in space, Figure 2 the two sides.

**Why physics behaves like modern algebra.** The claim of the paper is that the forms of several different physical phenomena are recovered from this one discrete update law and a small number of assumptions, Inside by derivation and Outside at the clicks, and the reason is stated in three steps that the paper then follows. First, Outside is arithmetic in  $\mathbb{Q}$ : every measured quantity is a count at a click or a ratio of two counts (a velocity is Nodes apart over counts apart; a visibility, a correlation and a redshift are ratios of counts), and nothing measured is a real number. Second, Inside is integer and local: six operations on bounded integers at a Node and its six neighbours, each rule generic, vector and local. Written down, the rules put into the cells are the cyclic group of the phase, the group ring of the arrivals, the integer matrices of the split and the rotation, the




Figure 2: The mechanism, drawn from the definitions and no run. Inside (left): a GameBoard of Nodes joined by Links; an emitter puts one record of two rows on the board, each row on the digital line of its direction at one Link per interval, its phase turned per Link, nothing kept at a Node; a detector is a set of Nodes. Outside (right): the click, one comparison of the record's birth phase  $u$  against the rungs of its ladder (Eq. (9)); the cell  $u$  falls in is the outcome, and the record is deleted. Nothing else is read.

evaluation at the roots of unity and the translation group's shift. So the state is a free  $\mathbb{Z}$ -module and the interval a  $\mathbb{Z}$ -linear map on it (Section 2). Its couplings are tables: the circle's  $C$  and  $S$ , the split's and the rotation's integer matrices, the family table. Born's rule is a coupling computed from  $N_\phi$ , the rungs of Eq. (9), so a click is a comparison of integers. Third, the transformations between click families that preserve the passage of information are the Lorentz group up to scale, the click theorem's [47] (Section 9), so the symmetry Outside has is the algebra's and not a metric put in. Every result the paper calls exact is an identity of that algebra under its declared tables, matrices and family table, which are inputs (Table 1). Each is checked against the register's readings by kind and not chosen to fit a reading; where a form was declared, the paper says so. The decisive question is how much of that the law really forces: if every phenomenon needed a rule that puts its result in beforehand, the law would explain nothing; if the same rules force several results with little freedom, that is the contribution. So the paper keeps one ledger (Table 1): for every result, the rules of Eq. (1) it starts from, the assumptions added with their kind, the freedom left and the ground of the derivation. What can be established now is an explicit model with exact and conditional results and checks after a detector. The paper is built around the distance between that and the broad claim that Outside represents nature.

**How the simulator led to the results.** The simulator ran first and the derivations followed it: every run registered with its expectation written before the run, its world file and its source fingerprint, each registered number traced back to the operations of the map until a derivation stood or the number was marked a finding; a relation defined in advance is a definition or an input. The two roads, the algebra's and the simulator's, are one platform, stated in Section 9 so that a reader can arrive at a formula the same way.

**The reading rule and the notation.** Only a detector's reading is a measurement (the frame's assumption, P11): a click, a record's moments or an external thing's reading, as the world file declares. Every physics name in this paper (Gauss's, Newton's, Coulomb's, Born's, Planck's, Tsirelson's) names the known form that the law's result recovers, Inside by derivation or Outside at a detector. A number of the GameBoard's state is a diagnostic of the host, never compared with nature and never pinned. Every symbol is named in English at its first use, a scalar plain, a vector in bold lowercase, a matrix or an operator in bold uppercase; the symbol table is Appendix B, the program's glossary is on the tree [12], and every number has its one source [5].

**What this paper contributes, and what it does not claim.** New here: one explicitly stated update rule with its implementation, run from world files with every reading pinned before the run; exact conditional algebraic results on the lattice (the quadratic read-out characterization, the count marginals, the CHSH sum as an exact rational of the grain,  $181/64$  on the registered plateau); one platform on which every formula is reached by the algebra and by the simulator alike; and a confrontation register that states the failures with the same care as the passes. Not claimed: a derivation of the Lorentz group from the six operations, the atom's levels as a reading, a unique prediction of  $181/64$  without a rule that fixes the grain, or any comparison with nature beyond the rows of Table 2, each read by kind.

## 2 The law and its implementation

**The GameBoard as a system of information transfer.** The GameBoard is a system of information transfer, and its only objects are messages, the rows: a row is a tuple (Node, direction, age, phase, emitter, content per unit, record, label, amount, multiplicity), and a Link is a channel that carries at most one step of a row per interval in each direction. Nothing else exists Inside: no field at a Node, no register, no memory beyond the rows present. Each interval does six things to messages and nothing else: it moves a row one Link along its digital line (the translation), turns its phase (the translation on the circle), splits it at a splitter into rows with integer weights (the multiplication by a declared table), rotates a labelled pair (an integer matrix), merges the rows that meet at a Node with equal words, opposite phases cancelling (the addition), and, at a detector, reads a record once by one comparison and deletes it (the evaluation and the count). The books say that no message is lost or made in transit: released equals in transit plus absorbed plus escaped plus cancelled at every interval. Two messages of this paper: a row on the heading  $+x$  crosses its Links at the intervals 1, 3, 5, 7, 8 (the hand-worked update below), nothing kept at a Node; and at a balanced splitter one row of amount 1 becomes two of amount 1 and multiplicity 2, which meet again at a second splitter, at one port with equal phases (the words add) and at the other  $N_\phi/2$  apart (the message cancels), so one detector reads the record, 64 of 64 births, and the dark port is the cancelled message (series L, Section 8). Written down, a message is a basis element of a free  $\mathbb{Z}$ -module, the passage is the translation group's shift, the split and the rotation are integer matrices, the merge is the group ring's addition and the read-out is one bilinear form: the passage from physics to modern algebra is the passage from messages with attributes to vectors with coordinates. The lattice gases [52, 53] carry bits on links with collision tables, the cellular automata machines of [27] and the quantum cellular automata of [78] are local rules on cells, and Shannon's channel [7] is the message and its capacity; the GameBoard is of that family, with two differences: its messages carry a phase on a bounded circle and an integer amount, and its only read-out is the click.

**The road from the cells to the algebra.** How the things of physics entered: the amount  $w$  of a row is a positive integer, translated and merged; the phase  $p$  is an element of  $\mathbb{Z}_{N_\phi}$ , turned per Link, added in the group ring, evaluated at the roots of unity; the multiplicity  $\mathfrak{m}$  is a positive integer,  $A = \sum a_i^2$  at a split; the age  $\tau$  is a count at the rate 1; mass is the content  $M$ , a count of units per family and an input of the family table, which sets the drive's wall  $N_l N_w M + |p_a|$  (the division), scales the push's bilinear form and the release rate, and is the rest energy in whole units,  $E'_0 = E_0/c^2 = N_l N_w M$ ; charge is  $\rho$ , the declared charge per unit of content, entering only the coupling matrix  $\rho_A \rho_B$  of the push; momentum is an integer vector, translated by the push, read by the drive as a rate against a wall, its square compared and never rooted; the field a body reads is the label flow  $\mathbf{a} = \sum \text{amount} \times \mathbf{u}_D$ , bilinear in the arrivals; the family table (content, cost per phase step, charge, the strong column, lifetime, phase rate, hand) is read and never computed; and the couplings, the circle's tables at 1/256, the split and rotation tables and the click's Gram matrix, are integer matrices and one bilinear form. Every family of the register is one row of declared integers (P8) and nothing else: the quantum  $\hbar$  (1 on a free family, whose rows carry their energy as turns, 0 on a paid family, whose content is the mass), the charge per unit of content as a rational pair, the columns with their signs, the lifetime, whether the phase turns, the hand; the content  $M$  of a body is the world's declaration. Written down, a family is where its quantities live: the content on  $\mathbb{Z}$ , the non-compact scale, free; the phase on  $\mathbb{Z}_{N_\phi}$ , quantised; the charge and a column's coefficient as entries of the coupling matrix; the hand a pseudoscalar under the 48; a family's rows on the fan under the 48; a pair's arms in  $\mathbb{Z}^2 \otimes \mathbb{Z}^2$  under the settings' integer matrices; and the six verbs are the only operations on them. The mass angle  $m$  of the frame is the rest pair's turn,  $2\pi n/d$  per interval, a world's declaration and not the family's. Table 5 (Appendix D) lists every family of the register with its declared integers, its object, when it entered, the verbs that act on it and the click that reads it; a cell the tree does not name says so.

**Rules, parameters, initial conditions.** The law is stated as rules, apart from the parameters and the initial conditions a world supplies. The rules: (P1) every quantity of the state is a bounded integer and every operation integer arithmetic, no real number, root or floating point at run time; (P2) space is a cubic array of Nodes, each joined to six neighbours by Links through directional Ports, a Node's complete local information its NodeState, time a sequence of intervals; (P3) the rates and walls of the map are made of the six operations of Section 1 and nothing else; (P4) a rule reads its own record and the arrivals at its Node and its six neighbours, already delivered, and nothing kept at a Node beyond the events there, no register, remainder or draw; (P5) no row or body crosses more than one Link per interval; (P6) a record is read out once, by one comparison, at the click, the one step that deletes, and the click of a pair is one gather of one record from both settings, the law's one non-local operation. The parameters (P7, P8): the grains, the circle  $N_\phi$ , the pace's grain  $N_l$ , the sphere of directions, the birth wheel  $N_u$ , the fan's grain, the clock's pair, and the roundings declared at load, among them the circle's tables at 1/256; and the physical inputs, the family table (per family the content, the cost  $\hbar$  per phase step, the charge per unit of content, the strong column, the lifetime, the phase rate and the hand; its instance on the tree [10]) and the world's width  $N_w$ , which carries what physics calls Newton's constant. The initial conditions: the box and its boundaries, the bodies with their momenta and contents, the lamps, the detectors, the splitters and the gates of the world file. The law of this paper is the runtime before the generic entry of the bending: the commits of `main` that every run of this paper names in Appendix C, on the first-parent line of `main` up to the archived law's head 3a9a7109, the last commit before the entry's merge 5b855791; the generic entry (`main` at 5b855791 since 2026-09-22, the flight in the age wall's set at  $1 + \gamma$ ,  $\gamma$  an input [32]) is a later rule of the law, and the paper's only readings under

it are the ring's (Table 2, row 13). Four rules are chosen among few (P9) and are assumptions, not derivations: the flight is blind to the crowd, a row's rate in transit constant and its wall the direction's  $T_D$ ; a body's drive is the flight at the fraction its momentum earns,  $|p_a|/(N_l N_w M + |p_a|)$  per axis, each axis's count advancing every interval, the first axis whose count fires making the step and a later axis's coincident fire lost, its wall subtracted and nothing carried (the axes  $x$  before  $y$  before  $z$ ; the engine's rule [17]), so that a body crosses at most one Link per interval (P5) and the per-axis fraction is its pace only where the fires do not coincide: three equal momenta  $N_l N_w M$  would give  $1/2$  per axis,  $3/2$  in all, and the built body moves on  $x$  alone at  $1/2$ , the sum of a body's paces bounded by 1; the directional drive of the momentum's line, form B [6, record 652], exists as a declared identity off by default (the world key `drive_b`, series Y), no coincident fire lost, read inside its pins on six registered worlds; it becomes the default only by a declared decision of the author, not before; until then a diagonal body is the per-axis walk; the collision's permutation is the cyclic shift; and a clock counts the age moment of the rows at its Node (the potential's form, [6, record 394]) rather than their presence. One axiom of the apparatus (P10): the click stores one pointer per set, the least-rank member of the family Theorem 4 forces, which is Born's form with the fundamental harmonic; which harmonic is a relabelling of the declared rate, and the harmonic constants are not derived. One assumption of the frame (P11): the GameBoard, Inside, is read only through its emitters and detectors, Outside; every result Outside is a click, a number of the board's state is never a reading, and every comparison with experiment is a comparison of clicks with readings; a measurement is itself an event of the board, an emitter putting a record on it and a detector ending the record at the click, the one step that deletes Inside and the one row that Outside gains. Nothing outside this list is assumed.

**The map.** Here  $s$  is one accumulator of the state, a bounded integer on a circle or a line;  $r$  its declared rate, signed, a constant of the world for a message in transit and a reading of what arrives for a body;  $d$  its declared wall, a positive integer;  $a$  the component's cap, 1 where a cap is declared and  $a = \infty$  where none is (the whole part; the primitive's `at_most` = 0); and  $e$  the event count: every unit of  $e$  is an event (a Link crossed, a phase step taken, a count completed, a birth, a push), and nothing else happens; a reversed rate first cancels what was accumulated the other way. For a count without cap the third step is the division toward zero with its signed remainder kept, the Euclidean division for  $s \geq 0$ ; under the cap the residual above the wall is kept and fires one event at each following interval until spent. The cap  $a = 1$  is the drive's alone, one Link per interval; every other count takes the whole part and can fire several events in one interval, the turn, the owed count, the release, the lamp, the push, so that a rate of 5 against a wall of 2 fires 0, 2, 5, 7, 10 events over the intervals, as the high-rate runs read [6, record 340]. The 0/1 form  $e \leftarrow [s \geq d]$ , one event bit per interval, holds exactly where the rate never exceeds the wall: the flight ( $2S_1 N_l \leq 2T_D$ , at most one Link per interval), the drive by its cap and with the sign ( $e \in \{-1, 0, 1\}$ ), the birth wheel (the rate 1 against  $N_u$ ) and a clock whose rate is below its wall (the derivation's section 0 [19]). The rates and walls of every component are made of six operations and nothing else [18, 16]:

1. the translation of an accumulator by its rate (the flight, the phase, the age, every count);
2. the multiplication by a declared integer matrix or bilinear form (a split, a rotation; a coupling, the click's Gram matrix);
3. the addition in the group ring  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$  (the merge, where opposite phases cancel);
4. the permutation of the joint state (a collision, a gate);

5. the evaluation of the record at the  $N_\phi$ th root of unity,  $\mathbb{Z}[\mathbb{Z}_{N_\phi}] \rightarrow \mathbb{Z}[\zeta_{N_\phi}]$  ( $N_\phi$  the phase circle's width), and its norm (the click's one quadratic step);
6. the division toward zero with the remainder kept (the Euclidean division for a non-negative accumulator), and the comparison (the carry that is the event; the cap; the threshold).

Two places are not linear and there is no third: the carry, a bijection  $s \mapsto (e, s - ed)$  with  $|s - ed| < d$  and the remainder's sign that of  $s$ , which loses nothing, and the click's threshold, the one comparison read out per record, whose remainder is not read: the one place the law deletes by design. The cap and the sign are operation 6's comparison,  $\min(\lfloor |s|/d \rfloor, 1) = \lfloor |s| \geq d \rfloor$ ; the capped carry is the injection  $s \mapsto (e, s - ed)$ , which loses nothing either, so there is still no third place. Written as algebra, the six are: the carry a bijection  $s \mapsto (e, s - ed)$ ; the linear block's closed form  $\lfloor s_0 + rt \rfloor$ , the flight the translation group's shift along the digital line; the split's table with  $A = \sum a_i^2$  inverted by its conjugate transpose (Theorem 2); the rotation an integer matrix with  $U_s^\top U_s = n_s I$ ; the merge the addition in  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ , its cancel the quotient by  $x^{N_\phi/2} + 1$ , which for  $N_\phi$  a power of two is the ring of cyclotomic integers  $\mathbb{Z}[\zeta_{N_\phi}]$ , a free  $\mathbb{Z}$ -module of rank  $N_\phi/2$ ; and the click one bilinear form  $\mathbf{f}^\top \mathbf{G} \mathbf{f}$  with  $\mathbf{G}$  the tables' Gram matrix, then one comparison. The exact evaluation at the roots of unity and the evaluation by the tables at  $1/256$  are two maps: the cancel's zero, an equal pair at  $N_\phi/2$ , is a zero of both, the click's other zeros are the tables' own (Section 6), and each claim below says which of the two it is about. Modern algebra is not assumed here; it is what the cell rules are when written down, and every theorem below is a property of these six maps.

**The state.** The GameBoard is a box of Nodes in  $\mathbb{Z}^3$ , each joined to six neighbours by Links through six Ports; an axis may be periodic. Time is the interval  $t \in \mathbb{Z}$ . The state at an interval is a vector  $\mathbf{s}$  of bounded integers on a product of circles and lines, and it is the multiset of the messages in transit (the rows) and the bodies (the measured events), and nothing else: a Node holds no register, remainder or draw. A body carries its Node, its content  $M$ , its momentum vector  $\mathbf{p}$  and its table of counts, one accumulator per count (the fields per kind on the tree [10]).

**The components of Eq. (1).** The ladder's rungs  $b_k$  of Section 6 are walls of this kind. The components and their  $(r, d)$  [19]: a row's Manhattan count (the rate  $2S_1 N_l$ , the wall  $2T_D$ , started at  $T_D$ ;  $D$  the direction vector,  $S_1$  its Manhattan length,  $T_D = \lfloor \sqrt{3|D|^2 N_l^2} \rfloor$  its resolution,  $N_l$  the flight's grain), with three axis deficits choosing the axis of each Link; a row's phase (an integer per Link, or a declared rational  $\nu$  of steps per interval, written  $n/d$  in Section 6, its whole part read modulo  $N_\phi$ ); a row's age (the rate 1, no wall; bounded by the family's lifetime and the run); a body's drive on each axis (the rate  $|p_a|$ , the wall  $N_l N_w M + |p_a|$ ,  $M$  the content and  $N_w$  the world's width), whose speed  $p/(N_l N_w M + p)$  per axis where the axes' fires do not coincide, one Link per interval in all, saturates at one Link per interval, above the rows' pace; a body's counts (the turn, the owed count, the release, the lamp, the push per column, each a rate against a wall); a record's birth phase (the rate 1 per birth, the wall  $N_\phi$ ; the birth wheel, a lamp's declared rate  $r_w$  per birth against the wall  $N_u$ , [2531, 4096] on the registered two-slit world, the rate the nearest odd integer to  $0.618 \times 4096$  [2]); the exact phase at the click [2]; and the click's ladder, one comparison read once. Beside the six operations stand the roundings declared at load, each a constant of the law, among them the tables of the circle at  $1/256$ , the unit vector of a direction to the nearest integer vector,  $T_D$  by the integer square root, the click's rung to the nearest integer and the fan's weights at the grain  $N_\theta$  [16].

**The two blocks.**  $\Phi$  is linear where its rates are constants of the world and feeds back where a rate is a function of what arrives. The linear block is the rows in flight: the flight, the phase and the age at constant rates, the merge, the integer matrices of a split, a rotation or a gate, and the evaluation at the click. Every component of it has the closed form of its unwrapped count,  $[s_0 + rt]$  (the accumulator itself is that count's remainder by the wall), so a row's state at any interval is a formula of its birth and its age, and the limits of Sections 4 and 5 are limits of that formula. The feedback block is the bodies: the push, whose rate is the reader's content times the flow of the arriving rows (bilinear in the state); the owed count, whose rate is what the body reads (the presence, or the age moment); the turn and the release, whose rate is the content; the drive, whose rate is the momentum the push changes. These have no closed form; they are iterated, and their continuum limit is the differential equation  $ds/dt = \mathbf{r}(\mathbf{s})$ . The exact word for the whole map is therefore piecewise-linear in the rows with bilinear rates in the bodies [19].

**The read-out.** The click reads one comparison and nothing else leaves the rows: for a record whose rows have all ended, the apparatus sums the rows' integer phasors per cell through the tables, squares the sums (the one quadratic step, forced in form by Theorem 4), lays the cells on a ladder and compares the record's coordinate  $u$  with the rungs; the cell whose rung  $u$  falls under is the click. Between a birth and a click every step of the linear block is injective on the rows' weights, multiplicities and phases for the event history the record keeps (Theorem 3); the click is the one deletion.

**A hand-worked update.** A row on the heading  $+x$  has the flight accumulator's rate  $2S_1N_l = 128$  ( $S_1 = 1$ ,  $N_l = 64$ ) against the wall  $2T_D$  with  $T_D = \lfloor \sqrt{3 \cdot 64^2} \rfloor = 110$ , so  $2T_D = 220$ , started at  $T_D = 110$ . Interval by interval the accumulator reads 238 (one wall counted: a Link, the remainder 18), 146 (none), 274 (a Link, 54), 182 (none), 310 (a Link, 90), 218 (none), 346 (a Link, 126), 254 (a Link, 34): Links at the intervals 1, 3, 5, 7, 8,  $\dots$ , five Links in eight intervals, and in the long run  $128/220 = 0.5818$  Links per interval, the heading's pace of Section 3. The same three lines of Eq. (1) run the phase (the rate the family's turn, the wall  $N_\phi$ ), a body's owed count and every other component; only the rate and the wall differ.

**In the code.** The engine realises Eq. (1) as one accumulator per component with its rate and wall read from the world file, the carry as the event, in six steps per interval: departures become arrivals (the walk), the readings at each Node, the collision, the measured events' tables and the detectors, the self-creations, and the border's lifetime with the merge of identical rows [2, 17]. The same engine runs every world of this paper from its world file.

**When the transition was made.** The road from ordinary physics to this algebra was walked in twenty-four dated steps between 2026-09-17 and 2026-09-22, each a thing of physics becoming an object of the law, recorded with the reading that showed it [22]; Appendix C lists them.

**The ledger.** Table 1 is the paper's spine: for every result, the rules of Eq. (1) it starts from, the assumptions added beside them with their kind, the freedom left, and the ground of the derivation [18, 2026-09-22]: exact on the GameBoard (no limit, no average), a limit of the grain taken on paper with a rate, or a limit under a spatial condition, an average that covers the shell (over the shell's Nodes, over the directions of a fan declared in a ball, or over the sweep of an orbit), the condition then part of the claim. A result whose derivation is not closed, or whose only reading

is of the GameBoard’s state, is not in the table; the hypotheses beside the law are on the tree (Section 9).

Table 1: The forcing ledger: what the algebra shows, its ground word the rung; what a run measures is Table 2’s. Ground: exact, on the GameBoard, no limit and no average; limit, of the grain, taken on paper with its rate; fan, a limit under an average that covers the shell, the condition part of the claim. A detector reading never lifts a claim from fan to exact: what a run establishes is a lattice-exact claim about the world at hand, whose agreement with the continuum’s form within the named ripple is evidence for the fan claim, not its proof. Every row is derived from the update rules implemented in the simulator; the third column names what is added beside them with its kind; a result resting on a hypothesis beside the law has no row. The section is this paper’s; the derivation’s section is [19]’s.

| Result                                                                                                              | Rules it starts from                                        | Assumptions added (kind);<br>freedom left                                                                                                                                                                                                                                                                                          | Ground                                                                                | Section     |
|---------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|-------------|
| The books’ conservation;<br>the continuity equation                                                                 | the carry (a bijection), the<br>split and rotation matrices | none                                                                                                                                                                                                                                                                                                                               | exact (the<br>lattice identity);<br>limit (the<br>differential<br>form)<br>exact      | 4; 25.4     |
| The group of the six<br>Ports, 48 and 24                                                                            | the six Ports, the hand bit                                 | none                                                                                                                                                                                                                                                                                                                               | exact                                                                                 | 3           |
| The pace of a direction<br>$N_l D /T_D$ , isotropic<br>within $1/T_D$ ; the<br>Manhattan bound<br>$S_1N_l \leq T_D$ | one Link per interval, the<br>digital line                  | none; the wall $T_D$ a rule<br>chosen among few (P9); $N_l$<br>free                                                                                                                                                                                                                                                                | exact                                                                                 | 3; 13.2 (a) |
| The value $c = 1/\sqrt{3}$<br>attained                                                                              | the same                                                    | none                                                                                                                                                                                                                                                                                                                               | limit (the grain<br>$N_l$ )<br>exact                                                  | 3; 1.4      |
| No dispersion, the<br>constancy of $c$                                                                              | the flight table indexed by<br>direction and age            | the rate in transit constant,<br>the flight blind to the phase<br>and to the crowd (P9, a<br>rule chosen among few; the<br>law before the generic<br>entry), not forced by the six<br>operations: a dispersive<br>flight passes every<br>requirement (the<br>derivation’s 27); (A1) of the<br>frame asks the same of the<br>clicks | exact                                                                                 | 3           |
| Gauss’s law of a free<br>family’s flux                                                                              | the walk, the books                                         | a closed surface no periodic<br>axis crosses, no absorber<br>inside (a declared world<br>condition)                                                                                                                                                                                                                                | exact                                                                                 | 4; 3.1      |
| The equivalence<br>principle; the third law<br>at rest (the two readers’<br>pushes)                                 | the step rule (divides by the<br>content); the books        | none                                                                                                                                                                                                                                                                                                                               | exact; the third<br>law to the<br>apportioning’s<br>grain (a<br>GameBoard<br>reading) | 4; 3.3; 9.1 |
| Newton’s inverse square;<br>$G = K\eta F/(4\pi N_w)$                                                                | the bilinear push on the<br>arriving rows’ first moment     | two additivities (24.2); an<br>average that covers the<br>shell; the fan’s Manhattan<br>factor $F = \langle S_1/ \mathbf{D}  \rangle$ (1.287<br>on the registered fan); $N_w$<br>an input                                                                                                                                          | fan                                                                                   | 4; 3.2, 3.3 |
| Coulomb’s law with<br>$k_C = G$                                                                                     | the charge column of the<br>push                            | $\rho$ an input; the same<br>average                                                                                                                                                                                                                                                                                               | fan                                                                                   | 4; 3.4      |
| The retarded potential<br>and the retarded wave<br>equation; Poisson’s<br>equation                                  | the release, the flight, the<br>age moment                  | the same average, the dwell<br>direction-blind, the open<br>box                                                                                                                                                                                                                                                                    | fan                                                                                   | 5; 5.1, 5.5 |

| Result                                                                                                                                                 | Rules                                                                  | Assumptions; freedom                                                                                                      | Ground                                                                                      | Section                               |
|--------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|---------------------------------------|
| The Liénard–Wiechert potential of a moving source, first order                                                                                         | the same                                                               | the same, the fan’s grain                                                                                                 | fan                                                                                         | 12.1                                  |
| The clock’s rate<br>$1/(1 + a_\tau)$ , never 0 at a finite count (no horizon)                                                                          | the owed count                                                         | the age moment as the clock’s word (P9)                                                                                   | exact                                                                                       | 5; 5.2                                |
| The clock’s $1/r$ form; its calibration                                                                                                                | the same                                                               | the $1/r$ form the same average; the calibration $a_\tau = GM/rc^2$ an input of the dictionary                            | fan (the form); input (the calibration)                                                     | 5; 5.2                                |
| The click’s weight a positive quadratic form of power 2                                                                                                | the evaluation at the roots of unity, the phase rotation               | the balanced splitter’s conservation (an axiom of the apparatus); the harmonic constants free; the least-rank click (P10) | exact                                                                                       | 6; 6.5, 6.6                           |
| Born’s bound: a cell within $1/N_\phi$ , the cumulative within $1/(2N_\phi)$ ; Born’s rule as its limit                                                | the ladder’s rungs                                                     | the uniform birth phase (a fact of the source)                                                                            | exact (the bound); limit (the rule)                                                         | 6; 6.2                                |
| The entropy identity; the entropy produced only at the click, the cancel and the leaks, a count of torus points; Boltzmann’s $S = k \log W$            | the click’s one deletion                                               | the coarse graining (a definition); no second law claimed                                                                 | exact                                                                                       | 14, 16.1                              |
| The split an isometry; the interval injective on the amplitudes for a fixed event history                                                              | the six operations                                                     | none                                                                                                                      | exact                                                                                       | 6                                     |
| Planck’s $E = hf$ ; de Broglie’s $\lambda = h/p$                                                                                                       | the release’s cost rule, the turn rule                                 | the dictionary<br>$h = h_q N_\phi = h_A$ , an input                                                                       | exact (identities under the dictionary)                                                     | 6; 6.4; 21.2 rows 51, 52; 24.1 row 25 |
| Young’s spacing<br>The uncertainty relation’s bounds on $\mathbb{Z}_{N_\phi}$                                                                          | two paths’ phase difference<br>the evaluation                          | the fan’s grain<br>the circle as position, its transform as momentum: a hypothesis                                        | limit<br>exact (the bounds)                                                                 | 7.1<br>6; 22                          |
| Exact count marginals; no-signalling of the counts only (the order of the outcomes signals)                                                            | the one gather, the nearest rung                                       | equal weights; open beyond                                                                                                | exact                                                                                       | 7                                     |
| $S(N_\phi)$ exact, $181/64$ on the plateau; $2\sqrt{2}$ the limit                                                                                      | the rungs, the tables                                                  | the tables at $N_t = 256$ (P7)                                                                                            | exact (the rational); limit (the bound)                                                     | 7; 24.4                               |
| The Doppler on the axis by the crossing rule                                                                                                           | the crossing count, its boundary row                                   | none                                                                                                                      | exact (the count per Link); limit (the ratio $1 \pm v/c$ )                                  | 2.2, 2.7                              |
| A body’s dispersion, its momentum’s fraction per axis where the axes’ fires do not coincide, the sum of the paces at most 1; the moving clock’s rate 1 | the drive per axis, the first axis’s step, the owed count              | none: the law’s own predictions                                                                                           | exact                                                                                       | 4.3, 4.4                              |
| Newton’s cooling within one birth’s cost                                                                                                               | the release                                                            | none                                                                                                                      | exact (a bound)                                                                             | 25.9                                  |
| The bending’s coefficient $2(1 + \gamma)$ on the one constant, the $M/b$ form                                                                          | the flow counted per Euclidean Link (flow-link-v1), the ring of starts | a hypothesis beside the law; $c_f$ and $\gamma$ inputs; the two stated factors                                            | conjectured from the algebra; the ring read inside its pins, 4 not closing within the grain | 5                                     |

| Result                                                                                | Rules                                                                                              | Assumptions; freedom                                                       | Ground                                                                                                                                              | Section |
|---------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| The atom's ladder<br>$a_j/a_i = (j/i)^2$ ,<br>$E_j \propto 1/j^2$ , Balmer's<br>27/20 | the closure congruence of the<br>action rows, the balance of<br>the shell mean, the virial<br>form | the shell mean's limit;<br>atom-level-v1 (a<br>hypothesis) for the release | conjectured<br>from the<br>algebra;<br>computed from<br>the algebra, not<br>read on the<br>board; a lattice<br>able to hold the<br>scale is not run | 9       |

### 3 Geometry and symmetries: the directions and $c = 1/\sqrt{3}$

**The six Ports, the group and the front.** Everything geometric in the law follows from one Node and its six Ports,  $\pm x$ ,  $\pm y$ ,  $\pm z$ , and from one Link per interval. The maps of the six Ports onto themselves that keep opposite Ports opposite are the signed permutations of the three axes:  $3!$  orderings of the axes times  $2^3$  choices of sign, 48 in all, the symmetry group of the cube and of the octahedron (Theorem 1). The determinant of a signed permutation is  $\pm 1$ , so it splits the 48 into two halves of 24: the rotations, of determinant  $+1$ , which are the permutations of the cube's four body diagonals; and the improper maps, of determinant  $-1$ , the reflections with the central inversion among them. The hand of a row is that determinant made a column of the state: one pseudoscalar bit that a map  $g$  sends to  $\det(g)$  times itself, so a world with a hand reads which half a map belongs to. The same six Ports bound the pace (Proposition 1 below); that the rows' pace  $c$  sits at that bound and not below is the flight's declared wall  $T_D$ , a rule chosen among few (P9), and given the wall the value is the flight operator's norm, attained exactly on the body diagonals (the derivation's 24.1, row 9).

**Theorem 1** (the 24 of the name). *The maps of the six Ports onto themselves that send opposite Ports to opposite Ports form the group of the signed permutations of the three axes, of order  $2^3 \cdot 3! = 48$ . The determinant is a homomorphism of this group onto  $\{+1, -1\}$ ; its kernel, the rotations of the cube, has index 2 and order 24, and is the group of the permutations of the cube's four body diagonals; the other coset, the reflections, has 24 elements too. The hand, the one pseudoscalar column of a row, which a map  $g$  sends to  $\det(g)$  times itself, tells the 24 rotations from the 24 reflections, as does an axial vector.*

The proof is Appendix A.

**The symmetries.** The law is invariant under the translations of the box and of the interval, and under the 48 signed permutations of the axes, the group of the octahedron of the six Ports and of the cube ( $3! \times 2^3$ ; the 24 of determinant  $+1$  its rotations, the 24 of  $-1$  its reflections, told apart by the hand, a pseudoscalar of the group), up to the law's two declared ties, the digital line's axis order and the collision's Port order ([2], note 39 and section 4); under the circle  $\mathbb{Z}_{N_\phi}$  exactly on the turns between events and at the click only up to the tables' rounding; and under the permutations of the families, each with its columns of the family table carried along [10]. A boost, which mixes a step in space with a step in time, is not among the 48: Lorentz's symmetry is not a symmetry of the GameBoard but of its clicks Outside, under the amplitude split (A2) that the law as built lacks, to second order in the velocity, the lattice's corrections from the fourth (Section 9). The 48 are the GameBoard's stand-in for the rotation group; angular momentum is conserved only up to the GameBoard's grain, and no Noether theorem for the map is claimed [6]. The name of the program counts them.




Figure 3: The Nodes one interval from a Node: the six neighbours are the vertices of the octahedron  $|x| + |y| + |z| \leq 1$ , the causal front of one interval; its inscribed sphere, of radius  $1/\sqrt{3}$ , touches the eight faces on the cube diagonals and is the rows' pace  $c$  (Proposition 1); the faint cube shares the 48 signed permutations of the axes. Drawn by `octahedron.py` from the definitions; no run.

**The pace of the rows, and the octahedron.** Two rules are the whole kinematics of a row: at most one Link per interval (the locality of the GameBoard) and the digital line of its direction, so that after one period of  $S_1 = |D_x| + |D_y| + |D_z|$  Links its displacement is  $D$  itself (straightness). The octahedron  $|x| + |y| + |z| \leq 1$  of the six neighbours (Figure 3), scaled by the age, is the causal front, inside which every row stays; a row's mean Euclidean pace lies within  $1/T_D$  of its inscribed sphere of radius  $1/\sqrt{3}$ , which touches the faces on the cube diagonals (Proposition 1), so the vertex's pace, one Link per interval along a heading, is reached by no row's mean pace: the front is a bound and not a signal.

**Proposition 1** (The pace). *For every primitive direction  $D$ ,  $S_1 N_l \leq T_D$ : the Manhattan pace  $S_1 N_l / T_D$  is at most one Link per interval on every digital line, and  $S_1 = \sqrt{3} |D|$  exactly when  $|D_x| = |D_y| = |D_z|$ . A pace  $v$  that is the same in every direction and at most one Link per interval on every line satisfies  $v \leq |D| / S_1$  for every  $D$ , hence  $v \leq 1/\sqrt{3}$ , the radius of the octahedron's inscribed sphere; and the table's Euclidean pace obeys  $1/\sqrt{3} \leq N_l |D| / T_D < (1/\sqrt{3})(1 + 1/T_D)$  for every  $D$ .*

*Proof.*  $S_1^2 = (|D_x| + |D_y| + |D_z|)^2 \leq 3(D_x^2 + D_y^2 + D_z^2)$  is the Cauchy–Schwarz inequality against  $(1, 1, 1)$ , an equality exactly when the three sizes are equal; so  $S_1 N_l \leq \sqrt{3|D|^2 N_l^2}$ , and since  $S_1 N_l$  is an integer,  $S_1 N_l \leq \lfloor \sqrt{3|D|^2 N_l^2} \rfloor = T_D$ . A row at the Euclidean pace  $v$  on the line of  $D$  makes  $S_1$  Links per length  $|D|$ , so  $v S_1 / |D|$  Links per interval on average, and at most one Link per interval bounds this by 1:  $v \leq |D| / S_1$ . The least of  $|D| / S_1$  over directions is  $1/\sqrt{3}$ , on the cube diagonals ( $1/\sqrt{n}$  in  $n$  dimensions).  $T_D \leq \sqrt{3} N_l |D| < T_D + 1$  gives the pace's interval.  $\square$

So  $c = 1/\sqrt{3}$  Links per interval is the largest isotropic pace the two rules allow, and the table sits at it; that it sits at the supremum and not below is the flight's declared wall  $T_D = \lfloor \sqrt{3|D|^2 N_l^2} \rfloor$ , the design's third statement, a rule chosen among few (P9; the derivation's 24.1, rows 7 and 9), not a consequence of the two; given the wall the value is the flight operator's norm, so the value rests on locality, straightness and the declared wall. Stated exactly: the two rules give  $c \leq 1/\sqrt{3}$  (one Link per interval allows the pace 1 on a heading and only  $1/\sqrt{3}$  on a cube diagonal, three Links for the displacement  $\sqrt{3}$ ), and what is derived is the largest uniform pace the GameBoard allows under them; why the rows realise it, and what a detector in motion reads of it, are not derived

here (Section 9). The anisotropy is the table's rounding, below  $1/T_D$  in every direction and so of order  $1/(\sqrt{3}N_l)$  (Table 2, row 5a, the bound on  $N_l$ ; the bound assumes the rounding's anisotropy is of order  $1/N_l$  at every  $N_l$ , as computed at 64; an upper bound alone gives a sufficient  $N_l$ , not a necessary one); nothing measured enters the value. The number is the bound of the Courant condition for the wave equation,  $c\Delta t/\Delta x \leq 1/\sqrt{n}$  [54], by this same argument, and the lattice Boltzmann sound speed  $c_s^2 = 1/3$  [55]; the paper adds the exact integer table at the bound and its rounding's anisotropy as a bound on the grain, and claims no novelty for the number. Stated with its ground: the pace of a direction,  $N_l|D|/T_D$ , isotropic within  $1/T_D$ , and the Manhattan bound  $S_1N_l \leq T_D$  are exact in integers; the isotropic bound  $1/\sqrt{3}$  is a theorem about a pace the same in every direction, attained on the eight diagonals, above on every other registered direction, and a limit of the grain  $N_l$ .

**What a detector measures.** Two registered runs read the pace after a detector (Appendix C): series Q, a lamp releasing one row on each of the 290 primitive directions with  $|a| + |b| + |c| \leq 6$  toward detectors on the six faces, every click at the derived interval, Node and face, the pace 0.5718 to 0.5893 Links per interval at finite ages against the table's asymptotic 0.5774 to 0.5818; and series L7, the cone, a row of 17 Links on an axis and a row of 24 on the plane diagonal, the same Euclidean distance to one percent, at their counters at the same age.

## 4 Conservation, the flux and the forces

**The books.** Every message released is on exactly one line of the books at every interval, released equals in transit plus absorbed plus escaped plus cancelled, per family, in amount and in content, exact at every interval and at every operation (P3, P5): the carry is a bijection  $s \mapsto (e, s - ed)$  that loses nothing, and the split and the rotation are integer matrices whose conjugate transposes invert them (Theorem 2). From the books the continuity equation of the content follows exactly on the GameBoard (the derivation's 25.4 [19]); it is the first row of the ledger. A paid message carries its label out of its emitter at birth and into its reader at its end, so for paid messages momentum is conserved and the third law holds message by message; a free message, a field's row, takes no recoil at birth and pushes its reader by its label times the reader's content. For a free family the third law is a symmetry of two readers, body A pushed by B's field and B by A's, the two pushes  $-M_A\mathbf{a}_B$  and  $-M_B\mathbf{a}_A$  ( $M$  the content,  $\mathbf{a}$  the acceleration the other's field gives) equal and opposite, and not a balance per message; the momentum lines of the free books are a report of the GameBoard, and the momentum of the bodies and the field together is not booked as conserved (the derivation's 9.1 and 12.4 [19]).

**The operations.** Six operations, of three of Section 2's kinds (the translation, the bilinear form, the Euclidean division), make the whole of gravitation and electricity in the law. The free release: a body of content  $M_B$  holding a free family releases at each self-creation the carry of the count of rate  $M_B\eta$  against the wall 1,  $\eta$  the release rate per unit of content per direction, on every declared direction, so a fan of  $K$  directions releases  $q = KM_B\eta$  per interval. The walk: each unit walks its digital line at the pace of Proposition 1. The reading: the label flow  $\mathbf{a} = \sum \text{amount} \times \mathbf{u}_D$  of the rows arriving at the reader's Node,  $\mathbf{u}_D$  the integer unit vector at the scale  $N_l$  ( $|\mathbf{u}_D| = N_l$  within 1.35 percent on every direction). The coupling, a bilinear form with a declared integer matrix: per axis  $\text{push}_A = M_A(\rho_A\rho_B - 1)\mathbf{a}$ , the gravity column  $-M_A\mathbf{a}$  and the charge column  $+\rho_A\rho_B M_A\mathbf{a}$ ,  $\rho$  the declared charge per unit of content. The momentum:  $\mathbf{p}_A \leftarrow \mathbf{p}_A + \text{push}_A$ . The step, the Euclidean division: one Link per  $d_p/|p_a|$  self-creations on the axis  $a$ , the wall  $d_p = N_l N_w M_A + |p_a|$ ,

$N_w$  the world's width, so that for a body whose clock owes nothing the speed on the axis is  $|p_a|/(N_l N_w M_A + |p_a|)$  per interval.

**Gauss's law, exact.** The walk is a translation: a free unit released inside a closed surface that no periodic axis crosses, with no absorbing reader inside it and within the unit's lifetime, crosses the surface outward once more than inward (a non-convex surface it may leave, re-enter and leave again, the signed crossings summing to one, the crossing counted with the sign of the Port's outward normal; no splitter, absorber or cancelling pair inside the surface; no collision acts on a fan or on lone beams), so once the front has passed, the net amount crossing the surface per interval equals the release inside it for a constant release, exactly, and for a varying release the crossings at the interval  $t$  equal the releases at the retarded intervals, direction by direction. The crossings of an interior surface are the host's count (GAMEBOARD, a diagnostic); a face's border clicks are a detector's (DETECTOR). Proved from the definitions under those conditions; no detector reading of it is registered.

**The inverse square as a shell mean.** On a beam the flow per Node is the same at every Node of the digital line: along a beam the field falls as  $1/r^0$ , and off every beam it is 0, by the walk. The inverse square is the density of beams over a shell. A shell at distance  $r$  of unit radial width holds  $\mathcal{N}(r)$  Nodes, and a beam of direction  $\mathbf{D}$  crosses it at  $S_1/|\mathbf{D}|$  of them (one Link per Node,  $|\mathbf{D}|/S_1$  of Euclidean advance per Link), each reading the beam's arrivals, so the mean over the shell of what a body reads is

$$\langle a \rangle(r) = \frac{q N_l F}{\mathcal{N}(r)} \rightarrow \frac{q N_l F}{2\pi r} \text{ on the plane,} \quad \frac{q N_l F}{4\pi r^2} \text{ in space,} \quad F = \langle S_1/|\mathbf{D}| \rangle_{\text{fan}}, \quad (2)$$

as  $\mathcal{N}(r) \rightarrow 2\pi r$  and  $4\pi r^2$ , with  $F$  the fan's Manhattan factor, 1 on a heading,  $\sqrt{3}$  on a body diagonal, 1.287 in the mean over the registered fan [33],  $4/\pi$  for a fan uniform in angle on the plane and  $3/2$  for one uniform in solid angle in space, a factor that vanishes neither with  $r$  nor with the fan's grain (the presence, Eq. (5), carries the dwell  $T_D/(S_1 N_l)$  per Link that cancels it into  $\tau_L$ ; flow-link-v1, rows 13 and 14 of Table 2, divides every push by the same mean; the derivation's check of the shell mean on the six headings, series C, has the factor 1 on every heading and does not test it, and no non-heading fan check of the coefficient is registered); the departure at finite  $r$  is the GameBoard's count of Nodes on a shell, Gauss's circle problem,  $\mathcal{N}(r) = 2\pi r + O(r^\theta)$  with the exponent  $\theta \leq 131/208$  [67], so the shell mean's relative ripple vanishes as  $r^{\theta-1}$ . No detector reading of the shell mean is registered;  $\mathcal{N}(r)$  is the GameBoard's count of Nodes on a ring, a number of the derivation and not a reading. Derived from the update rules implemented in the simulator, in the shell mean, with the ripple named; on one Node the inverse square exists only in the limit of every direction, and a fan declared as a cube of directions has in that limit a cubic anisotropy of the declaration,  $3^{3/2}$  between a corner and a face centre, absent for a fan declared within a ball.

**Newton's law and the place of  $G$ .** A body  $A$  of content  $M_A$  at the mean flow  $\langle a \rangle$  takes per interval the push  $-M_A \langle a \rangle$ , and its speed on an axis is  $|p_a|/(N_l N_w M_A + |p_a|)$ ; for  $|p_a| \ll N_l N_w M_A$  the acceleration is push/ $(N_l N_w M_A)$  Links per interval squared, so

$$a = -\frac{G M_B}{r^2}, \quad G = \frac{K \eta F}{4\pi N_w} \text{ in space;} \quad a = -\frac{G' M_B}{r}, \quad G' = \frac{K \eta F}{2\pi N_w} \text{ on the plane,} \quad (3)$$

in Links cubed (squared) per unit of content per interval squared:  $G$  is the source's release rate per unit of content per direction,  $\eta$ , times the number of directions  $K$ , over  $4\pi$  and the world's width

$N_w$ . No detector reading of either form is registered; the decisive reading, a closed orbit's period at two radii, is NOT MADE. Nothing of  $A$  enters: the equivalence principle is exact (the step rule divides by  $M_A$ ). The third law at rest, the two bodies' pushes equal and opposite, holds to the grain of the apportioning: in series C, item 2, the two sources' momenta agree to six parts in  $10^6$ , a GameBoard reading, a diagnostic (the derivation's 9.1 [19]); it is a symmetry of the two readers and not a balance per message, the free release taking no recoil. Two departures from Newton's law as written, both stated: the field is retarded at  $c$  (the front at  $r$  arrives at the interval 1 plus the flight table's first age at  $r$  Links on the heading), so the limit is the retarded inverse square and not action at a distance; and the speed the push builds, per axis where the axes' fires do not coincide and one Link per interval in all, saturates at one Link per interval, above the rows' pace, so a body can outrun its own field's rows on the drive as built. Derived from the update rules in the shell mean,  $G$  named, the width  $N_w$  an input, a choice of units (Table 1).

**Coulomb's law and the one constant.** The charge column adds  $+\rho_A\rho_B M_A \mathbf{a}$  per axis; with  $q_A = \rho_A M_A$ ,  $q_B = \rho_B M_B$  and  $\langle a \rangle$  proportional to  $M_B$ , the electric push in the shell mean is

$$\text{push}_e = q_A q_B \frac{K \eta N_l F}{4\pi r^2}, \quad a_e = \frac{q_A q_B}{M_A} \frac{G}{r^2}, \quad (4)$$

Coulomb's inverse square with the same constant as gravity,  $k_C = G$  in the law's units, repulsive for like signs; the ratio of the two forces on one body is  $-\rho_A\rho_B$  at every  $r$ . Derived under the same average as Newton's; no detector reading of it is registered. Nature's ratio for two protons,  $1.24 \times 10^{36}$ , is  $\rho_p^2$  here,  $\rho_p = 1.1 \times 10^{18}$  units of charge per unit of content: the hierarchy is the declared  $\rho$ , an input, not a formula of the law.

**The ground of these derivations.** Gauss's law of the flux and the equivalence principle are exact on the GameBoard; the third law at rest is the two readers' symmetry, to the apportioning's grain (a GameBoard reading, six parts in  $10^6$ ). The inverse square, Newton's and Coulomb's, is a limit under an average that covers the shell, in one of three forms: the shell mean (the ripple  $O(r^{\theta-1})$ ,  $\theta \leq 131/208$ ), the fan dense at one Node for a fan declared in a ball, or the mean over one closed turn of an orbit, the only one readable after a detector, which equals the ring mean at one radius of the loop,  $\langle r^2 \rangle_\theta / \langle r \rangle_\theta$  on the plane and  $\sqrt{\langle r^2 \rangle_\theta}$  in space ( $\langle \cdot \rangle_\theta$  uniform in angle), exactly in the continuum for every closed turn by the area rule of a radial push ([46], Theorem B; Kepler's second law the comparison, not an input); against the ring mean at the angle-mean radius  $\bar{r}$  the orbit's mean is smaller by  $\bar{r}^2 / \langle r^2 \rangle_\theta$ , between  $1 - e^2$  and 1, equal only on the circle; on the lattice it holds up to four grain terms, the ring's count against  $2\pi r$  (0.90, 0.95 at  $r = 12, 24$ ), its incidence against the fan's L1 factor (1.07, 0.97), the drive's anisotropy (0.069) and one Node's dwell per crossing of a line (1.55 at the registered pace; its mean over the fan's crossings small, assumed), GAMEBOARD by formula; on one Node beside a source the field is a comb of beams, the average part of the claim (the ledger's caption). After a detector the equivalence principle is measured (series D3) and the  $1/r$  force's scale symmetry reads consistent, 1.997 for  $2.00 \pm 0.18$ , on loops that are not similar figures, resting on the  $1/r$  push's scale symmetry and not on the equality above; the turns'  $e = (r_{\max} - r_{\min}) / (r_{\max} + r_{\min})$  is not on record, the run's extremes bound it by 0.96 and 0.70 at  $r = 12, 24$  (floors 0.07, 0.51 on the factor, GAMEBOARD by formula), which decides nothing ([46], section 5); the inverse square's decisive reading is NOT MADE and  $G$ 's value is not read; Coulomb's run is not made; Poisson's interior is read after a detector (series X, Section 5).

## 5 The delay field: the retarded potential, Poisson's equation and the clock

The rows of Table 1 on the delay field are written out here from the derivation's section 5 [19]; the register's series E reads probes on the GameBoard, diagnostics this paper does not carry, and the field's one measurement after a detector is the clock's form at two distances, series T (row 12 of Table 2; read before the generic entry of 2026-09-22 [32], not readable under the law as it stands, being re-read in the weak field).

**What is delayed.** Nothing in flight is delayed under the law of this paper, the runtime before the generic entry (P9): a row moves by the flight table at one pace, blind to the crowd (series K, Table 2). What a crowd delays is a clock. A body reads at its Node the presence of every row of another number (the engine's default) or, on an entry that reads the age, their age moment (the clock's word, P9) and owes after each self-creation the whole part of  $a_r n_c/d_c$  intervals before the next,  $a_r$  what it read and  $[n_c, d_c]$  the clock's pair of Section 2 ( $[1, 1]$  in the scalar world,  $[1, 2]$  in the age world), so that its rate is  $1/(1 + a_r n_c/d_c)$ ; its push reads the flow of the same rows.

**The two fields of one stream.** Let a source release  $q(t)$  rows per interval on a fan of every direction (the limit of Section 4's shell mean), each row moving at  $c$  and carrying its age. The presence  $\Pi$  at the distance  $r = |\mathbf{x}|$  from the source at the time  $t$  is the rows dwelling there, and the age moment  $\mathcal{A}$  is the presence times the age they carry,  $r/c$ :

$$\Pi(\mathbf{x}, t) = \frac{q(t - r/c) \tau_L}{4\pi r^2}, \quad \mathcal{A}(\mathbf{x}, t) = \frac{q(t - r/c) \tau_L}{4\pi c r}, \quad (5)$$

$\tau_L$  the radial residence factor, the intervals a row spends per unit of Euclidean distance,  $T_D/(N_l|D|) = 1/v_D$  (1.72 on a heading,  $\sqrt{3}$  in the limit of every direction); the intervals per lattice Link along a row's staircase are  $T_D/(S_1 N_l)$ , a different factor (1 on a body diagonal), and a finite shell's coefficient depends on its digital geometry, so the coefficient of (5) is the heading's on a beam and the limit's on the sphere. The presence is the retarded flux of a conserved stream; the age moment is the retarded potential of the source, since  $f(t - r/c)/(4\pi r)$  is the retarded Green's function of the wave operator, so  $\mathcal{A}$  obeys the inhomogeneous scalar wave equation with the release as its source,

$$\frac{1}{c^2} \frac{\partial^2 \mathcal{A}}{\partial t^2} - \nabla^2 \mathcal{A} = \frac{\tau_L}{c} q(t) \delta^3(\mathbf{x}), \quad (6)$$

$\delta^3$  the three-dimensional Dirac delta, and in the static limit Poisson's equation  $\nabla^2 \mathcal{A} = -(\tau_L/c) q \delta^3(\mathbf{x})$ , solved by  $\mathcal{A} = (\tau_L q / 4\pi c) / r$ . The moments are sums over rows, so the field of many sources is the sum of their fields: the theory is linear. The flow the push reads is the stream's flux vector,  $\mathbf{a} = (q N_l / 4\pi r^2) \hat{\mathbf{r}}$ ,  $\hat{\mathbf{r}}$  the unit vector from the source and  $N_l$  the label each unit carries, which is  $-(N_l c / \tau_L) \nabla \mathcal{A}$  in the static case; the gravity column reads  $-M_A \mathbf{a}$  (Section 4), so a body's acceleration, the push over  $N_l N_w M_A$  (the drive's conversion of a push to a pace, as in Eq. (4)'s  $a_e$ ), is  $+(c / (N_w \tau_L)) \nabla \mathcal{A}$ , up the age moment's field, toward the source. Derived from the update rules, in the limit of every direction: the age moment is the retarded potential of the release and obeys the wave equation there. After a detector the interior of a shell reads flat within the tolerance,  $k(2)/k(4) = 1.0029$  for the map's pin 1.0039 (series X, DETECTOR), the shell theorem's interior; the finite- $r$  ripple is MEASURED ONLY.

**The clock's redshift.** A clock at the age moment  $a_\tau = C_a M/r$ ,  $C_a$  the constant of Eq. (5) times the clock's pair, runs at  $1/(1 + a_\tau)$ ; general relativity's clock at the potential runs at  $\sqrt{1 - 2GM/rc^2} = 1 - GM/rc^2 - \frac{1}{2}(GM/rc^2)^2 - \dots$ . Under the calibration  $a_\tau = GM/rc^2$ , an input of the dictionary (by the derivation's 5.1 and 3.3 the clock's constant and Newton's  $G$  have the exact ratio of the suspension pair times the width, so the calibration is a constraint on the inputs), the two agree at first order: the redshift between two clocks,  $1 - a_\tau(r_1) + a_\tau(r_2)$ , is the  $1/r$  law. Derived at first order, the calibration an input. At second order the GameBoard reads  $1 - a_\tau + a_\tau^2$  against  $1 - a_\tau - a_\tau^2/2$ , and its rate never reaches zero at a finite  $a_\tau$ : a different law in the strong field, no horizon (the crowd only slows; nothing stops a clock). No detector reading of the strong-field form is registered; the weak-field confrontation ( $a_\tau \ll 1$ ) is a run the register names as the follow-up, neither pinned nor made. Which word the clock counts, the presence or the age moment, was put to the GameBoard as series T, the pins written before the run [4, 13]: a lamp at rest, two sources at 3 and at 6 Links, the detector at  $x = 110$ ; under the presence word the detector reads  $1 + z = 1.3000$  at both distances; under the age word 2.6517 and 4.1500, the ratio of the two  $a_\tau$  being 1.907 for the lines' pin 1.909, nature's 2.00 being 4.6 percent away; every pin met, measured and replicated [9]; read under the law before the generic entry of 2026-09-22; not readable under the law as it stands; the four worlds are being redeclared in the weak field (the pair [1, 16384], the crowd's age moment at most 1024 per Node on the light's path) and re-read, the new numbers to replace these; the gravitational redshift's ratio is a reading of the law before the entry, not of the law as it stands. What the detector establishes is the form at two Nodes, a lattice-exact reading; the continuum's  $1/r$  is its limit under the average that covers the shell, and the reading is evidence for it, not its proof. The age moment is the clock's word, an assumption of the law (P9; [6, record 394]), not a derivation; the law as built counts the presence by default and the age moment on an entry that reads the age; the presence word is the alternative refuted on the form.

**Light: no optical metric.** A row reads nothing of the crowd: beside a mass it is neither bent nor delayed (series K, three worlds, the deflection 0.000 pixel and the delay 0.00 interval, DETECTOR; FAIL against nature's 1.75 arcseconds, Table 2); the law as built here is the law before the generic entry of 2026-09-22 (on main at 5b855791 since that day [32]), and the paper's readings under the entry are the ring's alone (row 13); the delay field is a metric for clocks and for the bodies' pushes, not for the flight (Section 9, the bending under the key). What the algebra gives, on the law's integers and before any run (COMPUTATION): under the key `optical` the walk of a light row past a held mass by the law's steps, read at a ring of Nodes at the impact distance  $b$ , recovers the  $M/b$  form,  $C_{\text{ring}} = abc^2/(GM) = 5.12$  at  $b = 6$  (5.44, 5.86 at  $b = 3, 8$ ), flat in  $b$  within 15 percent and not 4 within the ring's grain, the factor left the fan's  $F_{L1} = 1.421$  of the flow ( $3/2$  in the isotropic limit) [34]; with the flow counted per Euclidean Link of its line (flow-link-v1, the one constant: the push's constant equal to the clock's, 0.990 against 0.993) the ring reads  $C_{\text{ring}} = 3.65, 3.82, 3.92$  at  $b = 6, 3, 8$  against 4, that is  $2c_f$  within the finite path's  $L/\sqrt{L^2 + b^2}$  (0.974 at  $b = 6$ ) and the shells' 0.990, one to three grains;  $4 = 2(1 + \gamma)$  with  $\gamma$  an input read back, not derived [35]. Stated as a conjecture, labelled so and not read: on the one constant the law's deflection coefficient is  $2(1 + \gamma)$  with  $\gamma$  the declared input, the  $M/b$  form with Einstein's factor 2 at  $\gamma = 1$ , conjectured from the algebra (the step algebra's simulation of the ring, 3.65 to 3.92 against 4, closing to  $2c_f$  within the two stated factors to one to three grains, COMPUTATION), not recovered and not read: what is read is series K's 0.000 pixel under the law as built (FAIL), the  $\gamma = 0$  ring inside its four pins, and, on the split ladder, the deciding  $\gamma = 1$  ring read at 0.731 against  $0.731 \pm 0.025$  with every arrival Node the map's (DETECTOR), the constant through the stated factors outside the one-grain pin

at the three  $\gamma = 1$  rings by 2.40, 1.07, 1.40 grains (COMPUTATION): the ring's rule read, the bending's 4 not closing within the grain. The run after the algebra is Table 2, row 13.

## 6 Measurement and the quadratic form

The worlds of this section have no body: the law of Section 2 on rows alone, and the measurement is the one threshold that is read out. Every interval applies three of the six operations to the state, the translation (the flight, the phase, every count and the birth), the multiplication by a declared integer matrix (the split and the rotation) and the sum in the group ring (the merge, where opposite phases cancel); between clicks the rates are constant, so the map has a closed form and every count of Section 8 was computed from it before its run. The click is the one place with a fourth operation: the apparatus sums a record's rows through the tables and squares the sum, the norm, and only then applies its threshold to the birth phase  $u$ .

**Definition 1** (Row and record). A *row* is  $(x, D, \tau, p, e, h, r, \ell, w, \mathfrak{m})$ : Node, direction, age, phase  $p \in \mathbb{Z}_{N_\phi}$ , emitter, content per unit, record  $r$ , branch label  $\ell$ , amount  $w \geq 1$ , multiplicity  $\mathfrak{m} \geq 1$ . Its amplitude is

$$z = \frac{w}{\sqrt{\mathfrak{m}}} \frac{C_{N_\phi}[p] + iS_{N_\phi}[p]}{256}, \quad |z|^2 = \frac{w^2}{\mathfrak{m}} \cdot \frac{C_{N_\phi}[p]^2 + S_{N_\phi}[p]^2}{65536}. \quad (7)$$

Two rows equal in every field but  $w$  are one row with the amounts added; two rows of one record and label equal in every field but a phase difference of  $N_\phi/2$  are one row with the amounts subtracted (an equal pair is no row). The state of record  $r$  is an element  $\psi_r$  of the free  $\mathbb{Z}$ -module  $M_r$  on the basis  $(x, D, \tau, p \bmod N_\phi/2, e, h, \ell, \mathfrak{m})$  with  $|p + N_\phi/2\rangle = -|p\rangle$ ; the scaling  $(w, \mathfrak{m}) \rightarrow (kw, k^2\mathfrak{m})$  leaves the amplitude and the norm unchanged and is an identification at the read-out, not a relation of the module (the quotient by it would not be free,  $b_1 = 2b_4 = 4b_{16} = \dots$  for the rows  $b_{\mathfrak{m}}$  of amount 1); the GameBoard's state is one normal form per element of  $\bigoplus_r M_r$ , and the multiplicity, which every split multiplies by  $A$ , is bounded by the law's bound of an amount, a multiplicity and a momentum component,  $2^{62} - 1$  (`events/world.py`, one bit spared for one more sum), inside the host's working bound  $2^{63} - 1$  for every intermediate (`core/integer.py`); a run that would pass either is refused (P1's bound, named here). In the row form of Theorem 2 a rotation multiplies  $\mathfrak{m}$  by 65536, so four in succession would pass the law's bound; the engine does not carry that factor, keeping a record's label set and applying the tables once at the click's form, the cell's weight reduced by its greatest common divisor, so the depth of successive rotations is not limited by the bound; where the bound binds, it is a limit of the implementation's domain and never a prediction. The norm is  $\langle \psi, \psi \rangle = \sum_{\text{rows}} w^2/\mathfrak{m}$ . The archived engine narrows the quotient at a detector: rows of one record with different  $\mathfrak{m}$  at one set are added when the ratio of the multiplicities is the square of a rational (the pointers rescaled by the integer roots) and refused otherwise; at the tree of the runs two multiplicities at one set were refused, and no reported world has them, so every set below reads one  $\mathfrak{m}$ .

The host, not any Node, keeps per live record the birth phase  $u \in \mathbb{Z}_{N_\phi}$ , the count of live units (amounts), an accumulated pointer  $(X, Y)$  per (Node, label) and a gathered flag. This is the apparatus's layer; the GameBoard never reads it. It is a memory no Node could own: on the two-slit world the two paths of one record reach a pixel up to 18 intervals apart, so no Node ever holds both rows [6].

**Definition 2** (The update rules of one interval). Applied in this order to every row.

- F Flight:  $(x, D, \tau, p) \mapsto (x + \text{pos}_d(\tau + 1) - \text{pos}_D(\tau), d, \tau + 1, p + \varphi(\tau))$ , where the family's phase turn is either an integer per Link, added only when the step is a Link, or a pair  $[n, d]$  per interval of age,  $\varphi(\tau) = \lfloor (\tau + 1)n/d \rfloor - \lfloor \tau n/d \rfloor$ , counted from the row's birth or last re-emission.
- C Collision: at a Node holding several rows of amount 1 the table permutes their directions as a function of the whole set present;  $r, \ell, \mathbf{m}$  unchanged; a bijection of states, not a linear map. No collision acts in any world of this paper.
- S Split, at a measured event with outputs  $\{(d_i, a_i, t_i)\}$ ,  $a_i \in \mathbb{Z}$ : the row  $(w, \mathbf{m}, p)$  is absorbed and re-emitted as the rows  $(wa_i, \mathbf{m}A, p + t_i)$  on  $d_i$  at age 0,  $A = \sum_i a_i^2$ . The balanced splitter is  $(1, 1)$ ,  $A = 2$ ; the reflection's turn is  $N_\phi/4$ .
- R Rotation, at a set with setting  $s$  and turn  $t$ , on a label pair  $(0, 1)$  read into the channels  $(+, -)$ :  $U_{s,t} = \begin{pmatrix} C'[s] & S'[s]v(t) \\ -S'[s] & C'[s]v(t) \end{pmatrix}$  with  $v(t)$  the turn by  $t$  steps on the label-1 column; a row of label 0 becomes  $(wC'[s], 65536\mathbf{m}, p)$  on  $+$  and  $(wS'[s], 65536\mathbf{m}, p + N_\phi/2)$  on  $-$ , a row of label 1 becomes  $(wS'[s], 65536\mathbf{m}, p + t)$  on  $+$  and  $(wC'[s], 65536\mathbf{m}, p + t)$  on  $-$  (the row form exact for  $t$  a multiple of  $N_\phi/4$ , the only turns used here; the archived code multiplies the column by the tables' phasor at  $t$ ). With  $t = 0$  this is  $U_s$  of Theorem 2; the GHZ bases below are  $X = (s, 0)$  and  $Y = (s, N_\phi/4)$ .
- M Normal form: merge and cancel as in Definition 1.
- B Birth: a lamp's self-creation at interval  $t$  releases record  $r$  with  $u$  the lamp's phase at the birth,  $(b-1) \bmod N_\phi$  for a lamp started at 0 that turns one step per birth,  $b$  its count of births (every world here;  $b-1 = t-1$  only while the lamp births at every interval; the archived code writes  $(b-1) \bmod N_\phi$  directly), rows of amount 1 on its  $k$  directions with  $\mathbf{m} = k$ , each row's phase starting at  $u$  (plus the lamp's own turn), and, for a pair, the declared labels with integer weights on separate arms.

**Definition 3** (The click). At a set that reads the sum, the rows of record  $r$  and label  $\ell$  that arrive are absorbed and the layer accumulates

$$X += \sum 32 w C_{N_\phi}[p], \quad Y += \sum 32 w S_{N_\phi}[p], \quad (8)$$

the phase  $p$  as the row carries it on arrival (the turns of rule F already in it). The pointer is kept per Node: interference is coherent within one Node only, and the offer of a set of Nodes (a detector, a face) and label  $\ell$  is the sum over its Nodes of  $(X_x^2 + Y_x^2)/\mathbf{m}$ , one cell of the ladder, the click landing at the Node within the set that  $u$  selects by the same rungs; all rows at a set share  $\mathbf{m}$ . When the record's live count reaches 0, with  $C_K = \sum R$  over its offers and the cumulative sums  $C_1 \leq \dots \leq C_K$  (the total) in the declared order,

$$\boxed{b_k = \left\lfloor \frac{2N_\phi C_k + C_K}{2C_K} \right\rfloor \quad (b_0 = 0, b_K = N_\phi), \quad \text{the click is the } k \text{ with } b_{k-1} \leq u < b_k,} \quad (9)$$

and the world's list gains one row: the record,  $u$ , the chosen set and label,  $R, C_K$ , the count of combinations before and after. For a record with arms at two rotated sets  $A$  and  $B$  (settings  $a, b$ ) the cells are the joint outcomes  $(o_A, o_B)$  in the arms' declared order, with the weights

$$R(o_A, o_B) = J(o_A, o_B)^2, \quad J(o_A, o_B) = \sum_\ell c_\ell U_a[o_A][\ell] U_b[o_B][\ell], \quad (10)$$

$c_\ell$  the labels' declared weights,  $(1, 1)$  in every world here; and the record is gathered once, when the rows at both arms have ended. The click of an arm has two parts. The arrival: an event at the arm's own detector at its own interval, the rows absorbed and their content placed there, the flight blind to the other arm's setting and length (P9), so the arrival interval at either arm is that arm's flight alone. The outcome: the cell of  $u$ ; for the party declared first (the arms' declared order, not the order of arrival) the coarse rung at  $N_\phi/2$ , a function of  $u$  alone,  $u$  the record's wheel reading fixed at its birth ( $u = \text{ordinal} \times r \bmod W$ ), so determined before either arrival and known at the first party's arrival from the record's own wheel under the hypotheses of Theorem 5, computed by nothing at that arm and written only at the completion; for the second the fine rung within it, fixed at the completion from the joint weights. The world's row carries both outcomes at the completion interval, so the near party's outcome is written at the far party's tick, a bookkeeping of the host and not a delay of any arrival. Reading the ladder in that order is the same as a coarse rung over  $o_A$  followed by a fine rung over  $o_B$  within it with the same  $u$ , which is how Theorem 5 states it. A set that reads without absorbing (the which-path read of the polariser) stores the record's residual selector keyed by its labels and lets the rows go on; it is a deferred offer, gathered at the record's one click on the labels present then, not an outcome at the interval of the read. A read, then a rotation of the labels, then a second read, taken as two outcomes in time, is not modelled: the first read's selector is applied once, at the gather, to the labels as the rotation left them [17].

**Lemma 1** (One click per record). *A record every row of which ends (at a set, a face or the border; not a row parked at rest or on a periodic axis without a lifetime) completes and is gathered exactly once: the live count reaches 0;  $b_K = N_\phi$  (with  $C_K > 0$ , the born row's norm up to the tables' rounding, Theorem 2; at  $C_K = 0$  the engine gathers nowhere, [6, record 363]) and  $b_0 = 0$ , so every  $u \in \mathbb{Z}_{N_\phi}$  lies in exactly one cell. Every record of the worlds below is of this kind; the delay between a record's last arrival and its completion is the record's, not the GameBoard's.*

Three things are put in here and not derived, and the paper claims none of them. Definition 3 recovers the Born rule's form: the weight is the squared length of the summed pointer, and the ruler over  $u$  is inverse-transform sampling of a uniform variable, so over births with  $u$  uniform on  $\mathbb{Z}_{N_\phi}$  the count in channel  $k$  is  $b_k - b_{k-1}$ , the difference of two cumulative rungs, Born to the resolution  $1/N_\phi$  per cell. No principle below the square is offered. The uniformity of  $u$  is an assumption about the source:  $u = (b - 1) \bmod N_\phi$  is uniform over  $N_\phi$  consecutive births, (one birth per  $u$  in every run below), and its independence of the settings is a fact of a world, not of the model:  $u$  is a clock and a chooser is a clock, so they are independent when the chooser's period is coprime to  $N_\phi$  and the run covers their common period, as in this paper's choosers' world (periods 3 and 5 against  $N_\phi = 64$ ). The tables at  $1/256$  are the third of these three, and a declared input of the law (Section 2, P7). What the paper derives is what follows from these with the GameBoard's rules: the interference of one record, the exact marginals, the  $\cos^2$  correlation of the pair, GHZ's zeros, and the finite- $N_\phi$  value of  $S$ . The click is the only step that removes combinations and adds a row to the world's list; the birth is the only step that adds combinations.

**The two theorems of the interval.** Two theorems of the linear block are stated here and proved in Appendix A.

**Theorem 2** (The split is an isometry whose conjugate transpose inverts it). *The split  $(w, \mathbf{m}, p) \mapsto ((w a_i, \mathbf{m} A, p + t_i))_i$  preserves  $\sum w^2 / \mathbf{m}$  for every nonzero integer vector  $(a_i)$ , and the conjugate transpose of the table (the same weights with the turns reversed) applied to its outputs, followed by the normal form, returns the input up to the scaling  $(k w, k^2 \mathbf{m})$  with  $k = A$ . The rotation satisfies  $U_s^\top U_s = n_s I$  exactly with  $n_s = C'[s]^2 + S'[s]^2$ , and multiplies a record's norm by  $n_s/65536$ : an isometry only up to the tables' rounding (65705/65536 at  $N_\phi = 64$ ,  $s = 1$ ).*

**Theorem 3** (The interval is injective between clicks). *Let  $Q$  be the quotient of  $\bigoplus_r M_r$  that forgets the age (rows differing only in their age identified), and fix the event history of the record, the intervals and Nodes of its births, splits, rotations and re-emissions, which the record keeps. For an interval without a click, an end at a face or the border, or a collision, and for splitters whose joint table  $B$  (one column per input, the entries  $a_i x^{t_i}$  in the group ring) satisfies  $B^H B = AI$ , the balanced splitter with the turn  $N_\phi/4$  among them,  $M \circ R \circ S \circ F$  induces an injective  $\mathbb{Z}$ -linear map of  $Q$  into itself: no rule of the interval deletes a weight, a multiplicity or a phase, and the state at any interval between two clicks, with the history, determines them at the birth. The age is not a coordinate of  $Q$  but a function of the history, the current interval less the interval of the row's last event; a split restarts it at 0 by the definition of age (Definition 2) and forgets nothing, since the interval of the split is in the history; the click is the one deletion. A collision, where it acts, is a bijection of the GameBoard's states that is not linear; none acts in the worlds of this paper. The theorem is of the rows' linear block; a body's drive, whose later coincident fire is lost under the engine's rule (P5, Section 2), is outside it and is not injective in that case.*

So a record's norm is conserved exactly by  $S$  and by the flight, and by  $R$  and by every read through the tables only within the tables' rounding (Appendix B). The normalisation by  $C_K$  in (9) makes the click's probabilities sum to one at completion; it does not restore the norm along the way, and the paper claims exact unitarity nowhere.

**The click's square as a positive quadratic form: a quadratic read-out characterization.**

The row of Table 1 on the click's weight is written out here from the derivation's section 6.5 [19]; the register's checks are series L's clicks (Section 8). The click's square, the norm of a record's summed pointer, is the one operation of the law beyond the translation, the wall, the declared matrix and the addition, and the place where the Born rule enters. Asked: whether the operations of Section 2, with one hypothesis on the reading, force the reading at a click to be that square, or what weaker statement holds and what the register pins.

**The objects.** A record's rows at one Node and label are one element  $f = \sum_p f_p x^p$  of the group ring  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ ,  $x$  the generator of one phase step and  $f_p$  the amount at the phase  $p$ ; the pointer is  $\text{ev}(f) = \sum_p f_p \zeta^p$ ,  $\zeta$  the primitive  $N_\phi$ -th root of unity, at the tables' scale. The merge's cancel identifies  $x^{N_\phi/2}$  with  $-1$ , so the rows after the cancel are an element of  $\mathbb{Z}[x]/(x^{N_\phi/2} + 1)$ ; for  $N_\phi$  a power of two (every registered  $N_\phi$ ) that quotient is the ring of cyclotomic integers  $\mathbb{Z}[\zeta_{N_\phi}]$  itself, a free  $\mathbb{Z}$ -module of rank  $N_\phi/2$ , and  $\text{ev}$  is the identity on it: the GameBoard's cancel and the ideal click's zero, the evaluation on  $\mathbb{Z}[\zeta_{N_\phi}]$ , are one relation, exactly (for an  $N_\phi$  with an odd factor they are not; the register has no such  $N_\phi$ ). The table click, Definition 3 with the declared integer tables, is a different map: the built click keeps the relation one way: every cancel reads  $(0, 0)$ , and the tables read  $(0, 0)$  on more, the table pointer being a  $\mathbb{Z}$ -linear map of  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$  onto a lattice of rank 2, its kernel of rank  $N_\phi - 2$  against the cancel's  $N_\phi/2$ ; at  $N_\phi = 8192$  only 1952 of the 8192 table pairs are distinct and two rows of amount 1 at the phases 0 and 4097 read  $(0, 0)$  where  $|\text{ev}(f)|^2 = 4 \sin^2(\pi/8192) = 5.9 \times 10^{-7}$ ; at  $N_\phi = 64$  no zero exists with at most three rows of amounts up to 4 ([31], a computation from the tables, not a run), a fact of the tables and not a relation of the module. The reading is a function  $R : \mathbb{Z}[\zeta_{N_\phi}] \rightarrow \mathbb{R}$  of a cell's element, up to one factor per record, which the multiplicity  $\mathfrak{m}$  books.

**The hypotheses.** (a) The phase rotation,  $R(xf) = R(f)$ : a rotation of the record's phase by one step changes no weight (the rungs take no  $u$ ;  $u$  only selects among them), an operation of Section 2. (b) The balanced splitter conserves the reading for every two inputs,  $R(x^{N_\phi/4}f + g) + R(f + x^{N_\phi/4}g) =$

$A_2(R(f) + R(g))$  for all  $f, g$  (the (1, 1) split with the quarter turn on the reflected row,  $A_2$  its multiplicity factor): a hypothesis on the reading, stronger than Section 2's split, which is an integer matrix on rows and says nothing of what a detector reads under it; the register checks it only on the registered pairs (the totals 64/0 and 32/32 of Section 8). (c) Counts,  $R(f) \geq 0$ . (d) The empty reads zero:  $R(0) = 0$  on the zero count vector. (e) Some input reads:  $R$  is not identically zero. Without (d) the hypotheses admit a detector that always reads 1, which satisfies (a), (b) with  $A_2 = 1$  and (c) and is no quadratic form; without (e) they admit the zero detector, under which (b) holds for every  $A_2$ ; so (a) to (c) alone do not force the square. (d) and (e) are what the engine does (the Gram form on the empty record is 0; a click needs a live unit) and what the proof's first step uses ([6, record 340]; the derivation's 6.5 restated with them). Not assumed: continuity, a dimension, the form of  $R$  on one row, or Theorem 2's  $A = \sum a_i^2$ .

**Theorem 4** (The quadratic read-out on the phase lattice, a Gleason-like characterization). *Under (a) to (e), with  $N_\phi$  a power of two,  $A_2 = 2$  and*

$$R(f) = \sum_{j \text{ odd}, 1 \leq j < N_\phi/2} c_j |\sigma_j(f)|^2, \quad c_j \geq 0, \quad \sigma_j(f) = \sum_p f_p e^{2\pi i j p / N_\phi}, \quad (11)$$

the  $\sigma_j$  the Galois conjugates of  $\text{ev}$ ,  $\sigma_1 = \text{ev}$ . Every such  $R$  is a positive quadratic form on the lattice, homogeneous,  $R(af) = a^2 R(f)$ ; it conserves the total at every declared split of one row or of two recombining inputs exactly if and only if the multiplicity factor is  $A = \sum a_i^2$ , and at every label rotation with the factor  $C'^2 + S'^2$ ,  $C'$  and  $S'$  the half-angle tables of Definition 2. Two rows  $ax^p$  and  $bx^{p+\Delta}$  at one Node,  $\Delta$  their phase difference in steps, read  $R = C_\Sigma(a^2 + b^2) + 2ab\mathcal{K}(\Delta)$  with  $C_\Sigma = \sum c_j$  and  $\mathcal{K}(\Delta) = \sum c_j \cos(2\pi j \Delta / N_\phi)$ ,  $\mathcal{K}(0) = C_\Sigma$ ,  $\mathcal{K}(N_\phi/4) = 0$ ,  $\mathcal{K}(N_\phi/2) = -C_\Sigma$  for every admissible  $R$ ; the Born form  $|\text{ev}(f)|^2$  is  $c_1 = 1$  and the other  $c_j = 0$ .

*Proof.* Sketch; the derivation's 6.5 [19] gives the steps in full under (a) to (e). (i)  $g = 0$  in (b) with (a) gives  $2R(f) = A_2(R(f) + R(0)) = A_2 R(f)$  by (d), so  $A_2 = 2$ , one  $f$  with  $R(f) > 0$  sufficing, and (e) supplies it. (ii) Replacing  $g$  by  $x^{-N_\phi/4}g$  in (b) and using (a) and  $x^{-N_\phi/2} = -1$  gives  $R(f - g) + R(f + g) = 2R(f) + 2R(g)$ , the parallelogram law on the abelian group  $\mathbb{Z}[\zeta_{N_\phi}]$ . (iii) The parallelogram law alone makes  $R$  a quadratic form: on a lattice the argument of Jordan and von Neumann [75] needs no continuity, and  $R(f) = B(f, f)/2$  with  $B$  a  $\mathbb{Z}$ -bilinear symmetric form, the restriction of the real form of a symmetric matrix  $\mathbf{G}$ , the click's Gram matrix (not the fan's grain  $N_\theta$  of Section 2). (iv) By (a),  $\mathbf{G}$  commutes with the signed permutation of the basis that multiplication by  $x$  is, whose eigen-functionals are the  $\sigma_j$ ,  $j$  odd; so  $\mathbf{G}$  is diagonal in that basis with real entries paired under  $j \leftrightarrow -j$ . (v) A negative  $c_j$  would make the form negative on an open cone containing lattice points, against (c). (vi) The conservation at a general split and at a label rotation follows from homogeneity, (a) and the cancellation of the cross terms between the two outputs. (vii) Two rows:  $|\sigma_j(a + bx^\Delta)|^2 = a^2 + b^2 + 2ab \cos(2\pi j \Delta / N_\phi)$ .  $\square$

**What is derived and what is not.** Derived from the update rules and the axiom (b): the reading is a positive quadratic form of the record's element, forced by the rotation, the balanced splitter's conservation and the counts alone, with no continuity and no dimension (Gleason's theorem [68] needs dimension three; the lattice version does not, the parallelogram law being granted by the balanced splitter); the multiplicity rule  $A = \sum a_i^2$  and the square as the only power are derived, and the cross term  $2ab\mathcal{K}(\Delta)$  is forced at the click. Not reached by the algebra: the constants  $c_j$ , the detector's response to the harmonics of the phase; a single harmonic  $j$  is the Born form under the relabelling  $p \rightarrow jp$ , a mixture a different law, the flat mixture the phase-blind detector, which




Figure 4: Interference read at a detector, from the runs of series L at  $N_\phi = 64$  (the figures' runs, Appendix C). Left, the two-slit world `slits_huygens`: above, the record's weight per pixel of the screen, a number of the board (GAMEBOARD, a diagnostic); below, the clicks over 64 births (DETECTOR) and the histogram conditioned on the screen, bright where the two rows' ages differ by whole turns. Right, the Mach-Zehnder worlds: the clicks over 64 births at the two ports and the absorber (DETECTOR), equal arms 64/0, a half turn 0/64, a quarter turn 32/32, the (1, 1) and (3, 4) splits, the unequal arms and the two Elitzur-Vaidman worlds (Table 2, row 2b; Appendix C).

interference excludes (Table 2, row 2a; the built click). As  $N_\phi$  grows the admissible readings are the same family, the cone of the even positive-definite functions on the circle with odd harmonics only; no  $N_\phi$  and no continuity forces  $c_j = [j = 1]$ , and the fundamental is what the register reads.

**Against the registered integers.** The built click carries the square: the (3, 4) split's 63/1, 31/1 and 125/3 at  $N_\phi = 64, 32$  and 128 [6, record 363] hold if and only if the power  $k$  lies in  $[1.917, 2.012]$ , containing 2 and excluding 1 and 3, a check that the implementation is the form of Theorem 4 and no evidence about nature's power (Table 2, row 2c); the norm range of Appendix B bounds one phasor's error between the ideal reading and the tables; the built weight is within  $A\sqrt{2R^*} + A^2/2$  of the exact weight  $R^*$  for a record of total amount  $A$  (a part in 181 at full scale, the one-row extreme a part in 276), with no relative bound where the record's rows cancel below  $A/2$  per component:  $256x^8 - 181(1 + x^{16})$  at  $N_\phi = 64$  reads 0, 196 and 25028 at three rotations against the exact 49 ([31], a computation from the tables, not a run). The two-slit period 23.5 pixels and Malus at 22.5 degrees (219/256 exact) admit the harmonic  $j = \pm 1 \bmod 8$  and exclude every mixture of weight above  $w^* = 1/1516 = 0.00066$  on  $j = \pm 3 \bmod 8$  ( $w$  the part of  $\sum c_j = 1$  off the fundamental's class, read as the count of 256 that falls from 219 to 218), and the two-slit bands a weight above about 0.02 on  $j = \pm 1 \bmod 8$  (one click at the peak of 51); below these weights the register does not exclude a mixture, Malus at 22.5, 45 and 90 degrees reading every member of  $j = \pm 1 \bmod 8$  alike ([31], a computation): to these weights the register pins the fundamental, and the least-rank click (P10) stores it.

**Interference, by the formulas alone.** Every phenomenon of this paper is read the same way: Outside, what a detector counts; down to Inside, the rows, their phases, the merge and the evaluation; back Outside, the click's counts over the births against the reading. Two slits: one record is born with two rows of amount 1, one per opening,  $m = 2$ ; each row's phase advances one turn per Link; at pixel  $x$  the two arrive with phases  $p_1(x), p_2(x)$ , their difference  $\Delta(x)$  the path difference times the turn; the merge adds them and an equal pair with  $\Delta = N_\phi/2$  cancels to no row; the click evaluates the sum and squares it, the offer  $R(x) = 512 [(C[p_1] + C[p_2])^2 + (S[p_1] + S[p_2])^2] = 1024 (65536 + C[p_1]C[p_2] + S[p_1]S[p_2])$  up to the tables' rounding, proportional to  $1 + \cos \Delta(x)$ ; the

rungs  $b_x$  of Eq. (9) lay the pixels on the ladder, the click is the pixel under which  $u$  falls, and over  $N_\phi$  births the count at  $x$  is  $b_x - b_{x-1}$ , Born's to  $1/N_\phi$ , with the exact tables. With the declared tables at  $1/256$  the weights vary with the birth phase within the tables' rounding (hypothesis (a) of Theorem 4 holds only to that rounding), so the count over the births departs from the rung difference by up to that rounding: two offers of equal ideal weight at the phases  $u$  and  $u + 1824$ ,  $N_\phi = 8192$ , read 4104 and 4088 for 4096 each, a departure of  $8/N_\phi$  (a computation from Eqs. (10) and (9), not a run); the built click's bound is the tables' rounding, the theorem's  $1/N_\phi$  its ideal. The spacing is the law's own:  $\lambda = cN_\phi d/n$  Links, one turn of the phase per age per fringe, Young's  $\lambda D/s$  only in the paraxial limit (rung 2, the fan) [20]; the bands' centres are 23.5 pixels apart for the exact two-path law's 23.3 (L2b, DETECTOR). What passes is a pixel with  $\Delta$  near 0, a wide rung; what does not is a pixel with  $\Delta = N_\phi/2$ , the rows cancelled, a rung of width 0 where no click can land; one slit is one row per pixel, nothing cancels, a flat count within the fan's grain and no dark pixel; the Mach-Zehnder's two ports are the same algebra at two Nodes, the bright port 64 of 64 clicks and the dark port 0 (row 2b; Appendix C).

**Planck and de Broglie, as identities.** Under the declared dictionary the release's cost rule reads as  $E = h_q s = (h_q N_\phi) f$  and the turn rule as  $\lambda = h_A/p$ : Planck's and de Broglie's relations are identities of the update rules with one constant,  $h = h_q N_\phi = h_A$ , an input (Table 1); no detector reading of them is registered.

**The uncertainty relation.** On the identification of the circle as position and its transform as momentum, a hypothesis of the program and not a derivation ([19], section 22), the support bound  $|\text{supp } f| |\text{supp } \hat{f}| \geq N_\phi$  [69], the entropic bound [74] and Kennard's form  $\Delta x \Delta p \geq \hbar/2$  in the limit [70, 73] are the transform's mathematics on  $\mathbb{Z}_{N_\phi}$  and nothing of the law; the entropy identity of the click, bits read plus bits not read of  $u$  equal to  $\log_2 N_\phi$  per record, is an identity of the wheel. What makes it an uncertainty is the one read-out: a record read at a Node gives its position and loses its phase, read on the fan's angle it gives its momentum's direction and not its Node.

## 7 Bell and CHSH: from weights to counts

The correlation at the settings  $(a, b)$  is  $E = (c_{++} + c_{--} - c_{+-} - c_{-+})/N_\phi$  over  $N_\phi$  births, one per  $u$ , and the CHSH sum is  $S = E(a, b) - E(a, b') + E(a', b) + E(a', b')$  at the labels  $(0, N_\phi/8, N_\phi/4, 3N_\phi/8)$ . Two theorems follow from the definitions, exact on the GameBoard.

**Theorem 5** (Exact count marginals). *For a pair with labels  $\{0, 1\}$  of equal weight and settings  $a, b$ , let  $J(o_A, o_B) = \sum_\ell U_a[o_A][\ell] U_b[o_B][\ell]$  and  $R(o_A, o_B) = J^2$ . Then the first party's outcome is  $+$  exactly for  $u < N_\phi/2$  and  $-$  for  $u \geq N_\phi/2$ , for every  $(a, b)$  and every  $N_\phi$ ; and the second party's count of  $+$  is exactly  $N_\phi/2$  for every  $(a, b)$  unless  $2N_\phi R(+, +)/C_K$  is an odd integer (a tie of the rung), in which case it is  $N_\phi/2 + 1$ . No tie occurs for any setting pair at the twelve grains from 8 to 1024 of [5], nor at the CHSH labels for any multiple of 8 up to 4096.*

*Proof.* Write  $n_a = C'[a]^2 + S'[a]^2$ . Since the rows of  $U_b$  are orthogonal on the integers,  $\sum_{o_B} J(o_A, o_B)^2 = n_a n_b$  for each  $o_A$ , so  $C_K = 2n_a n_b$  and the coarse cumulative after the first party's  $+$  is  $C_K/2$ :  $b = \lfloor (N_\phi C_K + C_K)/(2C_K) \rfloor = \lfloor (N_\phi + 1)/2 \rfloor = N_\phi/2$ . For the second party,  $R(o_A, o_B) = R(-o_A, -o_B)$  (the rows of  $U_s$  are each other's quarter turn), so with  $x = N_\phi R(+, +)/C_K$  the counts are  $c_{++} = \lfloor x + \frac{1}{2} \rfloor$  and  $c_{-+} = b_3 - N_\phi/2$  with  $b_3 = \lfloor (2N_\phi(C_K - R(+, +)) + C_K)/(2C_K) \rfloor = N_\phi + \lfloor \frac{1}{2} - x \rfloor$ . Hence  $c_{++} + c_{-+} = N_\phi/2 + \lfloor y \rfloor + \lfloor 1 - y \rfloor$  with  $y = x + \frac{1}{2}$ , which is  $N_\phi/2$  when  $y \notin \mathbb{Z}$  and  $N_\phi/2 + 1$

when  $y \in \mathbb{Z}$ . The tie condition is  $2x \in 2\mathbb{Z} + 1$ , that is  $2N_\phi R(+,+)/C_K$  odd. The absence of ties at the listed  $N_\phi$  is the computation over every setting pair [5].  $\square$

No-signalling in the arrival intervals and in the counts: the arrival interval at either arm is that arm's flight alone, independent of the other arm's setting and length (Definition 3, the flight blind) (that interval the host's tick at the arm's set, GAMEBOARD [6, record 768]; the arm's own count is read only under the later key `clock_stamp`, outside this paper's law); neither party's recorded outcome exists as a row before the completion, whose interval is the last arm's, so the availability of the record, unlike the arrival, waits on the far arm's length, and the paper claims no independence for that latency; the first party's outcome is determined by  $u$  at the record's birth (Theorem 5) and the second's at the completion by the joint weights, both written at the completion, with the marginal  $N_\phi/2$  for every setting (Theorem 5); so neither the counts nor the arrival times at either arm carry the other arm's setting or length, and the one non-local step of the law (P6) is the assignment of the second party's outcome from the joint weights at the completion, on the labels; its interval, the record's, is the last arm's. What the theorem does not cover, stated here. The wheel of the registered pair worlds runs at the rate 1 against  $N_\phi$ , one birth per  $u$  in order, so the first party's outcomes over the births are  $N_\phi/2$  of  $+$  followed by  $N_\phi/2$  of  $-$ , a square wave whose serial correlation over a cycle is  $1 - 4/N_\phi$  (0.9990 at 4096), a prediction of the sequence in time that no run reads and no row compares with nature's record of outcomes, which shows no such order. The marginal is for settings fixed or chosen independently of the wheel: a setting chosen from  $u$  ( $a = 0$  for  $u < N_\phi/2$  and  $a = N_\phi/2$  after,  $b = 0$ ) gives the second party  $+$  at every one of the 64 births at  $N_\phi = 64$ . On the register the settings are declared per world or balanced over every  $u$  (the choosers' 960 births meet every  $u$  with every pair once), so nothing chooses a setting from  $u$ ; nature's freedom-of-choice loophole has its counterpart here in the wheel, not closed by the register, and the reading that would test it is a marginal off  $N_\phi/2$  with the settings chosen from  $u$  [44]. And the order of the second party's outcomes carries the first party's settings even when they are independent of the wheel: the cells give  $B = A$  at  $a = b$  and  $B = -A$  at  $a = b + N_\phi/2$  exactly, so with  $b = 0$  a first party keeping  $a = 0$  leaves the second a serial correlation  $\langle B_t B_{t+1} \rangle = 15/16$  over 192 births at  $N_\phi = 64$ , and one cycling  $a = N_\phi/2, 0, 0$  (the period 3, coprime to  $N_\phi$ , the independence criterion above) leaves  $-5/16$ , the marginal 96/96 in both, the same at every shift of the wheel and of the chooser (a computation from the theorem's cells, not a run): the second party reads one bit of the first's choice from the order of its own outcomes. So the theorem's no-signalling is of the counts over a cycle and of the arrival times; the sequence in time signals under the sequential wheel, a result of the law as built and a FAIL of any claim beyond the counts, open until the wheel is other than a counter, which P4 (no draw) does not allow, or the claim is kept to the counts, as it is here. Stated plainly: the law as built, with its sequential wheel, does not meet operational no-signalling; its counts are blind to the other party's setting, the order of its outcomes is not, and until the wheel is other than a counter the pair's construction models a Bell experiment in its counts only, not in its record of outcomes. Read after a detector on the registered pair world (Table 2, row 1c): the second party's serial correlation 15/16 under a constant setting and  $-1/16$  under the period-3 cycle 0, 21, 42 (the cycle  $N_\phi/2, 0, 0$  the register's keys cannot declare; the declarable cycle reads the same channel with a smaller swing), the marginals 96/96 under both, the algebra met at 16 of 16 checks, nature's value 0: FAIL [28].

The strict-crossing rung  $N_\phi C_k > u C_K$  would give the second party 33/64 in 3944 of the 4096 setting pairs at  $N_\phi = 64$ , a signalling of  $1/N_\phi$  made by the rounding [3]; the nearest rung is what the theorem needs. The theorem's scope is what it states: a pair of two labels with equal weights, the CHSH form, and the rung of (9); unequal weights and more than two labels are not covered, and no-signalling for them is open. That the marginal at  $B$  is  $N_\phi/2$  for every  $a$  although  $B$ 's rung




Figure 5: The pair at  $N_\phi = 64$ :  $E(a, b)$  computed from the rule at every setting difference  $a - b$  (the counts of Theorem 5 with the repository's tables, a computation [5]), the tables' cosine  $C[a - b]/256$  dashed, and after a detector the registered CHSH worlds (circles) and the choosers' bins (squares) of series L (DETECTOR).

is computed from  $R(o_A, o_B)$  with  $a$  and  $o_A$  supplied by the layer (Definition 3) is the content of the theorem; the joint counts are not local, and the model does not claim they are.

**Theorem 6** (The finite- $N_\phi$  Bell value). *With the counts of Theorem 5 and off a tie,  $E_{N_\phi}(a, b) = 4c_{++}/N_\phi - 1$  and*

$$|E_{N_\phi}(a, b) - \cos(2\pi(a - b)/N_\phi)| \leq \frac{2}{N_\phi} + 4 \arcsin \frac{\sqrt{2}}{2\varrho} < \frac{2}{N_\phi} + 0.0111, \quad \varrho = N_t - \frac{\sqrt{2}}{2}, \quad (12)$$

so at the labels  $(0, N_\phi/8, N_\phi/4, 3N_\phi/8)$ ,  $|S(N_\phi) - 2\sqrt{2}| \leq 8/N_\phi + 0.0444$ .  $S(N_\phi)$  is an exact rational for each  $N_\phi$ ; it lies above  $2\sqrt{2}$  for 252 of the 512 multiples of 8 up to 4096 (among them  $N_\phi = 16, 32$  with  $S = 3$  and  $N_\phi = 128$  with  $S = 23/8$ ) and equals  $181/64 = 2.828125$  at every power of two from 512 through 8192 and  $5793/2048 = 2.828613$  at 16384 [5]; measured after a detector at 512, 1024, 2048, 4096, 8192 and 16384 [4] (Figure 6).

*Proof.*  $c_{++} = \lfloor x + \frac{1}{2} \rfloor$  with  $x = N_\phi R(+, +)/C_K$  gives  $|c_{++} - x| \leq \frac{1}{2}$  and  $E_{N_\phi} = 4c_{++}/N_\phi - 1$  from the mirror symmetry, so  $|E_{N_\phi} - (4R(+, +)/C_K - 1)| \leq 2/N_\phi$ . The vector  $(C'[s], S'[s])$  differs from  $256(\cos \theta_s, \sin \theta_s)$ ,  $\theta_s = \pi s/N_\phi$ , by at most  $\frac{1}{2}$  per component, so its direction differs from  $\theta_s$  by at most  $\delta = \arcsin(\sqrt{2}/(2\varrho))$  with  $\varrho = N_t - \sqrt{2}/2 = 255.29$ , below every table vector's length (each component within  $\frac{1}{2}$  of  $256 \cos$  or  $256 \sin$ ), so  $4\delta = 0.01108 < 0.0111$  and  $16\delta = 0.04432 < 0.0444$ ;  $J(+, +)/\sqrt{n_a n_b}$  is the cosine of the angle between the two directions,  $\cos(\theta_a - \theta_b + \epsilon)$  with  $|\epsilon| \leq 2\delta$ , and  $4R(+, +)/C_K - 1 = 2 \cos^2(\cdot) - 1 = \cos(2(\theta_a - \theta_b) + 2\epsilon)$ , within  $4\delta$  of  $\cos(2\pi(a - b)/N_\phi)$ . The four settings add the bounds. The values of  $S(N_\phi)$  are the computation from these formulas with the repository's tables [5], which reproduces the design's  $176/64$ ,  $720/256$  and  $2896/1024$  at  $N_\phi = 64, 256, 1024$  and the engine's three values below.  $\square$

What Theorem 6 does not say: that  $S(N_\phi) \leq 2\sqrt{2}$ , false for about half the  $N_\phi$ ; nor that the deficit is  $O(1/N_\phi)$  from below. The tables' term in (12) is the resolution  $1/256$  of the circle's tables and does not shrink with  $N_\phi$ ; at the CHSH labels the tables' entries are the same at every  $N_\phi$ , so the rounding gives  $181/64$  at the powers of two from 512 through 8192 and  $5793/2048 = 2.828613$

at 16384 and 32768 (measured at 16384, DETECTOR), and beyond the tables' bound  $2N_\phi \leq 65536$  the rungs on the fixed correlations are the closed form  $S(N_\phi) = 8(c_1 + c'_1)/N_\phi - 4$  with  $c_1, c'_1$  the first cell's counts, whose limit with the tables fixed is  $186034/65773 = 2.828425$ ,  $2.1 \times 10^{-6}$  below  $2\sqrt{2}$ ;  $2\sqrt{2}$  is the joint limit in  $N_\phi$  and  $N_t$  [5, 4]. At a tie  $c_{--} = c_{++} - 1$  and  $E_{N_\phi}$  drops by  $2/N_\phi$ ; no tie occurs at the CHSH labels up to 4096.

**The finite-grain value, and what would make it a prediction.** The construction's number is

$$S_{U24} = \frac{181}{64} = 2.828125 \text{ exactly,} \quad \Delta S = S_{U24} - 2\sqrt{2} = -3.02 \times 10^{-4}, \quad (13)$$

at every power of two from 512 through 8192, with the tables at  $N_t = 256$ , the wheel at  $N_u = N_\phi$ , the settings  $(0, N_\phi/8, N_\phi/4, 3N_\phi/8)$  and a pair of two labels with equal weights; the marginals exactly  $1/2$ ; the correlations at the two pairs of settings 725/1024 and 723/1024 at  $N_\phi = 4096$ , above and below  $1/\sqrt{2}$ ; 5793/2048 at 16384 and 32768. It was computed in advance and measured after a detector at  $N_\phi = 512, 2048, 4096, 8192$  and, the plateau's end, 16384 (the last three [4, at b34fe114]; Figure 6), under the click and the wheel with every pin met (the register's block of the derivation's 24.4); 512 and 4096 were re-run by the second runner [9]. The criterion: the law at a grain is rejected when a measured  $S$  differs from  $S(N_\phi)$  by more than five standard errors, under the identification of the experiment's settings and detection with the labels and the ideal read-out, which the model does not model; the comparison with nature is Section 8. It is a prediction of the law only once the pair's wheel meets no-signalling in the order of the outcomes (above), which the sequential wheel as built does not; until then it is the exact consequence of the declared construction at its grain.

## 8 The simulator's checks and the comparison with nature

Three things are kept apart: the gates on the code; what is shown Inside by algebra under named assumptions, which needs no experiment; and what is measured, a run Inside read Outside, compared with known experiments.

**The code implements the law.** Every registered world is run from its world file with its expectation written before the run and its source fingerprint; a second runner re-ran series L, T and fifteen more worlds bit by bit [9]; the books balance at every interval in every run; the rounding pins of the (3,4) split were met (Appendix C); the crossing rule's tests read their pinned counts. These are gates on the code, not measurements.

**The formulas after a detector.** Every number read after a detector is a count of clicks or a click's tick at a declared detector, registered with its source fingerprint (Appendix C; [4, 9]), its expectation written before the run; a number of the GameBoard's state appears nowhere. Seven formulas shown Inside (the pace, the click's power, Mach-Zehnder, the marginals, GHZ, Malus, light beside a mass) were confirmed by their runs (Appendix C); the readings that no formula gives are Table 3's (the clock's field at two Nodes, the equivalence after a detector, the covariant readings) and Figure 6's ( $S$  at seven grains).

**The comparison with nature.** The confrontation register [48] keeps one row per registered detector reading with a dimensionless counterpart in nature, one published source per value and one verdict per row (PASS within the stated uncertainty, FAIL with the number, BOUND where




Figure 6:  $S(N_\phi)$  at the CHSH labels. The curve is the closed form  $S(N_\phi) = 8(c_1 + c'_1)/N_\phi - 4$  computed from the rule at every multiple of 8 with the tables at  $N_t = 256$ , a computation [5]; the circles are detector readings (DETECTOR) at  $N_\phi = 64, 512, 1024, 2048, 4096, 8192$  and  $16384$  (series L, L6 and the register's block of the derivation's 24.4, the last three [4, at b34fe114]); the inset marks the plateau  $181/64$  from  $512$  through  $8192$  and its end  $5793/2048$  at  $16384$ ; the dashed line is  $2\sqrt{2}$ .

the comparison bounds a free parameter, NOT COMPARED where the law does not compute the observable); a FAIL is a result of the law and is stated with the same care as a pass. Table 2 carries the rows read after a detector; six rows of the register (4a, 5b, 10, 11a to 11c) rest on a pin whose run is not made and are not in this table, four of them FAIL by their pin (4a, 5b, 11a, 11c); every verdict is the law's as built at the commit of Appendix C, and a hypothesis's PASS beside the law never changes the law's FAIL.

Table 2: The confrontation register's rows read after a detector [48]. The criterion per row: PASS where the reading meets nature's value within the stated uncertainty, or, for a visibility, where the model's value lies above the apparatus-limited measurement (a bound met); the model's numbers are ideal, at the registered grain, not a prediction for a given instrument. The one count, computed from the rows and repeated verbatim in the abstract and in Section 9: three readings pass (1a, 2b, 9); twelve fail in the table, eleven in a registered run (1b, 1c, 2a, 3, 4b, 6, 7b, 8a, 8b, 13, 14) and one a declared input refuted (8c), with four more failing by a pin whose run is not made and not in the table (4a, 5b, 11a, 11c); two are bounds (5a, 7a); one is not compared (2c); one is history (12, the lattice pin met under the law before the entry, not a pass against nature); the hypotheses beside rows 6, 13 and 14 change no verdict of the law (the PASS rows replicated [9]). Each reading's kind is named: DETECTOR (a click), GAMEBOARD (the host's view, a diagnostic) or COMPUTATION (the algebra's); the criteria row by row, with the re-runs at head, are [44].

| Row | Observable and nature's value                                         | The law's detector reading                                                            | Verdict                                                              |
|-----|-----------------------------------------------------------------------|---------------------------------------------------------------------------------------|----------------------------------------------------------------------|
| 1a  | The CHSH sum of a pair: $2.42 \pm 0.20$ [86]; the quantum bound 2.828 | $S = 2.75$ at $N_\phi = 64$ , series L (DETECTOR; replayed at head by the test suite) | PASS: 1.65 standard errors above the measured, 0.078 below the bound |
| 1b  | The same, the phase-form window                                       | $S = 2$ exactly (DETECTOR, the counters' clicks)                                      | FAIL: the local bound, 2.1 standard errors below                     |

| Row | Observable and nature's value                                                                                                                                                                                                                                                                                                                                                    | The law's detector reading                                                                                                                                                                                                                                                                                                                                                                                | Verdict                                                                                                                                                                                                                                                                                                                           |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1c  | No signalling in the order: nature 0 on the second party's serial correlation at lag 1 under the first party's settings (the outcome order of a pair random, carrying no setting of the other party; the marginals' independence holds in nature and in the law alike)                                                                                                           | 15/16 under a constant $a = 0$ and $-1/16$ under the period-3 cycle $a = 0, 21, 42$ , cyclic over 192 births on the registered pair world at $N_\phi = 64$ (DETECTOR, the run of 2026-09-22 [28]); the marginals 96/96 under both (DETECTOR); the algebra's 15/16 and $-1/16$ met at 16 of 16 checks; the cycle $N_\phi/2, 0, 0$ of Section 7, $-5/16$ , a computation the register's keys cannot declare | FAIL: the law's determinism, the wheel a counter, shows in the order and not in the counts                                                                                                                                                                                                                                        |
| 2a  | Two-slit visibility of one quantum at a time: 0.98, the register's source [87] a Mach-Zehnder reading and not a two-slit one; a two-slit source, a Fresnel biprism's central fringe at 0.94 as reported [45], its figure to be verified against the source; the ideal 1; the criterion the clicks' visibility $(I_{\max} - I_{\min})/(I_{\max} + I_{\min})$ , no apparatus model | 0.966 in the clicks of <code>slits.huygens</code> (the birth wheel, 4096 births; L2b; DETECTOR), the pin derived for this world's fan by the register's own algebra 0.9659 (COMPUTATION) and met bit for bit by the re-run at head, 0 of 121 pixels differing; the record's weights read the same under the exact phase (the 0.954 of the built-phase pin of 2026-09-20 is history)                       | FAIL against the Mach-Zehnder's 0.98: 0.014 below the measured, 0.034 below the ideal; the cause the fan's discreteness, the digital lines' landings; against the two-slit source NOT COMPARED until its figure is verified, and then bound against bound, the biprism's central fringe apparatus-limited as the fan's clicks are |
| 2b  | Mach-Zehnder visibility: 0.98 [87]; the criterion the visibility $(b - d)/(b + d)$ of the bright and dark ports' offers of <code>mz_equal</code> , no apparatus model                                                                                                                                                                                                            | the offers 1681/1682 and 1/1682 (GAMEBOARD, the record's ledger); the clicks 64/0 over 64 births (DETECTOR; replayed at head by the tests)                                                                                                                                                                                                                                                                | PASS as a bound met, not a quantitative agreement: the clicks' visibility 1.000 (64/0, DETECTOR) above the apparatus-limited 0.98; the offers' 0.9988 a diagnostic of the apparatus layer                                                                                                                                         |
| 2c  | The power of the click's form: the Sorkin parameter $\kappa = 0.0064 \pm 0.0119$ [60]                                                                                                                                                                                                                                                                                            | the power's window [1.917, 2.012] from the click cells (DETECTOR), a read-back of the built click (COMPUTATION)                                                                                                                                                                                                                                                                                           | NOT COMPARED ( $\kappa$ not computed): the window an implementation gate, no evidence about nature; no three-opening world is registered                                                                                                                                                                                          |
| 3   | The deceleration parameter $q_0 = -0.53 \pm 0.01$ [61, 62, 63]                                                                                                                                                                                                                                                                                                                   | $-0.108$ coasting (series G2, the pointer's $z$ : the births DETECTOR per tick of the open-face detector, that tick the host's interval, GAMEBOARD [6, record 768], the detector's own pulse-and-return clock NOT READ); $-0.104$ at head ([6, record 408]; the re-run at head, the pin met, DETECTOR), the reading's own grain 0.004 between engines and 0.03 between windows                            | FAIL: 0.42 from nature's value, the coasting ideal $q = 0$ and the reading within its grain of it; the law has no term carrying $q$ toward $-0.5$                                                                                                                                                                                 |
| 4b  | A moving lamp's redshift with the clock's factor [57]                                                                                                                                                                                                                                                                                                                            | $z = 0.2636$ at $\beta = v/c = 0.2674$ (series G2: the births DETECTOR per tick of the open-face detector, that tick the host's interval, GAMEBOARD [6, record 768], the detector's own pulse-and-return clock NOT READ); 0.2647 at head (the re-run, the pin met, DETECTOR)                                                                                                                              | FAIL: 0.315 expected, 17 grains below; under covariant-readings-v1 0.3674 for the pinned $0.369 \pm 0.003$ (series S), the law's row FAIL; the frame: the two one-way factors apart by $1 - v^2$                                                                                                                                  |

| Row | Observable and nature's value                                                                                                                 | The law's detector reading                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         | Verdict                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 5a  | The anisotropy of $c$ by direction, below $10^{-18}$ [56]                                                                                     | the flight table's pace 0.5774 to 0.5818 at $N_l = 64$ (COMPUTATION; series Q's 290 of 290 clicks within it at finite ages, DETECTOR, Appendix C)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  | BOUND on $N_l$ : of order $5.8 \times 10^{17}$                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| 6   | Bohr's line ratio $\nu(\text{H}\beta)/\nu(\text{H}\alpha)$ of the Balmer series: 1.3500, 656.279 over 486.135 nm [43], Balmer's 27/20 exactly | by algebra first, on the law's integers (COMPUTATION [36]): a closed loop is an integer congruence of the action row, $\Delta A_a = 0 \bmod h$ on each axis with the quotients summing to $jN$ ; two closing radii stand as the square of two counts a detector reads, $a_j/a_i = (j/i)^2$ in the shell mean, checked by $T_j/T_i = (j/i)^3$ , no root, no float; from a closed circle one click reaches only the loops with $2i^2 \geq j^2$ and a release of content unbinds, so no line of Balmer's is one click of the law: NEEDS A NEW RULE. The law as built: the electron escapes at 3407 intervals (series H's world, DETECTOR; the cause the half-Link lag [30]). Under the key <code>centred_step</code> (a declared start of the drive, off by default; Section 9): the loop stays, no escape face click in 7500 intervals, 23 quarter crossings (C1 PASS, DETECTOR), tightening from 12 to 7 Links at two crossings over five turns, the period falling from 1431 to 1040 (C2, C3 FAIL as declared, DETECTOR; the run of 2026-09-22 [29]); no line read | FAIL, the law's: the loop opens at 3407; under the key C1 PASS, C2 and C3 FAIL, no pin moved; the lines NOT COMPUTABLE by any run of the law as built (no rule of the six quantises a body's loop; 27/20 waits on the new rule); the ladder $a_j/a_i = (j/i)^2$ , $E_j \propto 1/j^2$ and Balmer's 27/20 conjectured from the algebra in the shell mean's limit, computed from the algebra, not read on the board; a lattice able to hold the scale (the orbit's radius sixty thousand times the nucleus's) is not run (Section 9 (viii)) |
| 7a  | The deuteron's binding fraction 0.1185 percent [58]                                                                                           | the bond's two border clicks and the mass read 3673 (DETECTOR) against the declared 3677, 4 units, 0.109 percent (series N); the books' escaped 4 (GAMEBOARD, a diagnostic) the check; the same at head, every pin met                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             | BOUND on a declared input: nature's 4.35 units of 3677 reachable by no whole give                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| 7b  | The alpha's binding over the deuteron's, 12.72 [58]                                                                                           | 2.0 (series N, the border's four clicks against two, DETECTOR; the same at head)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | FAIL: a factor 6.4; the give once per body                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

| Row | Observable and nature's value                                                                                                           | The law's detector reading                                                                                                                                                                                                                                                                                                                                                                                                                       | Verdict                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|-----|-----------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 8a  | The neutron's decay curve, exponential, width over median 3.17 [66]                                                                     | a step: every neutron at its key on its own clock, the width 0 (DETECTOR, the 64 clicks of content 3); the width over the median in the lattice's clock 0.036 in the registered run and 0.08 to 0.13 at head at the cap 1024 (GAMEBOARD, the lattice's clock, counted in no criterion; against nature's 3.17 these widths would read a factor 88 in the registered run and 25 to 40 at head, numbers of the diagnostic that stand in no verdict) | FAIL: a step, the width 0 against nature's 3.17 (DETECTOR); the registered J1 run predates the one flight wall, under which the weak register's become blocks move ([6, records 931 to 933]), so the row is a reading of the law before it, kept as history, J1's redeclaration in the weak field after the paper                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| 8b  | The neutrino's passage through a second detector, about 1                                                                               | 16 of 1024 with 0 behind (series J2, a window; DETECTOR; the same at head, the pin met bit for bit)                                                                                                                                                                                                                                                                                                                                              | FAIL: 0 against nature's ratio 1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| 8c  | The heaviest neutrino mass state over the electron's mass: at least $9.8 \times 10^{-8}$ [64, 65]                                       | the $\nu$ family with no content, an input                                                                                                                                                                                                                                                                                                                                                                                                       | a declared input refuted by nature: massless states cannot carry the observed splittings; a content above 0 is an input the register does not declare                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| 9   | Malus's law at 45 degrees, $1/2$                                                                                                        | 128 of 256 exactly; 219 of 256 at 22.5 degrees (A12; DETECTOR, replayed at head by the tests)                                                                                                                                                                                                                                                                                                                                                    | PASS: exact at 45 and 90 degrees; $0.8555$ against $\cos^2 = 0.8536$ at 22.5                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| 12  | The clock's field at two distances at the same push: the potential's form, 2.00 (the GPS term 45.7 microseconds per day [14]; [15, 59]) | series T: the form measured at a detector at two Nodes (the births DETECTOR per tick of the open-face detector, that tick the host's interval, GAMEBOARD [6, record 768], the detector's own pulse-and-return clock NOT READ), the age clock 1.907 (the lines' pin $1.909 \pm 0.05$ ), the presence clock 1.000                                                                                                                                  | read under the law before the generic entry of 2026-09-22 [32]; not readable under the law as it stands; the four worlds are being redeclared in the weak field (the pair [1, 16384], the crowd's age moment at most 1024 per Node on the light's path) and re-read, the new numbers to replace these; HISTORY, not a pass: under the age word (P9) as the form at two Nodes the lattice pin $1.909 \pm 0.05$ is met (DETECTOR, a gate on the code, the law before the entry), the lattice's dwelling ages 5, 6 and 10, 11 pinning 1.909 for the continuum's 2.00; against nature the reading 1.907 is 4.6 percent off with no uncertainty stated on either side, so the caption's PASS criterion is met nowhere, PENDING the weak-field re-read; FAIL under the presence word |

| Row | Observable and nature's value                                                                                                                                             | The law's detector reading                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       | Verdict                                                                                                                                                                                                                                                                                                                                                        |
|-----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 13  | The bending of light beside a mass, the full deflection over the time part's alone, $1 + \gamma = 2$ at the Sun's limb, 1.75 arcseconds against the time part's 0.87 [42] | <p>the law as built: a row reads nothing of the crowd, the deflection 0.000 pixel and the delay 0.00 interval (series K, three worlds, DETECTOR). By algebra first, on the law's integers (COMPUTATION): the walk of a light row past a held mass under the key <b>optical</b>, read at a ring of Nodes at the impact distance <math>b</math>, recovers the <math>M/b</math> form with</p> $C_{\text{ring}} = \alpha bc^2 / (GM) = 5.12$ <p>at <math>b = 6</math> (5.44, 5.86 at <math>b = 3, 8</math>), not 4 within the ring's grain, the factor left the fan's <math>F_{L1} = 1.421</math> of the flow [34]; under the one constant (flow-link-v1, the flow counted per Euclidean Link of its line) <math>C_{\text{ring}} = 3.65, 3.82, 3.92</math> at <math>b = 6, 3, 8</math> against 4, that is <math>2c_f</math> within the finite path's <math>L/\sqrt{L^2 + b^2}</math> (0.974 at <math>b = 6</math>) and the shells' 0.990, one to three grains, <math>4 = 2(1 + \gamma)</math> with <math>\gamma</math> an input [35]. The run of 2026-09-22 on the split ladder (the wall's square reached without the root; the first run's deciding ring had been refused by the working bound) [38]: Under flow-link-v1 (the key <b>flow.link</b>, off by default) the six ring worlds read their mean radial shifts at the algebra's numbers (<math>b = 6, \gamma = 1</math>, the deciding pin: 0.731 Links against <math>0.731 \pm 0.025</math>; every arrival Node the map's, 168 of 168; DETECTOR), and the constant through the stated lever-arm factor,</p> $C_{\text{ring}} = 3.655, 3.819, 3.918$ <p>at <math>b = 6, 3, 8</math> and <math>\gamma = 1</math> (COMPUTATION), is the algebra's own walk to 0.01 and 2.40, 1.07, 1.40 grains from the continuum's <math>2c_f \times 0.990 \times L/\sqrt{L^2 + b^2}</math>, outside the one-grain pin: the engine walks the rule as the algebra walks it, and the bending's 4 does not close within the grain (3.655, 3.819, 3.918 bare against 4); <math>c_f = 2</math> is an input, <math>nS = d</math> a declaration, 4 on the comparison side; nothing here is compared with nature.</p> | <p>FAIL, the law's: 0 against 1.75 arcseconds (series K); the ring's rule read (DETECTOR, inside its pins), the bending's constant DOES NOT CLOSE within the grain (the README's rule, the deciding pin inside beside it), <math>\gamma</math> an input; the coefficient <math>2(1 + \gamma)</math> on the one constant stays conjectured from the algebra</p> |

| Row | Observable and nature's value                                                                                                                                                                                                                                                                                              | The law's detector reading                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Verdict                                                                                                                                                                                                                                                                                                                                              |
|-----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 14  | Newton after a detector: the ratio of the periods of two closed loops at $r = 24$ and 12 under the $1/r$ push on the plane, 2.00 (the form's scale symmetry; nature's inverse square in space $2^{3/2}$ ); the register's bracket $2.00 \pm 0.18$ ; the run's loops never closed, so its ratio is not compared with nature | by formula first (COMPUTATION [35]): under the one constant every push is divided by the plane fan's mean of $S_1/ \mathbf{D} $ , $F_{\text{plane}} = 1.2871$ , so the circular balance solves to the whole $n = 8$ and the periods $T = 2\pi r(S + n)/n$ move from 343, 687 to 384, 768 (377, 754 at the whole 8), the ratio 2.00 unchanged, the factor a constant of the line cancelling in it. The run of 2026-09-22 under the key <b>flow.link</b> , series D3's four worlds, the pins by the generator before any run [39]: the controls inside (every click at $x = c_x + r$ ; the escape tick 305 in 299 to 311; DETECTOR); the probes escaped, <b>r12.flow</b> through the face $+y$ at the tick 1208 and <b>r24.flow</b> through the face $-x$ at 2452 (face clicks, DETECTOR, read from the registered events by the diagnosis [40]; the run's first record had read none), so no loop closed, and the readings called periods are the means of the recurrences of $x$ before the escape, two and three unequal spacings, outside: $T(12) = 588.7$ against 343.1 to 410.9, $T(24) = 987.3$ against 686.1 to 821.8, $\omega^2 = 7.775 \times 10^{-5}$ and $3.112 \times 10^{-5}$ against 2.338 to $3.354 \times 10^{-4}$ and 5.845 to $8.386 \times 10^{-5}$ , the amplitudes 32.5 and 45 against 11 to 14 and 23 to 26 (DETECTOR); the ratio $T(24)/T(12) = 1.677$ against 1.82 to 2.18 (COMPUTATION from two means of unclosed loops' spacings, not a reading of the force's exponent; the two loops are not similar figures, their mean radii 24.3 and 37.6, CONVERSION, so the $1/r$ form's scale symmetry is not tested by them); 4 inside, 7 outside, none moved, 0 record checks failed; the registered D3 row at $n = 9$ (7 inside, 9 outside; Section 4's 1.997) stays as history beside it | FAIL as read under the one constant (1.677 against $2.00 \pm 0.18$ ), the law's at the launch pace 0.34 c: the probes left a confining push and no loop closed, the face clicks the reading; the ratio not compared with nature; the law as built: 1.997 in [1.82, 2.18] on the plane's $1/r$ form (Section 4), the inverse square in space NOT MADE |

**The prediction, and the failures.** Against the photon-pair value  $S = 2.82759 \pm 0.00051$  [85], under four assumptions the paper states and does not defend (nature's pairs this model's pairs at some  $N_\phi$ , fair sampling, the four settings the model's labels, the settings independent of the wheel), the law's 181/64 stands 1.05 standard errors above, 16384 and beyond at 1.5 to 2.0;  $N_\phi = 64$  and 256 are excluded; the loophole-free [86] excludes no  $N_\phi$ . The proximity is a compatibility, not an advantage over quantum mechanics; what refutes it is a CHSH measurement at an uncertainty below  $1 \times 10^{-4}$  reading a deficit below  $2 \times 10^{-4}$  or above  $4 \times 10^{-4}$ , and a separation of 181/64 from  $2\sqrt{2}$  at five standard deviations needs an uncertainty below  $6 \times 10^{-5}$  before systematic errors. The value is the model's at its declared table scale  $N_t = 256$  and at the registered grains 512 to 8192; why nature's grain would lie there, or its table scale at  $N_t = 256$ , the paper does not say; but the dependence is itself algebraic and not a fit:  $S(N_\phi)$  is the exact rational of Theorem 6 at every grain, 181/64 at every power of two from 512 to 8192 and 5793/2048 from 16384 to the tables' bound, so what the model predicts is that family of rationals, computable without a run, and a measurement at the stated uncertainty decides among them or refutes the family as a whole: a value below 2.828125 or above 2.828613 by five standard deviations refutes the registered grains, the powers of two from 512 to 32768, at once; the other multiples of 8 give other rationals of the same theorem, 17/6 at 768 and, over the enumerated grains from 1024 to 4096, from 2.82443 to 2.83212 (the check's scan [5], COMPUTATION, not a bound beyond it), so such a measurement decides among the family's members and refutes the plateau, not every grain; and nature's  $2.82759 \pm 0.00051$  sits 1.05 standard errors below the plateau. The failures are the law's own results, every FAIL row of Table 2 by its number: 1b the phase-form window, 1c the order channel, 2a the two-slit visibility, 3 the deceleration, 4b the moving lamp's redshift, 6 the atom's loop opening, 7b the strong ratio, 8a the neutron's step, 8b the neutrino's passage, 8c the massless neutrino, 13 light unbent under the law before the generic entry (Section 5, against 1.75 arcseconds), 14 Newton's periods under the one constant; and by their pins, not in the table, 4a the unsloped clock, 5b the arms' anisotropy, 11a and 11c the far lamp's brightness and Tolman's test; each refutes the law as declared on its own, and no count of passes weighs against them.

## 9 Discussion and conclusions

**Inside and Outside, and how a click arrives at Lorentz.** This paragraph and the two after it are the conditional extension of the law, the click frame under (A1) and (A2): where (A2) enters, the result is not a consequence of the law as built, which meets (A1) only, and each line names it; the  $r$ -free lines follow from (A1), (A3) and the conversion alone. A packet moved Inside, from place to place, and what it says Outside: Inside, the law propagates records as a beam by the algebraic formulas of Section 2, where no one measures; Outside, the game above the board, there are detectors and their clicks only, the beam passing to them with amplitudes at the click (P11). So a thing propagates Inside as a beam, rows on digital lines at one Link per interval, and shows Outside as a wave: what a detector reads is the click's weight, the quadratic form of the rows' phase counts (Definition 3's  $X^2 + Y^2$ , the same integer as the square of a phased sum), which adds and cancels as a phased sum at the roots of unity does, so Young's bands and the Mach-Zehnder's ports are the beam's rows read at the click and nothing else (Section 6); the wave is the read-out of the beam, not a second thing on the board. A detector's clock is its own record's count of what it emits and receives back, no Node holding a time and an open face without a body having none; a velocity is Nodes apart over counts apart between neighbouring detectors; the read-out is the click (Definition 3). The frame [47]: "To convert the lattice and the momentum inside it into motion Outside, the game above the board, one uses clicks that pass information, a click being the

passage of information from Node to Node at most one Node per interval (A1) and its content an amplitude with a phase that splits each interval between staying and hopping, the mass the staying share (A2); Lorentz of the world is assumed, and what is shown is that this conversion brings it. The symmetry of the conversion, the transformations between click families that preserve (A1) and (A2), is the Lorentz group up to scale, acting on the amplitudes of the passing packet as the Dirac walk's covariance, so that the passage of clicks is Lorentz up to corrections of order  $m^2v^2$ , exact in the continuum limit, a detector's own rate  $\sqrt{1-v^2}(1-\kappa^2/6)$  with the mass angle  $m$  the spacing over the reduced Compton wavelength; in three dimensions the lattice's anisotropy begins at the fourth order." (A1) and the one line, the relativity of the two directions ( $k_{AB} = k_{BA}$ ), are special relativity's two postulates in the click language, so what is arrived at is Einstein's road walked on clicks, and what the board adds is (A2), the amplitudes that carry  $\sqrt{1-v^2}$  into a detector's own rate. The law as built meets (A1), one Node per interval with the flight blind to the phase (P9), and not (A2), its hop a whole-record schedule with no staying amplitude; so its clicks carry the group and its counter runs at the rest rate at every speed (rows 4a, 4b), the two one-way Doppler factors read through records apart by  $1-v^2$ , the measurable that decides. No wall of the law and no declared identity enter the reading; Eq. (14) is an Inside formula on the record, read by no detector, whose Outside formulas the conversion gives. What the system gives is a computation Inside whose passage Outside is the clicks, at the quantum level as well; how far Outside represents reality is what Tables 3 and 2 show, result by result.

**The platform: how a formula is arrived at, by the algebra and by the simulator.** Every formula of this paper is arrived at on one platform, by two roads that meet at the detector. The algebraic road has three steps. The Inside step: the six verbs on the record's attributes, the family's declared integers naming which attributes the step reads (Section 2; Table 5). The conversion: an emitter puts a record on the board and a detector ends it at a click, so that Outside there are counts and Nodes only, the detector's own count the age wall's member at coefficient 1 (P11). The Outside form: a ratio or a difference of those counts, the image of the Inside step, with its rung (exact, in the fan's limit, in the shell mean) and its order in the grain; a known form, Einstein's or Young's, stands on the comparison side only. The simulator's road runs the same three steps: a world file declares the families, the emitters and the detectors, and the reading is pinned before the run. A detector at a Node reads each arriving record once, by one comparison, at its own count (series L, Q, X and T; T's four worlds were read before the generic entry of 2026-09-22, are not readable under the law as it stands, and are being re-read in the weak field). A lamp in motion is read at a detector at rest (G, G2, S). A detector in motion reads Nodes apart over counts apart between neighbouring detectors, its own one-way ratio, the measurable of the frame, a reading NOT MADE. A lamp on a body and a line of detectors at rest read the body's own counts as births and its place as the birth's Node (D3). Every reading named here is DETECTOR in its count, the self-creations' ticks of series S GAMEBOARD; where a count is taken per tick of an open-face detector without a body (the redshift rows 3, 4b and 12) the time base is the host's interval, GAMEBOARD [6, record 768], the detector's own clock NOT READ; and a number of the board's state is a diagnostic, never a reading. A formula enters the paper with both roads: the algebra's chain with its inputs and its word (Table 6), the simulator's reading with its series and kind (Table 3 and Figure 6), and nature on the comparison side (Table 2). The platform is the paper's offer beyond its list: to arrive at a further formula, declare the family and the attributes it reads, write the Inside step on them, name the detector and what it counts, take the ratio Outside, pin it before the run, and read it by kind; the three chains below do this in full.

**The step beneath and the step above.** The Inside step of this chain is not Eq. (1) but the click frame's walk, one amplitude per record that splits each interval between staying and hopping (A2), whose exact invariant is  $\cos \omega = \cos m \cos \kappa$ ; Eq. (1), which hops whole, carries  $W$  only as the declared identity of Eq. (14) (Table 6). The paper's centre is the step beneath the lattice, Eq. (14) in its frame, the six operations on the record with  $W$  its invariant; the direction is one way: the step beneath is the axiom, the conversion of (A1) and (A2) is the map, and the step above is its image, a theorem named as what Einstein wrote. From  $W$ , by the conversion, the step above follows:  $Wc^4$  is  $E^2 = E_0^2 + p^2c^2$ , the gate  $E_0/E'$  is  $1/\gamma$ , a detector's own rate, the drive's fraction is the click's velocity, and the two Doppler factors are the step above read at two detectors. The click frame's section 7 [47] shows the chain with no square declared: the walk's exact invariant  $\cos \omega = \cos m \cos \kappa$  gives  $\omega^2 = m^2 + \kappa^2 - m^2\kappa^2/3 + O(6)$ ; the law's own Planck map ( $E = h_q n/d = h_A f$ ,  $p = \hbar \kappa$  per Link,  $E_0 = \hbar m$ , the load identity  $3hn = Qsd$ ) gives  $E^2 = E_0^2 + c^2 p^2$ ; the whole unit  $E' = E/c^2$  gives  $W = E_0'^2 + 3\mathbf{p} \cdot \mathbf{p}$  with  $3 = 1/c^2 = d$  exactly on the body diagonal (2.954, 2.971, 3.000 by direction, the root's rounding), the corrections of relative order  $m^2\beta^2$ , exact in the continuum limit. So  $W$  is shown to that order as the conversion's identity under (A1) and (A2), no square declared anywhere in the chain [23], and Einstein's relation follows from the Inside step under (A1) and (A2), to second order, an extension the law as built does not meet, its rows hopping whole without (A2); for the rows' dynamics, which hop whole and lack (A2),  $W$  remains the declared identity of [6, record 270], the reading's identity and not yet the rows'. In one chain, with nothing of Einstein's put in: the Inside step, the click frame's walk on the record with  $W$  its invariant (one amplitude per record that splits each interval between staying and hopping, (A2), whose exact invariant is  $\cos \omega = \cos m \cos \kappa$ ; Eq. (1), which hops whole, carries  $W$  only as the declared identity of Eq. (14)); the conversion, an emitter and a click at one Node per interval (A1) and the amplitude's split (A2); the symmetry of that conversion, the Lorentz group up to scale (the click theorem); and its image, the Outside step, a detector's place apart and count apart, whose most general form above the board is Einstein's step as physics writes it,  $d\mathbf{p}/dt = \mathbf{F}$  with  $\mathbf{p} = \gamma m \mathbf{v}$ ,  $E^2 = E_0^2 + c^2 p^2$ ,  $d\tau/dt = \sqrt{1 - v^2}$ , and the geodesic's first terms ([21], section 3). Lorentz of the world is assumed, through (A1) and the relativity of the two directions ( $k_{AB} = k_{BA}$ ), and what the chain shows is that the conversion brings it: the Lorentz group up to scale is the symmetry of the click families that preserve (A1) and (A2), and Einstein's equations are the form of the Outside step, one step above the board; nothing here derives the Lorentz group from the six verbs, and "nothing of Einstein's put in" says that no factor of Einstein's enters the chain's steps, not that the symmetry is derived: the step's  $r$ -free lines (the round trip, the composition, the aberration) are Einstein's exactly from (A1), (A3) and the conversion alone, and its scaled lines are Einstein's if and only if  $r^2 = 1 - v^2$ , which (A2) gives to order  $m^2 v^2$  and the law as built, at  $r = 1$ , does not. The same road serves Newton (below) and, in the same font, what it does not reach: length contraction, the perihelion's second order, the bending of light, waves, horizons and the field equations [21].

**The smallest thing above.** What the Outside step is made of follows from (A1) and the click's definition with no run: a click is a Node, a detector's own count and what arrived, and the counts of a family add, so their ratios lie in  $\mathbb{Q}$  [47]. Hence the least distance a click can report is one Link, Nodes apart being an integer; the least time a detector times is one count, a pulse and its return two; the least step of a velocity at  $k$  counts is one Node per  $k$  counts;  $c$  is one Link per the least count; and the amount  $w$  of a row is the unit that arrives. Every reading Outside is a whole number of these, which is why the world above the board is quantized, a theorem of (A1), (A3) and the conversion (Theorem 3 of [21]; the velocity's quantum not read as such): physics above cannot take this from its own formulas, and the Inside gives it. The speed of light Outside is then

the arrival count over the Euclidean distance,  $c_D = N_t|D|/T_D$  by a row's age or by a pulse and its return, its anisotropy  $1/c_D^2 = 2.954, 2.971, 3.000$  on the heading, the face diagonal and the body diagonal and  $1/\sqrt{3}$  in the limit [20].

**Newton from the small step.** The same way, with five assumptions of the law as built and no (A2) (the crowd read as the age moment, the age wall at coefficient 1, the push, the flight blind, the reading a pulse and its return) and the spreading of  $K$  beams over a shell of  $N(r)$  Nodes, which is arithmetic and not a law, the fall, the equivalence principle, Kepler's ratio and the clock in a crowd come out Outside. The inverse square comes at the shell mean's rung and the equivalence exactly,  $M_A$  cancelling record by record. The register's D3, T and X are their confirmation, and the inverse square's decisive reading is not made [47]. Every result Outside is thus the image of one Inside formula under the conversion, and Table 3 lists the map row by row with the reading that sits on the Outside side.

**The road to each, and the check that nothing was derived from them.** The three roads are the one chain above, each from the Inside step, the six verbs, (A1) and, where it enters, (A2), with (A3), locality Outside (nothing Outside passes faster than a chain of clicks or jumps a Node), a definition of Outside and not a law [21]: to Lorentz, the boost from two legs at one Node per interval with  $k_{AB} = k_{BA}$ , the scale  $r$  free until (A2) fixes it at  $\sqrt{1-v^2}$  (the click frame's section 2); to Einstein, the rate  $1/\gamma$  as  $r$  read from two factors and  $E^2 = E_0^2 + c^2p^2$  as the line of the rate with  $W$  shown to second order (Theorems 1 and 2 and rows II.1, II.3 and II.6 of [21]); to the step above itself, the Outside step as the image of the Inside step, Einstein's step its most general form where  $r$  is supplied, its quantum a theorem of (A1), (A3) and the drive's one Link per self-creation, Newton's step its limit at  $v \ll c$  (section 3 of [21]). Einstein's, Lorentz's and the Outside step's formulas appear in these chains only as the thing compared with; no chain takes them as an input, and the Outside step is never a second formula set beside the Inside step. The word is “follows” or “recovered” where a known form is arrived at, “shown to second order” for  $W$  under (A1) and (A2), “declared” for  $W$  on the rows' dynamics and for the unit  $E_0 = mc^2$ ; “derived” stands nowhere. Table 6 (Appendix D) lists the check row by row, the inputs, the file and line where each chain starts and the word, read against the certification tables of the two sources and audited in their review.

**From an Inside formula to an Outside formula, family by family.** The same passage, written three times in full with the attributes of the row tuple of Section 2 (Node, direction, age, phase, emitter, content per unit, record, label, amount, multiplicity) named at each step, so that a reader sees which attribute each formula reads and where the conversion takes it Outside. *Light through two slits* (the family **light**,  $h = 1$ , the circle turns): the lamp puts one record on the board with two rows, one per opening, each with amount 1, multiplicity 2, the lamp's phase  $p_0$  at birth and its own direction on the fan; Inside, each row's phase advances by the declared turn per Link along its digital line, so at the pixel  $x$  the two arrive with the phases  $p_1(x)$  and  $p_2(x)$ , the difference  $\Delta(x)$  the path difference times the turn (the attributes read: direction, age, phase; the content plays no part,  $h = 1$ ); the merge adds them,  $f = x^{p_1} + x^{p_2}$  in  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ , and an equal pair at  $\Delta = N_\phi/2$  cancels; the click evaluates  $f$  at the root and squares it,  $R(x) = 1024(65536 + C[p_1]C[p_2] + S[p_1]S[p_2])$  (the formulas of Section 6) (the attributes read: amount, multiplicity, phase); the rung lays the pixels on the ladder and the birth wheel selects one, the record is deleted. Outside a detector holds only its counts per pixel: over  $N_\phi$  births the count at  $x$  is  $b_x - b_{x-1}$ , proportional to  $1 + \cos \Delta(x)$  within  $1/N_\phi$ , bright where the two ages differ by a whole turn, which is Young's law with the spacing set by the turn per Link; the reading is the bands' centres 23.5 pixels apart for the exact law's 23.3 (L2b, DETECTOR). *A massive body's clock and its fall* (the families **m** and **mass**,  $h = 0$ , no circle): a body carries its Node, its content  $M$ , its momentum **p**, an integer 3-vector, and its counts table,

each count an accumulator  $(s, r, d)$ ; Inside, its drive moves it at the pace  $|p_a|/(N_l N_w M + |p_a|)$  per axis where the axes' fires do not coincide, one Link per interval in all (the attributes read: content, momentum), its clock counts at the rate  $1/(1 + a_\tau n/d)$  with  $a_\tau$  the age moment of the crowd's rows at its Node (the crowd's attributes read: age, amount, direction; the body's: its counts), and the push reads the label flow  $\mathbf{a} = \sum \text{amount} \times \mathbf{u}_D$  of the arrivals into  $\mathbf{p}$  (the attributes read: amount, direction, content). The conversion is a lamp on the body and a detector at rest: the lamp's births are the body's own counts, the detector counts their arrivals with its own count, and Outside there are only the two counts and the births' Nodes. So  $1 + z$  is the ratio of the two counts. The ratio of two lamps' shifts at 3 and 6 Links is the clock's form at two distances, 1.907 for the pin 1.909 (series T, DETECTOR; read before the generic entry of 2026-09-22, not readable under the law as it stands, being re-read in the weak field). The fall is the birth's Node per count read at a line of detectors, the same Node on 138 of 139 births with the held mass four times (D3, DETECTOR), the equivalence read with no mass in the reading. *The pair* (the choosers  $\mathbf{sa}$  and  $\mathbf{sb}$ ; the pair's record carries the labels  $\{0, 1\}$  with integer weights on two arms): Inside, a setting  $a$  acts on an arm's label pair by the integer matrix  $U_a$  (the attributes read: label, amount, phase on each arm; the record's identity joins the arms), the joint weight is  $J(o_A, o_B) = \sum_\ell U_a[o_A][\ell] U_b[o_B][\ell]$  and the cell's offer  $R = J^2$ , Theorem 5; the conversion is the one gather of the record from both settings at the completion interval (P6), the rung over the four cells and the wheel's choice. Outside the two detectors hold four counts per setting pair:  $E_{N_\phi}(a, b) = 4c_{++}/N_\phi - 1$  and  $S(N_\phi)$  from them, Theorem 6, with each party's marginal  $N_\phi/2$  exactly; the reading is  $S = 2.75$  at  $N_\phi = 64$  and  $32/64$  in every bin (series L, DETECTOR). In each of the three, Outside holds counts and Nodes and nothing of the attributes themselves: the phase, the amount and the multiplicity of one record never pass singly, the content passes as a wall's scale, and what passes is what the click compares. Table 3 names, row by row, the family and the attributes each Inside formula reads.

Table 3: Inside formulas and Outside formulas, the conversion between them with its order, and the register's reading on the Outside side; each Inside cell opens with the family and the attributes the formula reads (the click frame's section 7 [47]; SHOWN is algebra under the named assumptions, MEASURED a run Inside read Outside).

| Inside (from the six operations)                                                                                                                                               | Outside (a detector reading)                                                                                              | The conversion; its order                                                         | The reading (kind)                                                                                                                     |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------|
| <i>a body</i> ( $\mu$ ; content, momentum): $W = E'_0 + 3\mathbf{p} \cdot \mathbf{p}$ on the record, $E'$ the whole root by comparisons; on the law no square, the wall linear | the energy a click reads, $\gamma$ , the rate $r = E'_0/E'$ (the identity) or 1 (the law)                                 | (A1), (A2), the Planck map; shown to order $m^2\beta^2$ ; the law's $r = 1$ exact | series S: the 64th self-creation at 70 and 124 (GAMEBOARD); $r$ not read                                                               |
| <i>a body</i> ( $\mathbf{m}$ ; content, momentum): the pace $ p_a /(N_l N_w M +  p_a )$ per axis; under the identity $p/E'$                                                    | the velocity: Nodes apart over counts apart between neighbouring detectors                                                | (A1); exact within $1/T_D$                                                        | D3's controls at their pace on every click (DETECTOR)                                                                                  |
| <i>light</i> (phase, age): the phase per age $n/d$ , $E = h_q n/d = h_A f$ , $\lambda = h_A/p$                                                                                 | the frequency a detector counts, $1 + z$ ; $k_{BA} = (1 + v)/r$ , $k_{AB} = r/(1 - v)$ , the round trip $(1 + v)/(1 - v)$ | (A1) and the count, rung 1                                                        | row 4b 0.2636 (FAIL); series S 0.3674 (MET in its domain); $k_{AB}$ not read                                                           |
| <i>a body's counts; the crowd's rows</i> (age, amount): the age wall, the crowd $a_\tau$ at coefficient 1, the rate $1/(1 + a_\tau n/d)$                                       | the clock rate in a crowd by a pulse and its return, the ratio of two lamps' $1 + z$                                      | (A1) and the count, rung 1                                                        | series T 1.907 for $1.909 \pm 0.05$ ; series X $k = 0.9089$ at $r = 4$ for the pin 0.9108 (DETECTOR; T before the entry of 2026-09-22) |
| <i>light</i> (direction, age): the flight table $T_D$ , the pace $N_l D /T_D$                                                                                                  | the arrival count at a detector $L$ Links away; $c$ one Link per the least count                                          | (A1); rung 1 in the count, rung 2 in the isotropy                                 | series Q 290 of 290; series S 369, 345 (DETECTOR)                                                                                      |
| <i>light, the pair</i> (phase, amount, multiplicity, label): the click's bilinear form $\mathbf{f}^T \mathbf{G} \mathbf{f}$                                                    | Born's form, the count of clicks over many records                                                                        | (A2) at the click; rung 1                                                         | the two-slit fringes, Malus (DETECTOR); the amount and the phase of one record never pass singly                                       |

| Inside                                                                                                                                                      | Outside                                                                                                 | The conversion                                                        | The reading                                                                             |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| <i>the crowd's rows</i> (amount, age, direction): the spreading, the presence $q\tau_L/(4\pi r^2)$ and the age moment $q\tau_L/(4\pi cr)$ in the shell mean | the clock's field, the ratio of two shifts 2.00; a shell's flat interior                                | rung 2, the shell mean; the ripple MEASURED ONLY                      | series T 1.907; series X 1.0029 for 1.0039 (DETECTOR; T before the entry of 2026-09-22) |
| <i>a body; the arrivals</i> (content, momentum; amount, direction): the push $-M_A \mathbf{a}$ and the drive                                                | the fall, $a = -GM_B/r^2$ , $G = K\eta F/(4\pi N_w)$ ( $F$ the fan's Manhattan factor); the equivalence | (A1) and the count; the equivalence rung 1, the inverse square rung 2 | D3: 138 of 139 common birth ticks (DETECTOR); the inverse square's reading not made     |
| <i>a body</i> (Node, momentum): the circular momentum under the push, $v^2/r = GM_B/r^2$                                                                    | Kepler's period, $T(24)/T(12) = 2$ on the plane                                                         | rung 2, $v \ll c$                                                     | D3: 1.997 in [1.82, 2.18] (DETECTOR)                                                    |

**What the Inside step gives Outside that the continuum cannot state.** Formulas of the six verbs taken Outside by the conversion, each with the click that reads it and its kind; none of them is a formula of the continuum; the full lists, each candidate under the three tests with its reading, are [25] for quanta and bodies and [26] for light. (i) The quantum of the Outside step, stated above (one Link, two counts,  $1/k$  with its quantum  $1/(k(k+1))$ ; Theorem 3 of [21]), read as a least step in series S's face clicks 369 and 345 within two of the flight table's and D3's 138 of 139 births at the same Node (DETECTOR); the velocity's quantum and the pulse's two counts not read as such. (ii) The pair's sum at the CHSH labels is the exact rational the rung gives,  $|S(N_\phi) - 2\sqrt{2}| \leq 8/N_\phi + 0.0444$ , above the bound at  $N_\phi = 16$  and 32 ( $S = 3$ ), the plateau 181/64 from 512 through 8192 below it by  $3.02 \times 10^{-4}$  and 5793/2048 above at 16384 (Theorem 6; the frame's section 10); measured at seven  $N_\phi$  (Figure 6, DETECTOR). (iii) The perihelion: the drive's pace departs from  $p/m_i$  at first order per axis, so a bound orbit's apsides move per revolution by an angle of order  $\beta$  with the lattice's symmetry, where Einstein has  $6\pi\beta^2$  isotropic, larger than nature's by  $1/(6\pi\beta)$ , about 330 times at Mercury's pace if its coefficient is of order one, and of the wrong symmetry: FAIL on the law, pinned in order and symmetry, the coefficient NOT MADE; under the identity  $\pi\beta^2$ , one sixth; the deciding reading, the apsides of a lamp's orbit at two orientations to the axes, NOT MADE, its pin written before any run, an apsidal motion of order  $\beta$  that changes with the orientation and not with the mass held ([21], II.12). (iv) The anisotropy of  $c$  by direction, stated above (series Q, 290 of 290, DETECTOR; [20]). (v) The equivalence's constant: the clock in a crowd and the accelerated detector read one shift,  $\delta k = (nS/d)(gY/c^2)$ , Einstein's  $gY/c^2$  exactly when  $nS = d$  (times the grain's  $\tau_{LC} = 0.993$  on a heading, 1 in the limit of every direction), a condition on an Inside declaration and not a result; the register's clock and crowd worlds declare  $nS/d = 16$  (GAMEBOARD, the world files), so the condition is not met by them, the form's readings are series T's 1.907 and series X's  $k$  (DETECTOR; T read before the generic entry of 2026-09-22, not readable under the law as it stands, being re-read in the weak field), and the constant's reading is NOT MADE ([21], II.10). (vi) Light's bending: on the law of this paper, the runtime before the generic entry, 0, the flight blind to the crowd (series K, 0.000 pixel, DETECTOR; FAIL against 1.75 arcseconds); under the key `optical`, the flight in the age wall's set with a declared  $c_f$ , the delay  $c_f(nS/d)(GM/c^3) \ln(4r_1r_2/b^2)$  and the bending  $\alpha = 2c_f(nS/d) GM/(c^2b)$ , Einstein's at  $nS = d$  and  $c_f = 2$ , NOT COMPARED, the route chosen ([21], II.11). The chain, algebra first and the run after (Section 5; Table 2, row 13): the step algebra's ring on the law as built,  $C_{\text{ring}} = 5.12$  at  $b = 6$ , the  $M/b$  form recovered and 4 not reached (COMPUTATION [34]); under the one constant 3.65 to 3.92 against 4,  $2c_f$  within two stated factors (COMPUTATION [35]); the run of 2026-09-22 under `flow_link`, the  $\gamma = 0$  ring at  $b = 6$  inside its four pins (DETECTOR) and, on the split ladder, every ring read inside its pins (the deciding pin 0.731 against  $0.731 \pm 0.025$ , DETECTOR), the constant through the stated factors outside the one-grain pin by 2.40, 1.07, 1.40

grains at  $\gamma = 1$ : DOES NOT CLOSE within the grain [38]. The conjecture, labelled so: on the one constant the deflection coefficient is  $2(1 + \gamma)$ ,  $\gamma$  the declared input, conjectured from the algebra and not read (Section 5). (vii) Dark energy's shape: the diagram of redshift against flight time read at a detector carries, beside the throw's Doppler  $1 + z = 1/(1 - x)$ ,  $x = H\tau$  (Milne's coasting form,  $q = 0$  exactly), one factor  $g(\tau) = (1 + k_A(\tau))/(1 + k_B)$ , the emitters' clock stretch by their crowd over the detector's own; with  $g = 1 + g_1x + \dots$  the deceleration parameter a reader fits is  $q_{\text{eff}} = -2g_1/(1 + g_1)$ , an apparent acceleration for every  $g_1 > 0$  with nothing accelerating on the board, a property of the conversion and not of any dynamics ([24]; rung 2, the throw's continuum). The register read that shape (series G, the accelerating form the nearest of three in all six pushing windows, DETECTOR) and the coasting throw's  $q = -0.108$  in  $0 \pm 0.25$  (G2, MET; FAIL against nature's  $-0.53$ , row 3). What the law does not give, stated in the same font: the size,  $g_1 = 0.36$  for  $-0.53$ , an input of the crowd (a declared pair, four orders above the known wells); the sign from its own accumulation of the crowd,  $q_{\text{eff}} = 2k_0(3 - k_0)/(1 - k_0)^2 > 0$ , the older light born in a thinner crowd; and the brightness of a standard lamp, which reads  $q_{\text{eff}} = 2/(1 + g_1)$ , a deceleration, one factor of  $1 + z$  short of the expanding form because a click carries the content of its birth (row 11a, FAIL). The model has no cosmological constant and needs none as a rule; it states the supernova diagram as a failure, not a resolution. (viii) The atom. What the algebra gives first, on the law's integers (COMPUTATION [36]): the closing radius is an integer congruence of the action row,  $\Delta A_a = 0 \bmod h$  on each axis with the three quotients summing to  $jN$ ; two closing radii stand as the square of two counts a detector reads,  $a_j/a_i = (j/i)^2$  in the shell mean, checked by  $T_j/T_i = (j/i)^3$ , no root and no float; and a theorem of the click says that from a closed circle one click reaches only the loops with  $2i^2 \geq j^2$ , while a release of content unbinds, so the transition Bohr reads as one line is not an operation of the law: NEEDS A NEW RULE, named under its own identity (atom-level-v1, the release at a closure of the difference of two closures, the virial form, [37]). Stated as a conjecture, labelled so and not read [36, 37]: every closed loop with  $j$  circles per return has, in the shell mean,  $a_j = j^2 h^2 / (4\pi^2 \kappa m)$ , so  $a_j/a_i = (j/i)^2$  and  $T_j/T_i = (j/i)^3$ ; the loop's constant  $-\kappa/(2a_j)$  on the virial form,  $E_j = -\Delta A/(2NT_j) = -jh/(2T_j)$ , is proportional to  $1/j^2$  in the limit, so two lines would stand in the ratios  $(1/i^2 - 1/j^2)$ , 27/20 for Balmer's  $\beta$  over  $\alpha$ , free of every declared constant; the conjecture is that the law's closed loops form Bohr's ladder in the shell mean's limit, conjectured from the algebra, not recovered. What is read: the  $r = 12$  loop's staying under the centred step and its tightening (DETECTOR); no level and no line. The atom enters the paper by the algebraic formula alone, and the paper says plainly why: its levels cannot be read on any lattice that can be run. In nature the electron's orbit radius over the nucleus's radius is about sixty thousand (the Bohr radius  $5.3 \times 10^{-11}$  m over the proton's  $0.84 \times 10^{-15}$  m), while the lattice worlds put the electron at  $r = 3$  to 12 Nodes from the proton, where the fan's grain (about 17 degrees per shell at  $r = 3$ , the flux 17 percent below the inverse square) is the named limit; a lattice of order fifty thousand on a side would be needed and is not run ([6, record 997, pending merge]). So the ladder is the algebra's convergence to Balmer's form in the limit: computed from the algebra, not read on the board; a lattice able to hold the scale is not run; not shown in nature. No run of the levels is cited as a reading, and row 6 stays NOT YET on the line. The run after it: the hydrogen loop at  $r = 12$  (series H's world, the electron's held content on the proton's push) escapes under the drive as built at 3407 intervals (DETECTOR), the law's own FAIL, beside which the key's PASS below stands. The cause, read from the baseline's records by algebra [30]: the count primitive fires a body's step when its accumulated motion reaches a whole wall, so the body's Node lags its accumulated motion by half a Link in the mean on each axis, toward the motion's sign, and the push read at the lagging Node has a part along the motion,  $(S_1(\mathbf{v})/|\mathbf{v}|)/(2r)$ , about 5 percent of the radial at  $r = 12$ , which pumps the loop's angular momentum and energy (up 30 and 45 percent over the first three quarters of a turn) until the loop opens. The centred step (`centred_step`, a

declared start of the drive under its key, off by default, every registered world byte identical without it: the count of a body's step the nearest whole number of its accumulator in units of the wall, the lag zero in the mean) is a hypothesis under its own name until the author makes it the law; under it, in the one run of 2026-09-22 [29], the loop stays: no escape face click in 7500 intervals, 23 quarter crossings (C1 PASS, DETECTOR), tightening from 12 to 7 Links at two crossings over five turns, the period falling from 1431 to 1040 (C2, C3 FAIL as declared, DETECTOR), no pin moved after the run; no line of Bohr's is read, and none is computable by a run of the law as built, no rule of the six quantising a body's loop (Table 2, row 6). The families of the register, each with its declared integers, its algebraic object and the click that reads it, are Table 5.

**What is proved.** Derived Inside, exact on the GameBoard, no experiment needed: the group of the six Ports and its 24 rotations; the books' conservation and the lattice form of the continuity equation; Gauss's law of a free family's flux under its world condition; the split's isometry and the injectivity of the interval on the amplitudes for a fixed event history; the pace of every direction, isotropic within  $1/T_D$ ; the click's weight a positive quadratic form of power 2; the exact marginals of a pair; the CHSH sum as an exact function of the grain and the tables; Planck's and de Broglie's relations as identities under the declared dictionary. Recovered in the limit of the grain, with a rate: Born's rule to  $1/N_\phi$  per cell and the cumulative rungs to  $1/(2N_\phi)$  with the exact tables (to the tables' rounding with the declared ones, Section 6), Tsirelson's bound to  $8/N_\phi$  plus the tables' 0.0444 ( $2\sqrt{2}$  in the joint limit of  $N_\phi$  and  $N_t$ ), Young's spacing, the attained value of  $c$ , the continuity equation's differential form. Recovered under an average that covers the shell, with the condition in the claim: Newton's and Coulomb's inverse square, the retarded potential and Poisson's equation, the clock's  $1/r$  form. Each names the known form the algebra recovers (Gauss's, Newton's, Coulomb's, Born's, Tsirelson's, Planck's, Einstein's); the paper's own are the integer road, the finite- $N_\phi$  values (Born's bound  $1/N_\phi$ , the plateau 181/64, the fixed-table limit), the quadratic read-out characterization on  $\mathbb{Z}_{N_\phi}$  (Gleason-like), the measurable  $1 - v^2$  and the entropy identity of the click (an identity of the wheel); the value  $1/\sqrt{3}$ , Tsirelson's limit and the dictionary's identities claim no novelty. Conjectured from the algebra and not read, labelled so wherever they appear (Sections 5 and 9, Table 1): the bending's coefficient  $2(1 + \gamma)$  on the one constant, and the atom's ladder  $a_j/a_i = (j/i)^2$ ,  $E_j \propto 1/j^2$ , with Balmer's 27/20 in the shell mean's limit; the ring's run reads its rule inside its pins and its constant outside the one-grain pin, and the ladder is computed from the algebra, not read on the board. After a detector every reading sits inside an expectation written before the run; against nature three readings pass (1a, 2b, 9); twelve fail in the table, eleven in a registered run (1b, 1c, 2a, 3, 4b, 6, 7b, 8a, 8b, 13, 14) and one a declared input refuted (8c), with four more failing by a pin whose run is not made and not in the table (4a, 5b, 11a, 11c); two are bounds (5a, 7a); one is not compared (2c); one is history (12, the lattice pin met under the law before the entry, not a pass against nature); the hypotheses beside rows 6, 13 and 14 change no verdict of the law (Table 2). What no rule of the law forces, with the check that decides each. Newton's inverse square and Poisson's law closed after a detector ( $G$  not read; D3): decided by a closed orbit's period at two radii under the directional drive (form B, built off by default, series Y), the escape's count at a face detector and the clock's form at more distances. The Lorentz factor, the law meeting (A1) and not (A2): decided by the ratio of the two one-way Doppler factors through records,  $1 - v^2$  under the law and 1 under Lorentz, and by the split of a row between staying and hopping (A2), a hypothesis not opened. The bending of light, the flight blind to the crowd: decided by a rule under which a row's wall reads the crowd's age moment (a hypothesis, its coefficient an input) against nature's 1.75 arcseconds. The masses' values, every rule linear in the content and every equal split a fixed point: decided by a nonlinear

closure on the amounts under its own identity, the family table an input until then. The click's harmonic constants and the circle as position (a hypothesis, Section 6): decided by the least-rank click, an axiom of the apparatus pinned by the two-slit period and Malus at 22.5 degrees, and by the single opening against its pin (row 10). And a CHSH measurement at  $10^{-4}$ , deciding 181/64 against  $2\sqrt{2}$ .

**The limits, and what does not return.** The content and  $\sum w^2/\mathfrak{m}$  are conserved exactly, the offered norm only within the tables' rounding. The covariant readings [6, record 270] read one Inside formula Outside on one axis, series S against its pins: a body's record carries the exact square of its energy,

$$\boxed{W = E_0'^2 + 3\mathbf{p} \cdot \mathbf{p}, \quad E_0' = E_0/c^2 = N_l N_w M, \quad c^2 = 1/3} \quad (14)$$

compared and never rooted, with  $E'$  the largest integer whose square is at most  $W$  and every count of the body gated by  $E_0/E'$ ; multiplied by  $c^4$  it is the energy-momentum relation of special relativity (series S, Table 3, inside their pins). Eq. (14) is a declared identity of the record's representation in the law as built, not derived from the rows' dynamics; under (A2), an extension the law does not meet,  $W$  is shown to second order (Section 9). The hypotheses beside the law are on the tree [11].

**What the law names and has not computed.** Each item below is a computation the law's rules define, on a declared world with a declared click, that this paper has not made: the register holds no reading for it, nothing is claimed for it, and its verdict is NOT YET, NOT COMPARED or absent until a run is made and read by kind. (i) The lines of the atom: not a run not yet made but a rule the law lacks, no rule of the six quantising a body's loop and one click reaching only the loops with  $2i^2 \geq j^2$  (Table 2, row 6; [36]); Bohr's 27/20 waits on that rule, under its own identity; the ladder itself,  $a_j/a_i = (j/i)^2$  and  $E_j \propto 1/j^2$  with 27/20, is conjectured from the algebra, not computed (item (viii) of the list above). (ii) Third-order interference, Sorkin's  $\kappa$  on a three-opening world (row 2c; no such world is registered). (iii) The single opening's spread against its pin (row 10), and the rows pinned without their run (4a, 5b, 11). (iv) The quarks and the nucleon: the quark families are in the catalog with their rational charges and their content per record (Table 5, u), and no detector reading of a quarks world is registered; the nucleon as three measured events bound by the strong column, its read mass against the sum of its parts, is the reading that would test them [6, record 248]; colour, a  $\mathbb{Z}_3$  label that passes the three tests and does nothing beside a colour force between rows that fails the vector and the local test, is designed under its own identity and not built [6, record 270]. (v) The unseen content: on the law's own rules the field at a galaxy's edge is Newton's in the shell mean and the detector adds nothing there; the one route to a flat rotation curve without unseen content, a short periodic dimension ([11], entry 6), fails nature's Tully-Fisher slope on a host map ([41], no run); what the law admits is unseen content as an input family, as nature's model does; no rotation curve is read after a detector and no disc under a declared unseen family is run. (vi) The size of the crowd's stretch,  $g_1 = 0.36$ , that the supernova diagram needs, from a declared content over a flight (an input in the dark energy item of the list above; the crowd's own accumulation gives the wrong sign), and the periodic GameBoard against the open one on the measured numbers, declared and not decided [6, record 270]. (vii) The detector's constants  $c_j$  of the click's harmonics, not reached by the algebra (Section 6). (viii) The inverse square in space, pending its run (row 14); the bending's full deflection, its coefficient  $2(1 + \gamma)$  on the one constant conjectured from the algebra; the ring's rule read inside its pins and the constant 4 not closing within the grain (row 13). Apart from these stand the numbers for which the law has no formula

and this paper offers none: the hierarchy  $\rho_p$ , the constant  $h = h_q N_\phi$ , the families' content per unit and the grain  $N_\phi$ , inputs by the dictionary (Table 1).

**Positioning.** The click's assignment of the second outcome from the joint weights is a non-local step on the labels, its interval the last arm's (as after Theorem 5), forced by Bell's theorem because the marginals are exact and  $S > 2$  [49, 50]; no-signalling of the counts is Theorem 5 [51], and the order of the outcomes signals under the sequential wheel (Section 7; Table 2, row 1c, FAIL after a detector). The click's finite kinematics is Weyl's and Schwinger's on  $\mathbb{Z}_{N_\phi}$  [71, 72]; the transport a lattice gas's exact integer dynamics [52, 53], free of dispersion where the lattice automata of [76, 77, 78] disperse; the uniform  $u$  does the work of quantum equilibrium [79, 80]; the model is  $\psi$ -ontic [81, 82]; Theorem 6 puts  $S(N_\phi)$  on both sides of Tsirelson's bound [83, 84]. The earlier testbed [1] is history.

## Declarations

**Funding.** None.

**Competing interests.** The author declares no competing interests.

**Ethics approval.** Not applicable.

**Data and code availability.** The simulator, every world file, every registered run's record, the check scripts with their outputs, the figures' scripts and the derivation records cited in this paper are archived at Zenodo [88] (concept DOI 10.5281/zenodo.22738746); the commit of the archived version is named in Appendix C, and every number of the paper has one row in the ledger of that archive (`paper/general_formula/NUMBERS.md`) naming its kind and its source.

**Author contributions.** Sole author: the model, its rules, the worlds and the experiments, every reading and its verdict, and the decisions recorded in the archived log.

**Use of artificial intelligence.** A large language model (Claude, Anthropic, through Claude Code, 2026) was used as a tool under the author's direction for code, computation and the drafting of the text; the author reviewed every statement and is responsible for every claim. The author's statement: I laid down every postulate and hypothesis of the model myself and brought in nothing external; the forms of Newton, Kepler, Einstein, Lorentz, Bohr and Balmer that this paper cites are the things compared with, not inputs to the model.

## References

- [1] A. Gonen, A finite-integer causal-GameBoard testbed: which resources reproduce interference, causal collapse and Bell correlations on one GameBoard (2026), arXiv preprint.
- [2] The Beam Law and its implementation notes, `docs/BEAM_LAW.md` of the archived code [88].
- [3] The design of `amplitude-v1` and its check outputs, `docs/designs/amplitude-v1/` of the archived code [88].
- [4] The experiments register, `docs/EXPERIMENTS.md` of the archived code [88]; consolidated on the branch `register-paper-sources` (its head b12dd253, pending merge).

- [5] `paper/general_formula/checks/s_of_n.py`, `s_powers_of_two.py` (the pair's  $S$  and its four correlations at the powers of two from 512 to 32768, the closed form on the fixed correlations to  $2^{20}$  with the distances from [85], with the tables fixed), `which_path_window.py`, `tables_norm.py`, `information_transfer.py`, `limits.py` and `light_speed.py` with their outputs, `figures/summary.json` and `NUMBERS.md`, in the archived code [88].
- [6] The dated working log of the model, cited by the number of its record, `docs/LOG_2026-09-20.md` of the archived code [88].
- [7] C. E. Shannon, Bell Syst. Tech. J. 27, 379 and 623 (1948).
- [8] The one symbol table of the law without clashes with physics' letters, the derivation mathematician's proposal `docs/designs/notation/NOTATION.md` of the archived code [88].
- [9] The replications register, `docs/REPLICATIONS.md` of the archived code [88].
- [10] The families in the paper, the mathematician's audit of 2026-09-21, `docs/designs/paper_families/FAMILY_TABLE_AUDIT.md` with its host script's output `family_table.out`, of the archived code [88] at the commit named in Reproducibility.
- [11] Hypotheses under test, entry 27, the conditional derivations as declared hypotheses, `docs/HYPOTHESES.md` of the archived code [88].
- [12] Canonical simulation terminology, `docs/TERMINOLOGY.md` of the archived code [88].
- [13] The clock's word, the presence or the age moment: the physicist's read, `docs/designs/clock_age/NOTE.md` of the archived code [88].
- [14] N. Ashby, Living Rev. Relativity 6, 1 (2003).
- [15] R. V. Pound and G. A. Rebka, Phys. Rev. Lett. 4, 337 (1960).
- [16] The law as one vector algebra: every rule in force as a vector operation, with its integer form beside it, `docs/designs/vector_form/LAW.md` of the archived code [88].
- [17] The engine's document, `docs/ENGINE.md` of the archived code [88].
- [18] The project's decisions and dictionary, `docs/HIGHLIGHTS.md` of the archived code [88].
- [19] The derivations of the beam law, `docs/DERIVATIONS_BEAM.md` of the archived code [88].
- [20] Light Outside from the board: every equation of light Inside carried Outside by its own transformation, `docs/designs/light_outside/DERIVATION.md` of the archived code [88], on `main` at a62441fb.
- [21] Einstein Outside: the formulas of the special and the general theory brought from inside the GameBoard to above it, `docs/designs/einstein_outside/DERIVATION.md` of the archived code [88], on `main` at e8432e6c .
- [22] The transition from ordinary physics to modern algebra: a dated, cited history, `docs/designs/algebra_transition/HISTORY.md` of the archived code [88], on `main` at 412f5c61 .
- [23] The click frame, part 2 ("no square is declared anywhere in the chain"), `docs/designs/click_frame/DERIVATION.md` of the archived code [88] on `main` at af051aaf , one file with the click frame.
- [24] Dark energy Outside: what the passage from Inside to Outside puts into the Hubble diagram, and what it does not, `docs/designs/light_outside/DARK_ENERGY.md` of the archived code [88] on `main` at c8ac2ffb.
- [25] New formulas of quanta and bodies: candidates that the Inside step gives Outside and the continuum cannot state, `docs/designs/new_formulas/QUANTA.md` of the archived code [88] on `main` at 477daald .
- [26] New formulas of light: what the Inside step gives Outside that the continuum cannot state, `docs/designs/new_formulas/LIGHT.md` of the archived code [88] on `main` at 7370d524 .
- [27] T. Toffoli and N. Margolus, Cellular Automata Machines (MIT Press, Cambridge, 1987).
- [28] The order channel: the run and its readings, `docs/designs/order_channel/RUN.md` and `PINS.md` of the archived code [88], on `main` at 804a8da0 .
- [29] The one run of hydrogen at  $r = 12$  under the centred step, `docs/designs/atom_give/RUN_CENTRED.md` and `CENTRED_STEP.md` of the archived code [88], on `main` at a4794cc6 .
- [30] The cause of the widening at  $r = 12$ , read from the baseline's records by algebra, `docs/designs/atom_give/CAUSE.md` of the archived code [88], on `main` at 8ee973d6 .
- [31] The Gleason bound: the tables' rounding against hypothesis (a), the weight of a mixture the register cannot exclude, and the table click's zeros, `docs/designs/gleason_bound/BOUND.md` of the archived code [88], on `main` at e6809713 .

- [32] The generic entry of the bending: the flight in the age wall's set at  $1 + \gamma$  for every world,  $\gamma$  an input, the one form of the last Link's time, the square ladder in place of the root, of the archived code [88], on `main` at 5b855791 (the branch head it carried 3760754c).
- [33] The orbit's inward push read on the fan: a body at rest on a fan reads the fan's Manhattan mean  $S_1/|\mathbf{D}| = 1.287$  shell means, `docs/designs/orbit_read/NOTE.md` of the archived code [88], on `main` at 75efe7b5 (host arithmetic, no run).
- [34] The step algebra of light bending: the walk of a light row past a held mass by the law's integer steps, read at a screen of Nodes and at a ring of starts, `docs/designs/light_bending/STEP_ALGEBRA.md` of the archived code [88], on `main` at 23ef9ce9 .
- [35] The algebraic check of flow-link-v1: the step algebra beside a mass, the ring of starts and Newton's rows under the weighted flow, `docs/designs/flow_weight/ALGEBRA.md` of the archived code [88], on `main` at d9b6694a.
- [36] The atom's algebra: two closing radii as a ratio of counts at a detector, and the click that moves a row between them, `docs/designs/atom_algebra/ALGEBRA.md` of the archived code [88], on `main` at 643b7ae7 .
- [37] atom-level-v1: the release at a closure of the difference of two closures, the ladder of levels in the law's integers and what the rule gives against Balmer's ratio, `docs/designs/atom_levels/LEVELS.md` of the archived code [88], on `main` at d8afc87c (the merge of `atom-levels`; `LEVELS.md` unchanged from the branch head 83b1bf2e).
- [38] The ring worlds of flow-link-v1: the pins before the run, the one run of 2026-09-22 and the register's refusal, `examples/events/flow_link/README.md` and `expectations.json` of the archived code [88], on `main` at 23b0e630 (the merge of `flow-link-build`; the run on the split ladder read at the branch head b094c53b, its files unchanged by the merge).
- [39] Series D3 under the one constant: the four worlds, the pins by the generator before any run and the one run of 2026-09-22, `examples/events/orbit_lamp/README.md`, `expectations_flow.json` and `run_flow.out` of the archived code [88], on `main` at 7682c57d (the reading's files as at the branch head f44a69e5).
- [40] Why Newton's rows fell: the diagnosis of series D3's ratio 1.677 against the pin 2.00, the probes' escapes read from the registered events, `docs/designs/newton_diagnosis/DIAGNOSIS.md` of the archived code [88], on `main` at 5ce4eb06 (the merge of `newton-diagnosis`); in its own words, nothing there is compared with nature.
- [41] The dark sector against the law's own rules: no dark energy and no dark matter, does it close the picture, `docs/designs/far_lamp/DARK_SECTOR.md` of the archived code [88], on `main` at aa243019 (a host map, no run).
- [42] F. W. Dyson, A. S. Eddington and C. Davidson, Phil. Trans. R. Soc. A 220, 291 (1920).
- [43] A. Kramida, Yu. Ralchenko, J. Reader and the NIST ASD Team, NIST Atomic Spectra Database, National Institute of Standards and Technology; the ratio to verify at the fifth digit.
- [44] The criteria of the confrontation table, row by row, with the re-runs at head, `docs/designs/paper_criteria/CRITERIA.md` of the archived code [88], on `main` at f5fe8004 .
- [45] V. Jacques, E. Wu, T. Toury, F. Treussart, A. Aspect, P. Grangier and J.-F. Roch, Eur. Phys. J. D 35, 561 (2005); the visibility figure to be verified against the source.
- [46] The ring mean and the orbit's mean over one turn: the proof and the bound, `docs/designs/ring_mean/PROOF.md` of the archived code [88], on `main` at ab96e7e8 .
- [47] Lorentz from the clicks: the click frame, its section 0 the click theorem, `docs/designs/click_frame/DERIVATION.md` of the archived code [88], on `main` at 70e9781a (the file identical to its head 1dd81fef).
- [48] The confrontation register, `docs/NATURE.md` of the archived code [88].
- [49] J. S. Bell, Physics 1, 195 (1964).
- [50] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, Phys. Rev. Lett. 23, 880 (1969).
- [51] G. C. Ghirardi, A. Rimini and T. Weber, Lett. Nuovo Cimento 27, 293 (1980).
- [52] J. Hardy, O. de Pazzis and Y. Pomeau, Phys. Rev. A 13, 1949 (1976).
- [53] U. Frisch, B. Hasslacher and Y. Pomeau, Phys. Rev. Lett. 56, 1505 (1986).
- [54] R. Courant, K. Friedrichs and H. Lewy, Math. Ann. 100, 32 (1928).
- [55] Y. H. Qian, D. d'Humières and P. Lallemand, Europhys. Lett. 17, 479 (1992).
- [56] M. Nagel, S. R. Parker, E. V. Kovalchuk, P. L. Stanwix, J. G. Hartnett, E. N. Ivanov, A. Peters and M. E. Tobar, Nature Commun. 6, 8174 (2015).

- [57] B. Botermann et al., Phys. Rev. Lett. 113, 120405 (2014).
- [58] M. Wang, W. J. Huang, F. G. Kondev, G. Audi and S. Naimi, Chin. Phys. C 45, 030003 (2021).
- [59] P. Delva et al., Phys. Rev. Lett. 121, 231101 (2018).
- [60] U. Sinha, C. Couteau, T. Jennewein, R. Laflamme and G. Weihs, Science 329, 418 (2010).
- [61] N. Aghanim et al. (Planck Collaboration), Astron. Astrophys. 641, A6 (2020).
- [62] A. G. Riess et al., Astron. J. 116, 1009 (1998).
- [63] S. Perlmutter et al., Astrophys. J. 517, 565 (1999).
- [64] S. Navas et al. (Particle Data Group), Phys. Rev. D 110, 030001 (2024).
- [65] M. Aker et al. (KATRIN Collaboration), Nature Phys. 18, 160 (2022).
- [66] F. M. Gonzalez et al., Phys. Rev. Lett. 127, 162501 (2021).
- [67] M. N. Huxley, Proc. London Math. Soc. 87, 591 (2003).
- [68] A. M. Gleason, J. Math. Mech. 6, 885 (1957).
- [69] D. L. Donoho and P. B. Stark, SIAM J. Appl. Math. 49, 906 (1989).
- [70] E. H. Kennard, Z. Phys. 44, 326 (1927).
- [71] H. Weyl, The Theory of Groups and Quantum Mechanics, translated by H. P. Robertson (Methuen, London, 1931; Dover, New York, 1950), chapter IV, section 14.
- [72] J. Schwinger, Proc. Natl. Acad. Sci. USA 46, 570 (1960).
- [73] H. P. Robertson, Phys. Rev. 34, 163 (1929).
- [74] H. Maassen and J. B. M. Uffink, Phys. Rev. Lett. 60, 1103 (1988).
- [75] P. Jordan and J. von Neumann, Ann. Math. 36, 719 (1935).
- [76] I. Bialynicki-Birula, Phys. Rev. D 49, 6920 (1994).
- [77] D. A. Meyer, J. Stat. Phys. 85, 551 (1996).
- [78] P. Arrighi, Natural Computing 18, 885 (2019).
- [79] D. Bohm, Phys. Rev. 85, 166 and 180 (1952).
- [80] D. Dürr, S. Goldstein and N. Zanghì, J. Stat. Phys. 67, 843 (1992).
- [81] R. W. Spekkens, Phys. Rev. A 75, 032110 (2007).
- [82] N. Harrigan and R. W. Spekkens, Found. Phys. 40, 125 (2010).
- [83] B. S. Cirel'son, Lett. Math. Phys. 4, 93 (1980).
- [84] S. Popescu and D. Rohrlich, Found. Phys. 24, 379 (1994).
- [85] H. S. Poh, S. K. Joshi, A. Cerè, A. Cabello and C. Kurtsiefer, Phys. Rev. Lett. 115, 180408 (2015).
- [86] B. Hensen et al., Nature 526, 682 (2015).
- [87] P. Grangier, G. Roger and A. Aspect, Europhys. Lett. 1, 173 (1986).
- [88] A. Gonen, Universe24: a discrete simulator of local physical laws (Reality Theory), Zenodo (2026), doi:10.5281/zenodo.22738746 (concept DOI, resolving to the latest version).

## A Auxiliary proofs

**Proof of Theorem 1.** A map of the six Ports that keeps opposite Ports opposite is a signed permutation of the three axes,  $3!2^3 = 48$  of them, the hyperoctahedral group  $B_3$ ; the determinant is a homomorphism onto  $\{+1, -1\}$ , so its kernel, the rotations, has index 2 and order 24, and it acts faithfully on the cube's four body diagonals, so it is the symmetric group on them; a pseudoscalar column goes to  $\det(g)$  times itself by definition.

**Proof of Theorem 2.**  $\sum_i (wa_i)^2 / (\mathfrak{m}A) = w^2 / \mathfrak{m}$ . The conjugate transpose maps the outputs  $(wa_i, \mathfrak{m}A, p + t_i)$  to  $(Aw, A^2 \mathfrak{m}, p) \sim (w, \mathfrak{m}, p)$ ; for two inputs at one splitter the cross terms carry phases differing by  $N_\phi/2$  and cancel in the normal form, which the check with the pairs (20, 21), (3, 4, 5) and (119, 120, 169) confirms integer by integer [3], and the engine’s merge reproduces on its own rows [2]. That cancellation is the joint condition  $B^H B = 2I$  of the balanced splitter’s table  $\begin{pmatrix} 1 & x^{N_\phi/4} \\ x^{N_\phi/4} & 1 \end{pmatrix}$ , and the condition is needed: two inputs declared (1, 1) with the turns (0, 0) onto the same two outputs have the table  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ , which sends (1, −1) to 0 and merges the inputs (1, 3) and (2, 2) to the same (4, 4) (the exact witness of the independent read of 2026-09-22); each column keeps its own  $w^2 / \mathfrak{m}$  there and the joint map does not, and Rule S as a generic rule admits such a table; no world of this paper declares one. For  $U_s$  the diagonal of  $U_s^T U_s$  is  $C'^2 + S'^2$  and the off-diagonal  $C' S' - S' C' = 0$  in integers.

**Proof of Theorem 3.**  $F$  maps a basis element of  $Q$  to a basis element, the Node moved by an offset that is a function of  $D$  and of the age, the history’s, so with the history fixed it is injective on the basis;  $S$  absorbs the arriving row and re-emits its weight, multiplicity and phase on its outputs at age 0, injective on  $Q$  by Theorem 2 under the joint condition  $B^H B = AI$ ;  $R$  is injective because the turn is a bijection and  $U_s^T U_s = n_s I$ ;  $M$  is the identity on the module; a composition of injective linear maps is injective. The collision, a permutation of the amount-1 rows at a Node as a function of the whole set present, is a bijection of states and not a linear map, which is why it is excluded from the linear statement; the engine’s inverse interval is the witness on worlds without a measured event [2].

## B Technical details

**The tables and their rounding.** interval  $t \in \mathbb{Z}$ . The circle is  $\mathbb{Z}_{N_\phi}$  with  $4 \mid N_\phi$ . The tables are  $C_{N_\phi}[p] = \text{round}(256 \cos(2\pi p / N_\phi))$  and  $S_{N_\phi}[p] = C_{N_\phi}[p - N_\phi/4]$ , so  $C_{N_\phi}[p + N_\phi/2] = -C_{N_\phi}[p]$  and  $S_{N_\phi}[p + N_\phi/4] = C_{N_\phi}[p]$  exactly; the norm  $C_{N_\phi}^2 + S_{N_\phi}^2$  lies in  $[65536 - 351, 65536 + 361]$  for every power of two  $N_\phi \leq 65536$  [5]. The normalisation by  $C_K$  in Eq. (9) makes the click’s probabilities sum to one at completion, and the paper claims exact unitarity nowhere.

### The symbols.

Table 4: The symbols: physics’ letters keep physics’ meanings, the law’s grains are  $N$  with a subscript, a world key is in code font.

| Symbol                                         | Kind                                               | Name                                                                                                                                                                                                                                                                                                                      |
|------------------------------------------------|----------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $E, E_0, m, \mathbf{p}, c, v, \beta$           | scalars, one vector                                | the energy, the rest energy $E_0 = E'_0 c^2$ with $E'_0 = N_l N_w M$ the rest energy in whole units, the mass $m = N_l N_w M$ in label units, the momentum, the limit speed $c = 1/\sqrt{3}$                                                                                                                              |
| $M, N_l, N_w, N_\phi, N_D, N_\theta, N_u, N_t$ | scalars                                            | Links per interval, the speed, $v/c$<br>the content (a count of units), the label’s scale, the width, the phase circle, the direction bound, the fan’s angular grain, the birth wheel, the cosine tables’ scale (the law’s documents write $Q, S$ and $N$ for $N_l, N_w$ and $N_\phi$ until their writers’ next pass [8]) |
| $\Phi; s, r, d, a, e$                          | operator; scalars                                  | the interval’s map; one accumulator of the state, its rate, its wall ( $d_p$ the drive’s), its cap, its event count                                                                                                                                                                                                       |
| $\mathbf{G}, U_s, R(f), R(o_A, o_B)$           | a matrix, a matrix written plain, a form, a scalar | the click’s Gram matrix, the label rotation (Section 6’s matrix, written plain there), the click’s reading (weight), a cell’s weight                                                                                                                                                                                      |
| $D, \mathbf{u}_D, T_D, S_1, K$                 | an integer vector written plain, a vector, scalars | a direction, its unit vector at the scale $N_l$ , its resolution, its Manhattan length, the number of directions of a fan (the ladder’s $K$ of $C_K$ and $b_K$ is Section 6’s)                                                                                                                                            |

| Symbol                                                                                                                     | Kind                                                    | Name                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $z, G, \rho$                                                                                                               | scalars                                                 | the redshift, Newton's constant, a family's charge per unit of content                                                                                                                                                                                                                                                                                                                 |
| $\mathbb{Z}_{N_\phi}, \mathbb{Z}[\mathbb{Z}_{N_\phi}], \mathbb{Z}[\zeta_{N_\phi}], \mathbb{Z}^2 \otimes \mathbb{Z}^2, B_3$ | a cyclic group, a group ring, a ring, a module, a group | the phase circle, the turn its shift; the arrivals' group ring, the merge its addition; its quotient by $x^{N_\phi/2} + 1$ , the cyclotomic integers, where opposite phases cancel; the pair's arms under the settings' integer matrices; the six Ports' hyperoctahedral group, 48 signed permutations, 24 rotations (Theorem 1); the flight the translation group's shift (Section 2) |

## C Reproduction

The code, the worlds with their expectation files written before the runs, the check scripts with their outputs, the runs' summary and the figure scripts are archived [88]; the register carries every run's world, duration and source fingerprint, and NUMBERS.md beside the manuscript maps each number of this paper to its source [5]; the criterion of every row of Table 2, what is measured, its kind, what would refute it and the re-run at head, is [44]. The register's consolidation (its head `b12dd253` on the branch `register-paper-sources`, pending merge; the merge commit replaces it) keeps every world this paper cites, and the worlds it removes are cited by no row of this paper; NUMBERS.md marks the rows whose worlds the paper no longer cites. The interpreter is Python 3.14, no floating point in any physical module, the runs seconds each. To run a registered world from the archive: `python -m pip install -e .` and then `python -m event_universe --init examples/events/amplitude/bell_0.24.json --output artifacts/bell_0.24`; the worlds of this paper are under `examples/events/amplitude/`, the pair `bell_*.json`, the two slits `slits_huygens.json`, the Mach-Zehnder `mz_equal.json`, `mz_half.json` and `mz_quarter.json`; a run writes its metadata and its events beside the output, and the check scripts read them. The runs' fingerprints: the register's `ff5c382d672f` (series L), `L7 4bf55a62e6fd`, `S 24e0c1ba`, `T afb533a`. The figures' runs carry the fingerprint `731d0f56c9f9` (`figures/summary.json` [5]). The archived version cited at submission is the tag `paper-2026-09-22` on the merge commit of the pull request that carries this cut, one tree whose history holds the trees named above and the log records this paper cites; it replaces the concept DOI below. The paper's law is the runtime before the generic entry of the bending: the runs' own commits named above, on the first-parent line of `main` up to `3a9a7109`, the last commit before the entry's merge `5b855791`; the generic entry is a later rule of the law (Section 2, P9), and the ring's run (Table 2, row 13) is this paper's one reading under it. Every record, note, design and register row this text cites is on the tree of this cut (main merged into it at `87c7ec6b`).

The long form of this manuscript, with the derivation record of every formula, is the tree's paper at the commit `c15c1174`; the forty-page cut before it at `b0d1ebf7`.

**The seven confirmations.** Each a run's clicks against the formula shown Inside, the rows in Tables 2 and 5: the pace, series Q, 290 of 290 primitive directions ( $|a| + |b| + |c| \leq 6$ ) at the derived interval, Node and face, 0.5718 to 0.5893 at finite ages; the click's power, series L, the (3, 4) split 63/1, 31/1, 125/3 at  $N_\phi = 64, 32, 128$ , the far pair's cells 27, 5, 5, 27; Mach-Zehnder, 64/0 equal arms, 0/64 at a half turn, 32/32 at a quarter, over 64 births; the marginals, 32/64 per party in every bin at  $N_\phi = 64$ ; GHZ, the four allowed triples 16 each, the products +1 and -1, the others 0; Malus (A12), 128, 0 and 219 of 256 at 45, 90 and 22.5 degrees; light beside a mass, series K, three worlds, the deflection 0.000 pixel, the delay 0.00 interval [4, 9].

**The transition from ordinary physics to modern algebra, dated.** The twenty-four dated steps, each a thing of physics becoming an object of the law with the reading that showed it, are [22] (`docs/designs/algebra_transition/HISTORY.md` on `main` at `412f5c61`), their dates once: 2026-09-17 the shared resource deleted and the amplitude a phase with a content; 09-18 the Born table from  $N_\phi$  and a Node its six Ports; 09-19 the quantum an integer row, the click the one border, the push one bilinear form,  $E = hf$  a release's cost; 09-20 the readings by kind, the masses the initialisation, every count an accumulator, the fan a width of the law; 09-21 the six verbs, the record in  $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ , the lattice the translation group, the Lorentz factor refused as a seventh verb; 09-22 the detector's clock, the click theorem and the framing.

**Use of AI tools.** The code, the documents, the checks and the drafts of this manuscript were produced with AI coding agents under the author’s direction and review; the author verified every number against the archived runs and is responsible for the whole text; no AI system is an author. [The tool and version are named at submission.]

## D The families in the algebra, and the check of the roads

Table 5: Every family of the register [10]: its declared integers ( $h$ ,  $\rho$ , the columns, the lifetime, the circle, the hand; the content  $M$  on the measured worlds), its algebraic object [47, 22], when it entered, the verbs that act on it (1 translation, 2 bilinear form, 3 group-ring addition, 4 permutation, 5 evaluation, 6 division) and the click that reads it, by kind; “not named” is a cell the tree does not fill.

| Family                                                      | The declared integers                                                                                                                                                               | The algebraic object                                                                                                                             | When it entered; the verbs                                                                                      | The click that reads it                                                                                   |
|-------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|
| <b>light</b> (the photon)                                   | $h = 1$ ; no charge; the circle turns; $M$ per unit 1; the amplitude worlds’ births $2^{20}$ to $2^{33}$                                                                            | a free row in $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ born at the lamp, its amount the coefficient, its direction on the fan under the 48              | the amplitude a phase and a content, 2026-09-17 (entry 2); $E = hf$ , 09-19 (entry 8); verbs 1, 2, 3, 5         | the two-slit pixels (L2b), the Mach-Zehnder ports 64/0, series Q’s 290 of 290 face clicks (DETECTOR)      |
| <b>e</b> (the electron)                                     | $h = 0$ ; $\rho = -15$ ; the circle turns; $M = 1836$ (atoms, bohr), 1 (gallery)                                                                                                    | the content on $\mathbb{Z}$ ; the charge one entry of the coupling matrix; the momentum an integer 3-vector; the phase on $\mathbb{Z}_{N_\phi}$  | the charge a rational per unit, 09-19 (entry 7); the masses the initialisation, 09-20 (entry 11); verbs 1, 2, 6 | series S’s face clicks 369, 345 (DETECTOR); Bohr’s ratio (row 6) not run                                  |
| <b>beta, w</b> (the decay’s electron and boson)             | $h = 1$ ; a whole charge $-7344$ per unit of amount; <b>w</b> ’s lifetime 1, no circle; <b>beta</b> ’s circle turns, its hand $-1$ on the world <b>wu</b>                           | a free row pushing by its label; the lifetime an integer wall; the hand a pseudoscalar in $\mathbb{Z}_2$                                         | the catalog, records 30 and 113 (09-20); verbs 1, 2, 6                                                          | the neutron’s decay clicks, series J1, 64 clicks, width over median 0.036 (DETECTOR, row 8a)              |
| <b>p</b> (the proton)                                       | $h = 0$ ; $\rho = [1, 1]$ (a winding), 4 on the atoms and binding worlds; no circle; $M = 1834, 1836$                                                                               | the content on $\mathbb{Z}$ ; the charge a rational pair in the coupling matrix; clicks by no phase                                              | the catalog, records 30 and 113 (09-20); verbs 1, 2, 6                                                          | series N’s border clicks: the deuteron’s held bond 0.109 percent, the alpha’s 2.0 (DETECTOR, rows 7a, 7b) |
| <b>n</b> (the neutron)                                      | $h = 0$ ; no charge; no circle; $M = 1837, 1839$                                                                                                                                    | the content on $\mathbb{Z}$ ; the momentum an integer 3-vector                                                                                   | as above; verbs 1, 2, 6                                                                                         | series J1’s decay clicks (row 8a); series N (row 7a) (DETECTOR)                                           |
| <b>nuclear, glue, bond</b> (the strong column and the bond) | $h = 0, 0, 1$ ; the column <b>strong</b> $\sigma = 10000$ with the sign $-1$ (7000 on one nucleus world); the lifetime 3; no circle                                                 | the column’s signed scalar per unit, one entry of the coupling matrix; the lifetime an integer wall                                              | a force a column with a sign and a lifetime, 09-20 (entry 9); verbs 2, 6                                        | series N’s border clicks (DETECTOR, rows 7a, 7b)                                                          |
| <b>u</b> (the up quark)                                     | $h = 0$ ; $\rho = 1224$ ; no circle                                                                                                                                                 | the content on $\mathbb{Z}$ , the charge on $\mathbb{Z}$                                                                                         | the catalog, record 30 (09-20); verbs 1, 2, 6                                                                   | no detector reading registered (the quarks worlds)                                                        |
| <b>nu, nubar</b> (the neutrino)                             | $h = 0$ ; no charge; no content per unit (row 8c); the circle turns; the hand $+1$ ( <b>nubar</b> ), $-1$ ( <b>nu</b> ) on the world <b>nu_hand</b> ; $M = 4096$ on the weak worlds | the hand a pseudoscalar in $\mathbb{Z}_2$ under the 48, kept by the rotations and negated by the reflections; the phase on $\mathbb{Z}_{N_\phi}$ | the hand, <b>hand-v1</b> [22]; verbs 1, 4, 5                                                                    | series J2, 16 of 1024 with 0 behind (DETECTOR, row 8b); the massless input refuted (row 8c)               |

| Family                                                                                              | The declared integers                                                                  | The object                                                                                                                                                                             | When; verbs                                                                                                                               | The click                                                                                                                                                                      |
|-----------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>m, mass, probe, neutron</b> (the massive bodies)                                                 | $h = 0; \rho = 0$ ; no circle; $M$ from 1 to $2^{26}$ as declared per world            | the held content on $\mathbb{Z}$ ; a body's momentum an integer 3-vector; its counts accumulators ( $s, r, d$ ); the age wall's member                                                 | the content, record 15 of 09-19; the masses the initialisation, 09-20 (entry 11); the clock's word, 09-21 to 22 (entry 21); verbs 1, 2, 6 | D3's births at the line, 138 of 139, $T(24)/T(12) = 1.997$ ; series T's 1.907; series X's $k = 0.9089$ ; series G2's $z = 0.2636$ (DETECTOR; T before the entry of 2026-09-22) |
| <b>q</b> (the test charge)                                                                          | $h = 0; \rho = [0, 1], [-1, 2]$ on the coupling worlds; the circle turns; $M = 2^{24}$ | the charge a rational pair in the coupling matrix                                                                                                                                      | the charge a rational per unit, 09-19 (entry 7); verbs 2, 6                                                                               | no detector reading registered (the coupling worlds)                                                                                                                           |
| <b>sa, sb</b> (the pair's choosers)                                                                 | $h = 0$ ; no charge; the circle turns                                                  | a setting's integer matrix $U_s$ in $M_2(\mathbb{Z})$ acting on the arms' label pair; the pair's record in $\mathbb{Z}[\mathbb{Z}_{N_\phi}] \otimes \mathbb{Z}^2 \otimes \mathbb{Z}^2$ | the record an element of $\mathbb{Z}[\mathbb{Z}_{N_\phi}]$ , 09-21 (entry 17); verbs 2, 3, 5                                              | the pair's clicks, $S = 2.75$ at $N_\phi = 64$ , the marginals 32/64 (DETECTOR, row 1a)                                                                                        |
| the apparatus materials ( <b>wall, screen, counter, apparatus, carrier, detector, d</b> )           | $h = 1$ ; no charge; the circle turns ( <b>d</b> no)                                   | the click's readers: a detector's Nodes, its own count the age wall's member at coefficient 1, its reading the moments of order 0 to 2                                                 | the readings the moments, 09-19 (entry 7); a detector's clock, 09-22 (entry 22); verbs 5, 6                                               | every DETECTOR row: they are the clicks                                                                                                                                        |
| the sources ( <b>s</b> ; the 24 thrown sources; the 24 star sources)                                | $h = 0$ or 1; no charge; the circle turns; $M = 64$ , 1024, 4096, 8192                 | a lamp's counts table on $\mathbb{Z}_W$ (the birth wheel); a throw's direction on the fan                                                                                              | the amplitude law, the wheel, 09-20 (entry 12); verbs 1, 6                                                                                | series G2's $z = 0.2636$ (row 4b); series T's lamp; the hubble worlds' $q_0 = -0.108$ (row 3) (DETECTOR; T's lamp before the entry of 2026-09-22)                              |
| the inline families <b>mu</b> (the muon, the covariant worlds) and <b>matter</b> (the massive rows) | $h = 0$ ; <b>mu</b> 's $M = 13248$ (series S); <b>matter</b> 's content per unit       | the record's $W = E'^2 + 3 \mathbf{p} \cdot \mathbf{p}$ , $E'$ the whole root by comparisons (a hypothesis beside the law); a paid row's content per unit on $\mathbb{Z}$              | the covariant readings, 09-21 (entry 20); the massive rows, series W (09-21); verbs 1, 2, 6                                               | the 64th self-creation at 70 and 124 (GAMEBOARD) and the products' face clicks (DETECTOR, series S); series W's first-click ages 815 and 876 (DETECTOR)                        |

Table 6: The check of the roads (record 817): one row per formula the paper arrives at, the inputs its chain used, the file and line where the chain starts (the click frame on **main** at 70e9781a [47]; Einstein Outside at e8432e6c [21], whose section III is the same check in the source) and the word. No row lists Einstein's or Lorentz's formula among its inputs; “derived” stands in no row.

| The formula arrived at                                       | The inputs its chain used                                                                                                   | Where the chain starts                                                             | The word                                                                                                                              |
|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------|
| $W = E_0'^2 + 3 \mathbf{p} \cdot \mathbf{p}$ to second order | (A1), (A2); the walk's invariant $\cos \omega = \cos m \cos \kappa$ ; the Planck map on <b>main</b> ; the unit $E' = E/c^2$ | the click frame, section 7                                                         | shown to second order (the conversion's identity); declared for the rows' dynamics (record 270)                                       |
| the boost, Lorentz's form                                    | (A1); two legs at one Node per interval; $k_{AB} = k_{BA}$                                                                  | the click frame, section 2 (b) :278–297; section 0 :84–132 (the group up to scale) | follows; Lorentz's form recovered, the scale $r$ free; (A2) fixes $r$ at $\sqrt{1-v^2}$                                               |
| $1/\gamma$ , the rate $r$                                    | Theorems 1, 2; $r$ read from two factors                                                                                    | Einstein Outside, II.1 :647                                                        | follows in $r$ ; $\sqrt{1-v^2}$ recovered under the line, shown to second order under (A2), declared under the identity, 1 on the law |
| $E^2 = E_0^2 + c^2 p^2$                                      | $E = E_0/r$ , $p = Ev$ , the line of II.1; the frame's section 7 for $W$                                                    | Einstein Outside, II.6 :887                                                        | follows as the line; $W$ shown to second order; exact by declaration under the identity; $E_0 = mc^2$ a declared unit                 |

| The formula                                                                                                 | The inputs                                    | Where                                                                                           | The word                                                                       |
|-------------------------------------------------------------------------------------------------------------|-----------------------------------------------|-------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------|
| the Outside step, place apart<br>and count apart of a<br>detector, Einstein's step its<br>most general form | (A1), (A3), the Inside step,<br>Theorems 1, 2 | Einstein Outside, section 3<br>(a) :481 (Einstein's form<br>named at :497 as the<br>comparison) | follows; Einstein's form<br>recovered where $r$ is<br>supplied, never an input |
| Newton's step as the limit                                                                                  | Theorem 4, the shell mean, $v \ll c$          | Einstein Outside, section 3<br>(c) :554                                                         | recovered (the limit)                                                          |