

CW complexes

Cloudifold

February 19, 2022

0 Basic Definitions and Lemmas

Definition 0.1. A CW-complex is a space constructed by successively attaching cells:
For $n \in \mathbb{N}, n \geq 1$, there are maps $\{\varphi_i : S^{n-1} \rightarrow X^{n-1}\}_{i \in I_n}$ (called characteristic maps). The way to construct X^n (called n -skeleton of X) is :
(starting from $X^0 = \prod_{I_0} *$)

$$\begin{array}{ccc} \prod_{i \in I_n} S^{n-1} & \xrightarrow{\prod_{i \in I_n} \varphi_i} & X^{n-1} \\ \downarrow & \searrow \tau & \downarrow \\ \prod_{i \in I_n} D^n & \longrightarrow & X^n \end{array} \quad (\text{pushout})$$

and the resulting CW-complex X is $\text{Colim}\{X^0 \rightarrow \dots \rightarrow X^n \rightarrow X^{n+1} \rightarrow \dots\}$. The images of $\overset{\circ}{D}_i^n$ in X is called open cell e_i^n of X .

Definition 0.2. A is a subcomplex of CW-complex X iff for any open cell e_i^n of X , A satisfy:
 $A \cap e_i^n \neq \emptyset \implies e_i^n \subseteq A$.

Pair of X and subcomplex $A : (X, A)$ is called a CW-pair.

Definition 0.3. The Infinite Symmetric Product of a pointed space (X, x_0) is colimit of its n -th Symmetric Products ($\text{SP}^n X := (\prod_{\{0,1,\dots,n-1\}} X)/S_n$) :

$$\begin{aligned} \text{Colim}\{\dots \hookrightarrow \text{SP}^n X \hookrightarrow \text{SP}^{n+1} X \hookrightarrow \dots\} \\ \{x_1, \dots, x_n\} \mapsto \{x_0, x_1, \dots, x_n\} \end{aligned}$$

Definition 0.4. For $n \geq 1$, a map between pairs $f : (X, A) \rightarrow (Y, B)$ is an n -equivalence if:

- $f_*^{-1}(\text{Im}(\pi_0 B \rightarrow \pi_0 Y)) = \text{Im}(\pi_0 A \rightarrow \pi_0 X)$
- For all choices of basepoint a in A ,

$$f_* : \pi_q(X, A, a) \rightarrow \pi_q(Y, B, f(a))$$

is isomorphism for $1 \leq q \leq n-1$ and epimorphism for $q = n$.

Definition 0.5. A pair (X, A) of topological spaces is **n -connected** if $\pi_0(A) \rightarrow \pi_0(X)$ is surjection and $\pi_q(X, A) = 0$ for $1 \leq q \leq n$.

Definition 0.6. For topological spaces $A \hookrightarrow X$, A is a **strong deformation retract** of a neighborhood V in X if:

$\exists h : V \times I \rightarrow X$ such that

$$\forall x \in V, h(x, 0) = x$$

$$h(V, 1) \subseteq A$$

$$\forall (a, t) \in A \times I, h(a, t) = a$$

Definition 0.7. For topological spaces $i : A \hookrightarrow X$, A is a **deformation retract** of X if:

$\exists h : X \times I \rightarrow X$ such that

$$\forall x \in X, h(x, 0) = x$$

$$h(X, 1) = A$$

$$\forall (a, t) \in A \times I, h(a, t) = a$$

(That is, there are retraction $r : X \rightarrow A$ and homotopy $h : \text{id}_X \simeq i \circ r \text{ rel } A$)

And $r := h(-, 1)$ is called a **deformation retraction**.

Definition 0.8. For topological spaces $A \hookrightarrow X$, a neighborhood V of A is **deformable** to A if:

$\exists h : X \times I \rightarrow X$ such that

$$\forall x \in X, h(x, 0) = x$$

$$h(A \times I) \subseteq A, h(V \times I) \subseteq V.$$

$$h(V, 1) \subseteq A$$

A criterion of weak homotopy equivalence:

Lemma 0.1. *The following on a map $e : Y \rightarrow Z$ and any fixed $n \in \mathbb{N}$ are equivalent:*

1. *For any $y \in Y$, $e_* : \pi_q(Y, y) \rightarrow \pi_q(Z, e(y))$ is monomorphism for $q = n$ and is epimorphism for $q = n + 1$.*
2. *(HELP of (D^{n+1}, S^n)) Given maps $f : D^{n+1} \rightarrow Z$, $g : S^n \rightarrow Y$ and homotopy $h : f \circ i \simeq e \circ g$:*

$$\begin{array}{ccc} S^n & \xhookrightarrow{i} & D^{n+1} \\ g \downarrow & \swarrow h & \downarrow f \\ Y & \xrightarrow{e} & Z \end{array}$$

then we have extension $g^+ : D^{n+1} \rightarrow Y$ of g and $h^+ : f \simeq e \circ g^+$:

$$\begin{array}{ccc} S^n & \xhookrightarrow{\quad} & D^{n+1} \\ g \downarrow & \swarrow g^+ & \downarrow f \\ Y & \xrightarrow{e} & Z \end{array}$$

h^+

3. *Conclusion above holds when the given h is $id_{f \circ i}$.*

Proof. Trivially 2. implies 3.

Our first goal : 3. implies 1.

Fix $n \in \mathbb{N}$. $\pi_n(e)$ is monomorphism:

For $n = 0$, 3. says if we have path $e(y) \simeq e(y')$ then we have path $y \simeq y'$. That is to say e can not map two path-connected component to one.

For $n > 0$, 3. says if $e \circ g$ is nullhomotopic, then $g : S^n \rightarrow Y$ could be extend to $g^+ : D^{n+1} \rightarrow Y$, which can be used to construct nullhomotopy of g .

Fix $n \in \mathbb{N}$. $\pi_{n+1}(e)$ is epimorphism:

For $[f] \in \pi_{n+1}(Z, e(y)) \cong [D^{n+1}, S^n; Z, e(y)]$, let $g := s \mapsto y$, the extension g^+ satisfy $e_*([g^+]) = [f]$, that proves e_* is epimorphism.

Second goal : 1. implies 2.

Fix g, f, h in the condition of 2. first. And observe that $\pi_n(Y, y) = [S^n, *; Y, y]$, $\pi_{n+1}(Y, y) = [D^{n+1}, S^n; Y, y]$.

There is a map $f' : (D^{n+1}, S^n) \rightarrow Z$ homotopic to f defined by $f' = f \circ b(-, 1)$ where

$$b : CS^n \times I \rightarrow CS^n$$


$$((x, t), s) \mapsto \begin{cases} \overline{(x, 1 - 2t)} & t \leq \frac{s}{2} \\ (x, \frac{t-s/2}{1-s/2}) & t \geq \frac{s}{2} \end{cases}$$

(recall that $D^{n+1} \simeq CS^n$) Therefore we can replace f with f' . Using the epimorphism leads to $h' : e \circ g^{+'} \simeq f'$, using the monomorphism leads to $r : g^{+'} \circ i \simeq g$. Construct $g^+ := a(-, 1)$ using

$$a : CS^n \times I \rightarrow Z$$

$$((x, t), s) \mapsto \begin{cases} r(x, s - 2t) & t \leq \frac{s}{2} \\ g^{+'}(x, \frac{t-s/2}{1-s/2}) & t \geq \frac{s}{2} \end{cases}$$

And that is the end of the proof:



□

1 Properties and Examples

Theorem 1.1. *Homotopy Extension and Lifting property:*


A : a topological space

X : result of successively attaching cells on A of dimensions $0, 1, \dots, k$ ($k \leq n$)

$e : Y \rightarrow Z$: n -equivalence


$g : A \rightarrow Y, f : X \rightarrow Z$

$h : f|_A \simeq e \circ g$




Then there exists $g^+ : X \rightarrow Y$ extends g ($g^+|_A = g$)

and $h^+ : X \times I \rightarrow Z$ extends h , $h^+ : f \simeq e \circ g^+$



Proof. It suffices to prove the case $A = S^{k-1}, X = D^k$, e is inclusion. (replace Z by M_e) Apply HEP of (D^k, S^{k-1}) :



$f' := \hat{h}(-, 1)$, replace f with f' the diagram would be strictly commute. Therefore, f' is map of pairs $(D^k, S^{k-1}) \rightarrow (Z, Y)$, $k \leq n$ implies f' is nullhomotopic, suppose $h^+ : D^k \times I \rightarrow Z$ is the nullhomotopy, then $g^+ := h^+(-, 1)$ satisfy $g^+(D^k) \subseteq Y$.

□

Note. In HELP, at condition $Y = Z$ and $e = \text{id}$, HELP says (X, A) have HEP

Corollary. *If*

A : a topological space

X : result of successively attaching cells on A of any dimensions

Then, (X, A) have HEP.

Theorem 1.2. *If X is an CW-complex, $e : Y \rightarrow Z$ is an n -equivalence, Then $e_* : [X, Y] \rightarrow [X, Z]$ is a bijection if $\dim X < n$, and a surjection if $\dim X = n$. (Also valid for pointed case)*

Proof. Surjectivity:

Apply HELP of (X, \emptyset) ((X, x_0) for pointed case) to obtain $e_*[g^+] \simeq [f]$:

$$\begin{array}{ccc} \emptyset & \longrightarrow & X \\ \downarrow & \nearrow g^+ & \downarrow f \\ Y & \xrightarrow{e} & Z \end{array}$$

Injectivity ($\dim X < n$):

Suppose $[g_0], [g_1] \in [X, Y]$, $e_*[g_0] = e_*[g_1]$.

Let $f : e \circ g_0 \simeq e \circ g_1$ Apply HELP to $(X \times I, X \times \partial I)$:

$$\begin{array}{ccc} X \times \partial I & \longrightarrow & X \times I \\ \downarrow g & \nearrow g^+ & \downarrow f \\ Y & \xrightarrow{e} & Z \end{array}$$

□

Corollary. *If X is a CW-complex, $e : Y \rightarrow Z$ is weak homotopy equivalence, then $e_* : [X, Y] \rightarrow [X, Z]$ is bijection.*

1.1 CW-approximation

Definition 1.1. A **CW-approximation** of $(X, A) \in \mathbf{Top}(\mathbf{2})$ is a CW-pair (\tilde{X}, \tilde{A}) and a weak homotopy equivalence of pairs $\varphi : (\tilde{X}, \tilde{A}) \rightarrow (X, A)$.

Lemma 1.3. φ, ψ are CW-approximations of X, Y , $f : X \rightarrow Y$, then

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{\varphi} & X \\ \exists \tilde{f} \downarrow \text{dashed} & & \downarrow f \\ \tilde{Y} & \xrightarrow{\psi} & Y \end{array}$$

commutes up to homotopy, and \tilde{f} is unique up to homotopy.

Proof. Directly from $\psi_* : [\tilde{X}, \tilde{Y}] \rightarrow [\tilde{X}, Y]$ is bijection. □

Theorem 1.4. φ, ψ are CW-approximations of $(X, A), (Y, B)$, $f : (X, A) \rightarrow (Y, B)$, then

$$\begin{array}{ccc} (\tilde{X}, \tilde{A}) & \xrightarrow{\varphi} & (X, A) \\ \exists \tilde{f} \downarrow \text{dashed} & & \downarrow f \\ (\tilde{Y}, \tilde{B}) & \xrightarrow{\psi} & (Y, B) \end{array}$$

commutes up to homotopy, and \tilde{f} is unique up to homotopy.

Proof. Apply Lemma 1.3 to obtain map $\tilde{f}_A : \tilde{A} \rightarrow \tilde{B}$ and homotopy $h : \psi|_{\tilde{B}} \circ \tilde{f}_A \simeq f \circ \varphi|_{\tilde{A}}$ Use

HELP of (\tilde{X}, \tilde{A}) to extend it:

$$\begin{array}{ccc}
 \tilde{A} & \hookrightarrow & \tilde{X} \\
 \tilde{f}_A \downarrow & & \downarrow \\
 \tilde{B} & \xrightarrow{\tilde{f}} & X \\
 \downarrow & & \downarrow \\
 \tilde{Y} & \longrightarrow & Y
 \end{array}$$

ψ_* is bijection implies the uniqueness up to homotopy of \tilde{f} . \square

Theorem 1.5. (Whitehead's Theorem)

Every n -equivalence between CW-complexes whose dimension is lower than n , is homotopy equivalence. Every weak homotopy equivalence between CW-complexes is homotopy equivalence.

Proof. $e : Y \rightarrow Z$ induce bijections $[Y, Y] \rightarrow [Y, Z]$ and $[Z, Y] \rightarrow [Z, Z]$, $[f] = e_*^{-1}[\text{id}_Z]$ implies $[e \circ f] = [\text{id}_Z]$ and $[e \circ f \circ e] = [e]$ ($[f \circ e] = e_*^{-1}[e] = [\text{id}_Y]$). \square

Corollary. CW-approximation is unique up to homotopy.

Example 1.1. Polish circle (Warsaw circle) : closed topologist's sine curve. It is n -connected for all n but not contractible.

Definition 1.2. A cellular map between CW-pairs is $g : (X, A) \rightarrow (Y, B)$ such that $g(A \cup X^n) \subseteq B \cup Y^n$.

Theorem 1.6. For any map between CW-pairs $f : (X, A) \rightarrow (Y, B)$ there exists a cellular map g such that $g \simeq f \text{ rel } A$

Proof. Construct g inductively:

Start from $A \cup X^0$:

take paths $\gamma_i : f(x_i) \simeq y_i$, where y_i is any point in Y^0 and $x_i \in X^0 - A$.

Construct $h_0 : (X^0 \cup A) \times I \rightarrow Y : h_0|_A(a, t) := f(a)$, $h_0|_{X^0-A}(x_i, t) := \gamma_i(t)$. This is a homotopy from f to $g_0 := h_0(-, 1) : A \cup X^0 \rightarrow B \cup Y^0$

Inductive step:

Assume $g_n : A \cup X^n \rightarrow B \cup Y^n$ and homotopy $h_n : f|_{A \cup X^n} \simeq g_n$ is given, try to construct g_{n+1} :
For each characteristic map $\varphi_i : S^n \rightarrow X^n$, take the resulting cell map $\varphi_i^+ : D^{n+1} \rightarrow X^{n+1}$ and use HELP of (D^{n+1}, S^n) :

$$\begin{array}{ccc}
 S^n & \hookrightarrow & D^{n+1} \\
 \varphi_i \downarrow & & \downarrow \varphi_i^+ \\
 X^n & \xrightarrow{g_{n+1,i}} & X^{n+1} \\
 g_n \downarrow & \swarrow h_{n+1,i} & \downarrow f \\
 B \cup Y^{n+1} & \hookrightarrow & Y
 \end{array}$$

Glue all $g_{n+1,i}$ and $h_{n+1,i}$ to produce g_{n+1} and $h_{n+1} : f|_{A \cup X^{n+1}} \simeq g_{n+1}$.

Final stage:

Maps g_n determine a cellular map $g : X \rightarrow Y$ since X has the final topology determined by skeletons. \square

1.2 Operation of CW-complexes

Product of CW-complexes:

Example 1.2. Product topology of two CW-complexes does not coincide with the final topology (union topology):

X (star of countably many edges) : $X = X^1 = \bigvee_{n \in \omega} I_n$

Y (star of ω^ω many edges) : $Y = Y^1 = \bigvee_{f \in \omega^\omega} I_f$ ($(I_n, 0) \cong (I_f, 0) \cong (I, 0)$)

Consider subset H of $X \times Y$: $H := \{(\frac{1}{f(n)+1}, \frac{1}{f(n)+1}) \in I_n \times I_f \mid n \in \omega, f \in \omega^\omega\}$.

H is closed under the final topology since every cell of $X \times Y$ contains at most one point of H . But closure of H contains $(0, 0)$ at product topology:

Let $U \times V$ be an open neighborhood (at product topology) of $(0, 0)$, let $g : \omega \rightarrow \omega - 0$ be an increasing function such that for all $n \in \omega$, $[0, \frac{1}{g(n)}) \subseteq U \cap I_n$, let $k \in \omega$ be sufficiently large that $\frac{1}{g(k)+1} \subseteq V \cap I_g$, then $(\frac{1}{g(k)+1}, \frac{1}{g(k)+1}) \in U \times V \cap H$.

Note. Another way to realize $X \times Y$ as CW-complex is to change its topology to the compactly generated topology $k(X \times Y)$.

Proposition 1.7. X and Y are CW-complexes, $X \times Y$ is CW-complex if

X or Y is locally compact

or

both X and Y have countably many cells.

Quotient of CW-pair:


Proposition 1.8. For CW-complex X and subcomplex A , the Quotient space X/A have a CW-complex structure induced by X and A .

Proof. Suppose the characteristic maps of X are indexed by $\{I_n\}_{n \in \mathbb{N}}$ and of A are indexed by $\{I'_n\}_{n \in \mathbb{N}}$ ($I'_n \subseteq I_n$). Then the characteristic maps of X/A are indexed by $\{K_n\}_{n \in \mathbb{N}}$, which defined below:

$K_0 := (I_0 - I'_0) \cup \{i_0\}$ where i_0 is an arbitrary element in I'_0

$K_n := I_n - I'_n$ for $n > 0$.

Verify the maps determine the CW-complex structure:



□

Smash product of CW-complexes:

Proposition 1.9. If (X, x_0) , (Y, y_0) are pointed CW-complexes with both countably many cell, and $X^{r-1} = \{x_0\}$, $Y^{s-1} = \{y_0\}$, then $X \wedge Y := X \times Y / X \vee Y$ is an $(r + s - 1)$ -connected CW-complex.

Proof. $X \times Y$ is CW-complex with cells of the form $e_{i,X}^n \times \{y_0\}$, $\{x_0\} \times e_{j,Y}^m$ or $e_{i,X}^n \times e_{j,Y}^m$ for $n \geq r$, $m \geq s$. Cells of the first two forms are contained in $X \vee Y$, therefore $(X \wedge Y)^{r+s-1} = *$. □

Corollary. If X is a pointed CW-complex, then $\Sigma^n X$ is an $(n - 1)$ -connected CW-complex.

1.3 Properties of Infinite Symmetric Product

Functoriality:

Pointed map $f : X \rightarrow Y$ induces

$$\begin{aligned} f_n : \mathrm{SP}^n X &\rightarrow \mathrm{SP}^n Y \\ \{x_1, \dots, x_n\} &\mapsto \{f(x_1), \dots, f(x_n)\} \end{aligned}$$

$$\begin{array}{ccccc} \longrightarrow & \mathrm{SP}^n X & \longrightarrow & \mathrm{SP}^{n+1} X & \longrightarrow \\ & \downarrow f_n & & \downarrow f_{n+1} & \\ \longrightarrow & \mathrm{SP}^n Y & \longrightarrow & \mathrm{SP}^{n+1} Y & \longrightarrow \end{array}$$

Which induces map $\mathrm{SP} f : \mathrm{SP} X \rightarrow \mathrm{SP} Y$. And Functorial properties are directly from the constructions above.

Commute with directed colimit:

Suppose P is a directed poset (that is $\forall x, y \in P, \exists z \in P, x \leq z, y \leq z$) and X_i are pointed spaces indexed by P satisfying $i \leq j \implies X_i \subseteq X_j$.

Then $\mathrm{SP}^n(\mathrm{Colim}_i X_i) \approx \mathrm{Colim}_i(\mathrm{SP}^n X_i)$

(Proof is obtained by showing that $\mathrm{SP}^n f$ is continuous iff f is, which implies final topology on $\mathrm{Colim}_i(\mathrm{SP}^n X_i)$ agree on $\mathrm{SP}^n(\mathrm{Colim}_i X_i)$)

Suppose $i : A \hookrightarrow X$ is an pointed inclusion, then $\mathrm{SP} i : \mathrm{SP} A \hookrightarrow \mathrm{SP} X$ is also inclusion. Furthermore, if A is open (or closed) in X , then $\mathrm{SP} A$ is open (or closed) in $\mathrm{SP} X$.

Pointed homotopy $h : X \times I \rightarrow Y$ induces

$$\begin{aligned} h_n : \mathrm{SP}^n X \times I &\rightarrow \mathrm{SP}^n Y \\ (\{x_1, \dots, x_n\}, t) &\mapsto \{h(x_1, t), \dots, h(x_n, t)\} \end{aligned}$$

which induces $\mathrm{SP} h : \mathrm{SP} X \times I \rightarrow \mathrm{SP} Y$.

Then we observe:

$f \simeq g$ implies $\mathrm{SP} f \simeq \mathrm{SP} g$,

$e : X \rightarrow Y$ is homotopy equivalence implies $\mathrm{SP} e : \mathrm{SP} X \rightarrow \mathrm{SP} Y$ is,

X is contractible implies $\mathrm{SP}^n X$ and $\mathrm{SP} X$ is.

Theorem 1.10. (*Dold-Thom Theorem*)

If X is T_2 space and A is closed path-connected subspace of X , and there is neighborhood V deformable to A in X .

Then the quotient map $q : X \rightarrow X/A$ induces quasi-fibration $\mathrm{SP} q : \mathrm{SP} X \rightarrow \mathrm{SP}(X/A)$, which satisfy $\forall x \in \mathrm{SP}(X/A), (\mathrm{SP} q)^{-1}\{x\} \simeq \mathrm{SP} A$.

Corollary. If X, Y are T_2 spaces and Y is connected, $f : X \rightarrow Y$. Then consider $X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X$, the map $p : C_f \rightarrow \Sigma X$ induces quasi-fibration $\mathrm{SP} p : \mathrm{SP} C_f \rightarrow \mathrm{SP}(\Sigma X)$ with fiber $\mathrm{SP} Y$.

Corollary. If X is T_2 and path-connected, then for any $q \geq 0$, there is $\pi_{q+1}(\mathrm{SP}(\Sigma X)) \cong \pi_q(\mathrm{SP} X)$.

Proof. CX is contractible implies $\mathrm{SP} CX$ is contractible, use the exact homotopy sequence of quasi-fibration to see:

$$\longrightarrow \pi_{q+1}(\mathrm{SP} CX) \longrightarrow \pi_{q+1}(\mathrm{SP} \Sigma X) \xrightarrow[\partial]{\cong} \pi_q(\mathrm{SP} X) \longrightarrow \pi_q(\mathrm{SP} CX) \longrightarrow$$

□

Note. The inverse of the isomorphism ∂ above is given by

$$[S^q, \mathrm{SP} X] \ni [g] \mapsto [\Sigma g] \in [S^{q+1}, \Sigma \mathrm{SP} X]$$

$(\Sigma \text{SP } X \cong \text{SP } \Sigma X)$. Because ∂ is given by:

$$\begin{array}{ccccc} \pi_q(\text{SP } \Sigma X) & \longrightarrow & \pi_q(\text{SP } CX, \text{SP } X) & \longrightarrow & \pi_{q-1}(\text{SP } X) \\ \Psi & & \Psi & & \Psi \\ [f] & \longmapsto & [\hat{f}] & \longmapsto & [\hat{f}]_{S^{q-1}} \end{array}$$

$$[p \circ Cg] = [\Sigma g] \longleftarrow [Cg] \longleftarrow [g]$$

□

Corollary. If X is T_2 space and A is path-connected subspace of X , then the canonical map $\text{SP}(X \cup (A \times I)) \rightarrow \text{SP}(X \cup CA)$ is a quasi-fibration with fiber $\text{SP } A$.

Theorem 1.11. If X is T_2 space and A is path-connected subspace of X , and $A \hookrightarrow X$ is a cofibration.

Then the quotient map $q : X \rightarrow X/A$ induces quasi-fibration $\text{SP } q : \text{SP } X \rightarrow \text{SP}(X/A)$, which satisfy $\forall x \in \text{SP}(X/A)$, $(\text{SP } q)^{-1}\{x\} \simeq \text{SP } A$.

Proof. If $A \hookrightarrow X$ is cofibration, then $X \cup CA \simeq X/A$ and $X \cup (A \times I) \simeq X$. □

Proposition 1.12. The inclusion $S^1 \rightarrow \text{SP } S^1$ is homotopy equivalence, therefore $\pi_q(S^1) \cong \pi_q(\text{SP } S^1)$.

Proof. $S^1 \simeq S^2 - \{0, \infty\}$

$\text{SP}^n S^2 = \{\{a_1, \dots, a_n\} \mid a_i \in \mathbb{C} \cup \{\infty\}\} = \{\prod_{\{a_1, \dots, a_n\}} (z - a_i) \mid a_i \in \mathbb{C} \cup \{\infty\}\}$ where $(z - \infty) := 1$
 $\text{SP}^n S^2 = \{f \in \mathbb{C}[z] - \{0\} \mid \deg(f) \leq n\} = \mathbb{C}\mathbb{P}^n$

$\text{SP}^n(S^2 - \{0, \infty\}) = \{f \in \mathbb{C}[z] - \{0\} \mid \deg(f) \leq n, f_n \neq 0, f_0 \neq 0\} = \mathbb{C}^n - \mathbb{C}^{n-1} \times 0 = \mathbb{C}^{n-1} \times (\mathbb{C} - 0)$

it have the same homotopy type of S^1

Corollary. $\pi_q(\text{SP } S^n) = \mathbb{Z}$ if $q = n$, otherwise $\pi_q(\text{SP } S^n) = 0$. (use corollary of 1.10 to see $\pi_{q+1}(\text{SP } \Sigma X) \cong \pi_q(\text{SP } X)$)

2 Homology Groups

2.1 Reduced Homology Groups

Definition 2.1. For a path-connected pointed CW-complex X , define its n -th **reduced homology group** for $n \geq 0$:

$$\tilde{H}_n(X) := \pi_n(\text{SP } X)$$

Note. All reduced homology groups are abelian since $\tilde{H}_n(X) \cong \tilde{H}_{n+1}(\Sigma X)$. Thus, we can extend the definition above to those X which does not necessarily be path-connected.

As SP , \tilde{H}_n also satisfy functoriality. Furthermore, \tilde{H}_n maps homotopic maps $f \simeq g$ to identical maps $f_* = g_*$. (SP maps homotopic maps to homotopic maps)

Exact Property:

Proposition 2.1. For any pointed map between CW-complexes $f : X \rightarrow Y$, we have an exact sequence:

$$\tilde{H}_n(X) \xrightarrow{f_*} \tilde{H}_n(Y) \xrightarrow{i_*} \tilde{H}_n(C_f)$$

where C_f is the mapping cone of f , $i : Y \hookrightarrow C_f$.

Proof. $Z_f := Y \cup_f (X \times I) / \{x_0\} \times I$ is the **reduced mapping cylinder** of f .
 $q : Z_f \rightarrow C_f$ is defined by

$$\begin{aligned} y &\mapsto y \\ \overline{(x, t)}^{Z_f} &\mapsto \overline{(x, t)}^{C_f} \end{aligned}$$

By Dold-Thom theorem, the induced map $\mathrm{SP} q$ is quasi-fibration $\mathrm{SP} Z_f \rightarrow \mathrm{SP} C_f$ with fiber $\mathrm{SP} X$.
By definition of quasi-fibration, we have

$$\pi_n(\mathrm{SP} X) \cong \tilde{H}_n(X) \xrightarrow{f_*} \pi_n(\mathrm{SP} Z_f) \cong \tilde{H}_n(Y) \xrightarrow{i_*} \pi_n(\mathrm{SP} C_f) = \tilde{H}_n(C_f)$$

□

Proposition 2.2. *There does not exist retraction $r : \mathbb{D}^n \rightarrow S^{n-1}$.*

Proof. $id = r \circ i : \mathbb{S}^{n-1} \rightarrow \mathbb{D}^n \rightarrow \mathbb{S}^{n-1}$ induces

$$id_* = r_* \circ i_* : \mathbb{Z} \cong \tilde{H}_{n-1} \mathbb{S}^{n-1} \rightarrow \tilde{H}_{n-1} \mathbb{D}^n \cong 0 \rightarrow \tilde{H}_{n-1} \mathbb{S}^{n-1} \cong \mathbb{Z}$$

which lead to contradiction. □

Theorem 2.3. *Fix-point theorem:*

If $f : \mathbb{D}^n \rightarrow \mathbb{D}^n$ is continuous, then exist $x_0 \in \mathbb{D}^n$ such that $x_0 = f(x_0)$.

Proof. (non-constructive) No such x_0 implies $\forall x \in \mathbb{D}^n, f(x) \neq x$ therefore, we can construct continuous retraction $r : \mathbb{D}^n \rightarrow \mathbb{S}^{n-1}$ by

$r(x) :=$ the intersection of “ray starting from $f(x)$ to x ” and \mathbb{S}^{n-1} . Contradict to 2.2.

Definition 2.2. Let (X, A) be an CW-pair, define the **n -th homology group** for $n \in \mathbb{N}$ of (X, A) be:

$$H_n(X, A) := \tilde{H}_n(X \cup CA)$$

And for single space:

$$H_n(X) := H_n(X, \emptyset) = \tilde{H}(X + 1)$$

where $X + 1 := X \sqcup *$.

Note. Map between CW-pair $f : (X, A) \rightarrow (Y, B)$, induces map $\bar{f} : X \cup CA \rightarrow Y \cup CB$ defined by $(x, t) \mapsto (f(x), t)$, which induces $f_* : \tilde{H}_n(X \cup CA) \rightarrow \tilde{H}_n(Y \cup CB)$ for any $n \in \mathbb{N}$.

2.2 Axioms for Homology

Definition 2.3. A (Ordinary) Homology Theory (on **TOP** with coefficient $G \in \mathbf{Ab}$) is functors $\{H_n(-, -; G) : \mathbf{TOP}(\mathbf{2}) \rightarrow \mathbf{Ab}\}_{n \in \mathbb{N}}$, with natural transformations $\partial_{n, (X, A)} : H_n(X, A; G) \rightarrow H_n(A, \emptyset; G)$ (called connecting homomorphism) satisfying following axioms:

- **Dimension:**

$$H_0(*, \emptyset; G) = G, \text{ for any } n > 0, H_n(*, \emptyset; G) = 0.$$

- **Weak Equivalence:**

Weak equivalence $f : (X, A) \rightarrow (Y, B)$ induces isomorphism

$$f_* : H_*(X, A; G) \rightarrow H_*(Y, B; G)$$

- **Long Exact Sequence:**

For any $(X, A) \in \mathbf{TOP}(\mathbf{2})$, maps $A \hookrightarrow X$ and $(X, \emptyset) \rightarrow (X, A)$ induce a long exact sequence together with ∂ :

$$\cdots \rightarrow H_{q+1}(A; G) \rightarrow H_{q+1}(X; G) \rightarrow H_{q+1}(X, A; G) \rightarrow H_q(A; G) \rightarrow \cdots$$

where $H_n(X; G) := H_n(X, \emptyset; G)$.

- **Additivity:**

If $(X, A) = \coprod_{\lambda} (X_{\lambda}, A_{\lambda})$ in **TOP(2)**, then inclusions $i_{\lambda} : (X_{\lambda}, A_{\lambda}) \rightarrow (X, A)$ induces isomorphism

$$(\bigoplus i_{*,\lambda}) : \bigoplus_{\lambda} H_*(X_{\lambda}, A_{\lambda}; G) \cong H_*(X, A; G)$$

- **Excision:**

If $(X; A, B)$ is an **excisive triad** (that is, $X = \overset{\circ}{A} \cup \overset{\circ}{B}$), then inclusion $(A, A \cap B) \hookrightarrow (X, B)$ induces isomorphism

$$H_*(A, A \cap B; G) \cong H_*(X, B; G)$$

Note. An equivalent form of Excision Axiom:

If $(X, A) \in \mathbf{TOP}(2)$, U is subspace of A and $\overline{U} \subseteq \overset{\circ}{A}$, then inclusion $i : (X - U, A - U) \hookrightarrow (X, A)$ induces isomorphism

$$i_* : H_*(X - U, A - U; G) \rightarrow H_*(X, A; G)$$

There is a critical criterion about weak homotopy equivalence between excisive triads, we prove lemmas first:

Lemma 2.4. *For*

$$\begin{array}{ccc} Z & \xrightarrow{f} & Y \\ \downarrow i & \lrcorner & \downarrow i_* \\ X & \xrightarrow{f_*} & X \cup_Z Y \end{array}$$

if D is deformation retract of X and $Z \subseteq D \subseteq X$, then $D \cup_Z Y$ is deformation retract of $X \cup_Z Y$.

Proof. Let $h : \text{id}_X \simeq r \circ i$ where r is the deformation retraction $X \rightarrow D$. Define $h_* : \text{id}_{X \cup_Z Y} \simeq (i \cup_Z \text{id}_Y) \circ (r \cup_Z \text{id}_Y)$

$$\begin{aligned} h_* : (X \cup_Z Y) \times I &\rightarrow X \cup_Z Y \\ (x, t) &\mapsto f_*(h(x, t)) \\ (y, t) &\mapsto i_*(y) \end{aligned}$$

Observe that $(X \cup_Z Y) \times I = (X \times I) \cup_{Z \times I} (Y \times I)$, check that h^* is continuous:

$$\begin{array}{ccccc} Z \times I & \longrightarrow & Y \times I & & \\ \downarrow & & \downarrow & \searrow \text{id}_{\text{id}_Y} & \\ X \times I & \longrightarrow & (X \cup_Z Y) \times I & & Y \\ & \searrow h & \searrow h_* & \downarrow i_* & \\ & & X & \xrightarrow{f_*} & X \cup_Z Y \end{array}$$

□

Lemma 2.5. *For maps $i : C \rightarrow A$, $j : C \rightarrow B$ define the double mapping cylinder $M(i, j) := A \cup_{C \times \{0\}} C \times I \cup_{C \times \{1\}} B$. If i is cofibration, then the quotient map*

$$\begin{aligned} q : M(i, j) &\rightarrow A \cup_C B \\ a &\mapsto a \\ b &\mapsto b \\ (c, t) &\mapsto c \end{aligned}$$

is a homotopy equivalence.

Proof.

$$\begin{array}{ccc} C & \longrightarrow & B \\ \downarrow i & & \downarrow \\ A & \xrightarrow{i_A} & A \cup_C B \end{array}$$

The canonical quotient $r : M_{i_A} \rightarrow A \cup_C B$ is a deformation retraction with homotopy:

$$\begin{aligned} h : (B \cup_{C \times 0} (A \times I)) \times I &\rightarrow B \cup_{C \times 0} (A \times I) = M_{i_A} \\ (a, t, s) &\mapsto (a, (1-s)t) \\ (b, s) &\mapsto b \end{aligned}$$

Observe that $C \times I \cup_C A \times \{1\}$ is a deformation retract of $A \times I$, since $i : C \rightarrow A$ is cofibration. Then we have $M(i, j) = B \cup_{C \times \{0\}} (C \times I \cup_{C \times \{1\}} A \times \{1\})$ is a deformation retract of $B \cup_{C \times \{0\}} A \times I = M_{i_A}$. (use lemma 2.4)

Finally, an easy check shows that $M(i, j) \rightarrow M_{i_A} \xrightarrow{r} A \cup_C B$ is identical to q .

□

Theorem 2.6. For excisive triads $(X; X_1, X_2)$, $(X'; X'_1, X'_2)$ and map $e : X \rightarrow X'$, if

$$\begin{aligned} e|_{X_1} : X_1 &\rightarrow X'_1 \\ e|_{X_2} : X_2 &\rightarrow X'_2 \\ e|_{X_3} : X_3 &\rightarrow X'_3 \end{aligned}$$

are weak equivalences, (where $X_3 := X_1 \cap X_2$, $X'_3 := X'_1 \cap X'_2$) then e is.

Proof. Use an important criterion of weak homotopy equivalence, it suffices to show for all $n \in \mathbb{N}$, any commutative diagram below:

$$\begin{array}{ccc} S^n & \xhookrightarrow{i} & D^{n+1} \\ \downarrow g & & \downarrow f \\ X & \xrightarrow{e} & X' \end{array}$$

can be filled like:

$$\begin{array}{ccc} S^n & \xhookrightarrow{i} & D^{n+1} \\ \downarrow g & \swarrow g^+ & \downarrow f \\ X & \xrightarrow{e} & X' \end{array}$$

$\nwarrow h$

whose upper triangle commutes.

Let

$$\begin{aligned} A_1 &:= g^{-1}(X - \overset{\circ}{X}_1) \cup f^{-1}(X' - \overset{\circ}{X}'_1) \\ A_2 &:= g^{-1}(X - \overset{\circ}{X}_2) \cup f^{-1}(X' - \overset{\circ}{X}'_2) \end{aligned}$$

which are disjoint closed subsets of D^{n+1} . Choose CW-complex structure on D^{n+1} such that for each n -cell σ_i , $\overline{\sigma_i} \cap (A_1 \cup A_2) = \overline{\sigma_i} \cap A_1$ or $\overline{\sigma_i} \cap A_2$. Now define

$$\begin{aligned} K_1 &:= \bigcup \{ \overline{\sigma_i} \mid g(\overline{\sigma_i} \cap S^n) \subseteq \overset{\circ}{X}_1 \text{ and } f(\overline{\sigma_i}) \subseteq \overset{\circ}{X}'_1 \} = \bigcup \{ \overline{\sigma_i} \mid \overline{\sigma_i} \cap A_1 = \emptyset \} \\ K_2 &:= \bigcup \{ \overline{\sigma_i} \mid g(\overline{\sigma_i} \cap S^n) \subseteq \overset{\circ}{X}_2 \text{ and } f(\overline{\sigma_i}) \subseteq \overset{\circ}{X}'_2 \} = \bigcup \{ \overline{\sigma_i} \mid \overline{\sigma_i} \cap A_2 = \emptyset \} \end{aligned}$$

which are subcomplexes of D^{n+1} and satisfy $K_1 \cup K_2 = D^{n+1}$. By HELP, we have:

$$\begin{array}{ccc} S^n \cap K_1 \cap K_2 & \xrightarrow{i} & K_1 \cap K_2 \\ g|_{K_1 \cap K_2} \downarrow & \swarrow g_0 & \downarrow f|_{K_1 \cap K_2} \\ X_1 \cap X_2 & \xrightarrow[e|_{X_1 \cap X_2}]{} & X'_1 \cap X'_2 \end{array}$$

$\nwarrow h_0$

such that h_0 is $f|_{K_1 \cap K_2} \simeq e \circ g_0 \text{ rel } (S^n \cap K_1 \cap K_2)$. Apply HELP to:

$$\begin{array}{ccc} (S^n \cup K_1) \cap K_2 & \xrightarrow{i_2} & K_2 \\ g_{K_2} \downarrow & \swarrow h_{K_2} & \downarrow f|_{K_2} \\ X_2 & \xrightarrow{\quad} & X'_2 \end{array} \quad \begin{array}{ccc} (S^n \cup K_2) \cap K_1 & \xrightarrow{i_1} & K_1 \\ g_{K_1} \downarrow & \swarrow h_{K_1} & \downarrow f|_{K_1} \\ X_1 & \xrightarrow{\quad} & X'_1 \end{array}$$

where

g_{K_i} are defined by $g_{K_i}|_{S^n \cap K_i} := g|_{S^n \cap K_i}$ and $g_{K_i}|_{K_1 \cap K_2} := g_0$,
 h_{K_2} are defined by (h_{K_1} is similar):

$$h_{K_2} : ((S^n \cup K_1) \cap K_2) \times I \rightarrow X'_2$$

$$(x, t) \mapsto \begin{cases} e(g(x)) & x \in S^n \cap K_2 \\ h_0(x, t) & x \in K_1 \cap K_2 \end{cases}$$

We get:

$$\begin{array}{ccc} (S^n \cup K_1) \cap K_2 & \xrightarrow{\quad} & K_2 \\ g_{K_2} \downarrow & \swarrow g_2 & \downarrow f|_{K_2} \\ X_2 & \xrightarrow[e|_{X_2}]{} & X'_2 \end{array} \quad \begin{array}{ccc} (S^n \cup K_2) \cap K_1 & \xrightarrow{\quad} & K_1 \\ g_{K_1} \downarrow & \swarrow g_1 & \downarrow f|_{K_1} \\ X_1 & \xrightarrow[e|_{X_1}]{} & X'_1 \end{array}$$

$\nwarrow h_2$ $\nwarrow h_1$

Define g^+ and $h : f \simeq g \text{ rel } S^n$ by $g^+|_{K_i} := g_i$ and $h|_{K_i \times I} := h_i$.
 $h|_{S^n \times I} = (e \circ g) \times \text{id}_I$ (h is $\text{rel } S^n$) since $h_i(-, t)|_{S^n \cap K_i} = h_{K_i}(-, t)|_{S^n \cap K_i} = e \circ g|_{S^n \cap K_i}$. □

Note. The proof above can be easily modified to case each weak equivalence appear in the statement is an n -equivalence.

Following theorem allow us to use CW-triads to approximate excisive triads:

Theorem 2.7. *For any excisive triad $(X; A, B)$, there is a CW-triad $(\tilde{X}; \tilde{A}, \tilde{B})$ (A CW-triad $(X; A, B)$ is X and its subcomplex A, B such that $A \cup B = X$) and a map $r : \tilde{X} \rightarrow X$ such that*

$$\begin{aligned} r|_{\tilde{A}} : \tilde{A} &\rightarrow A \\ r|_{\tilde{B}} : \tilde{B} &\rightarrow B \\ r|_{\tilde{C}} : \tilde{C} &\rightarrow C \\ r : \tilde{X} &\rightarrow X \end{aligned}$$

are all weak homotopy equivalences (where $\tilde{C} := \tilde{A} \cap \tilde{B}$, $C := A \cap B$). Furthermore, such r is natural up to homotopy.

Proof. Choose a CW-approximation $r_C : \tilde{C} \rightarrow C$ and extend it to $r_A : \tilde{A} \rightarrow A$, $r_B : \tilde{B} \rightarrow B$.
 $\tilde{X} := \tilde{A} \cup_{\tilde{C}} \tilde{B}$. $i : \tilde{C} \rightarrow \tilde{A}$ and $j : \tilde{C} \rightarrow \tilde{B}$ are cofibrations, by lemma 2.5 we have homotopy

equivalence $q : M(i, j) \rightarrow \tilde{X}$, which induces homotopy equivalence of triads:

$$\begin{aligned} q : M(i, j) &\rightarrow \tilde{X} \\ q| : \tilde{A} \cup (\tilde{C} \times [0, \frac{2}{3})) &\rightarrow \tilde{A} \\ q| : \tilde{B} \cup (\tilde{C} \times (\frac{1}{3}, 1]) &\rightarrow \tilde{B} \end{aligned}$$

then we can deduce that $r \circ q$ is a weak homotopy equivalence by theorem 2.6. Consequently, r is weak homotopy equivalence. r is natural up to homotopy since each CW-approximation r_C, r_A, r_B is. □

Then we have:

Definition 2.4. A (Ordinary) Homology Theory on CW-complexes with coefficient $G \in \mathbf{Ab}$ is functors $\{H_n(-, -; G) : \mathbf{CW}\text{-pairs} \rightarrow \mathbf{Ab}\}_{n \in \mathbb{N}}$, with natural transformations $\partial_{n, (X, A)} : H_n(X, A; G) \rightarrow H_n(A, \emptyset; G)$ (called connecting homomorphism)

satisfying axioms with the excision axiom changed to:

- **Excision:**

If $(X; A, B)$ is an **CW-triad** (that is $X = A \cup B$ for subcomplexes A and B) then the inclusion $(A, A \cap B) \hookrightarrow (X, B)$ induces isomorphism

$$H_*(A, A \cap B; G) \cong H_*(X, B; G)$$

Proposition 2.8. The homology groups defined in definition 2.2 is a ordinary homology theory on CW-complexes with coefficient \mathbb{Z} .

Proof.

- Dimension: by a corollary, $H_q(*, \emptyset) = \pi_q(\mathrm{SP} S^0) = \begin{cases} \mathbb{Z} & q = 0 \\ 0 & q \geq 1 \end{cases}$.

- Weak Equivalence: SP preserves weak equivalence.
- Long Exact Sequence: use a corollary of Dold-Thom theorem.
- Additivity: For index set Λ , $P := \{S \mid S \subseteq \Lambda\}$.

Then define $Y_S := \bigvee_{\lambda \in S} X_\lambda \cup CA_\lambda = (\coprod_{\lambda \in S} X_\lambda) \cup C(\coprod_{\lambda \in S} A_\lambda)$,

and use fact that SP commutes with directed colimit, we have

$$\bigvee_{\lambda \in \Lambda} \mathrm{SP}(X_\lambda \cup CA_\lambda) = \mathrm{Colim}_{S \in P} \mathrm{SP} Y_S \approx \mathrm{SP}(\mathrm{Colim}_{S \in P} Y_S) = \mathrm{SP}((\coprod_{\lambda \in \Lambda} X_\lambda) \cup C(\coprod_{\lambda \in \Lambda} A_\lambda)) = \mathrm{SP}(X \cup CA).$$

Which induces $\bigoplus_{\lambda \in \Lambda} \tilde{H}_n(X_\lambda \cup CA_\lambda) \cong \pi_n(\bigvee_{\lambda \in \Lambda} \mathrm{SP}(X_\lambda \cup CA_\lambda)) \cong \pi_n(\mathrm{SP}(X \cup CA)) = \tilde{H}_n(X \cup CA).$

- Excision: For CW-triad $(X; A, B)$, $A/(A \cap B) \approx X/B$. Apply theorem 1.11 to $(Y \cup CZ, CZ)$ to show that $H_n(Y, Z) \cong \tilde{H}_n(Y/Z)$. □

3 Homotopy and Eilenberg-Mac Lane Spaces

Theorem 3.1. (Blakers–Massey) Homotopy Excision Theorem:

For pointed CW-triad $(X; A, B)$ such that $C := A \cap B \neq \emptyset$, if (A, C) is $(m-1)$ -connected and (B, C) is $(n-1)$ -connected where $m \geq 2$, $n \geq 1$. Then $i : (A, C) \rightarrow (X, B)$ is an $(m+n-2)$ -equivalence for pairs.

Note. We can replace the "CW-triad" with "excisive triad" in condition by theorem 2.7.

Proof. Define (pointed) the triad homotopy group for $q \geq 2$:

$$\pi_q(X; A, B) := \pi_{q-1}(P_{i_{B,X}}, P_{i_{C,A}})$$

where $i_{B,X} : B \hookrightarrow X$, $i_{C,A} : C \hookrightarrow A$ and P_f is the homotopy fiber

$$\{(y, \gamma) \in Y \times M(I, Z)_* \mid \gamma(1) = f(y)\}$$

of pointed map $f : Y \rightarrow Z$. Use long exact sequence of pairs:

$$\begin{aligned} \cdots \rightarrow \pi_q(P_{i_{B,X}}, P_{i_{C,A}}) \rightarrow \pi_{q-1}(P_{i_{C,A}}) \rightarrow \pi_{q-1}(P_{i_{B,X}}) \rightarrow \pi_{q-1}(P_{i_{B,X}}, P_{i_{C,A}}) \rightarrow \pi_{q-2}(P_{i_{C,A}}) \rightarrow \cdots \\ \cdots \rightarrow \pi_1(P_{i_{B,X}}, P_{i_{C,A}}) \rightarrow \pi_0(P_{i_{C,A}}) \rightarrow \pi_0(P_{i_{B,X}}) \end{aligned}$$

and observe that $\pi_q(P_{i_{X,B}}) \cong \pi_{q+1}(X, B)$ since for any $f : S^q \rightarrow P_{i_{X,B}}$ we have:

$$\begin{array}{ccccc} S^q & & & & \\ & \searrow f & & \searrow f' & \\ & P_{i_{B,X}} & \longrightarrow & M(I, X)_* & \\ & \downarrow g & \lrcorner & \downarrow & \\ & B & \longrightarrow & X & \end{array}$$

use the fact $f' \in M(S^q, M(I, X)_*)_* \cong M(S^q \wedge I, X)_* \ni f''$ and $S^q \wedge I \approx D^{q+1}$ with

$$\begin{aligned} S^q \hookrightarrow S^q \wedge I \approx D^{q+1} \\ s \mapsto (s, 1) \end{aligned}$$

the condition $f'(s)(1) = g(s)$ is equivalent to $f''((s, 1)) = g(s)$, that is have a map f is equivalent to have a map $f'' : (D^{q+1}, S^q) \rightarrow (X, B)$. With the analogue statement also valid for homotopies $S^q \times I \rightarrow P_{i_{X,B}}$, we have $\pi_q(P_{i_{B,X}}) = [S^q, *; P_{i_{B,X}}, *] \cong [D^{q+1}, S^q; X, B] = \pi_{q+1}(X, B)$.

Rewrites the long exact sequence of pairs above to:

$$\begin{aligned} \cdots \rightarrow \pi_{q+1}(X; A, B) \rightarrow \pi_q(A, C) \rightarrow \pi_q(X, B) \rightarrow \pi_q(X; A, B) \rightarrow \pi_{q-1}(A, C) \rightarrow \cdots \\ \cdots \rightarrow \pi_2(X; A, B) \rightarrow \pi_1(A, C) \rightarrow \pi_1(X, B) \end{aligned}$$

Conditions $m \geq 1, n \geq 1$ guarantees $\pi_0(C) \rightarrow \pi_0(A)$ and $\pi_0(C) \rightarrow \pi_0(B)$ are surjections.

$m \geq 2$ is equivalent to $\pi_1(A, C) = 0$, which implies $\pi_0(C) \rightarrow \pi_0(A)$ is bijection.

For $x \in \pi_0(A \cap_C B)$, we can always find $b \in \pi_0(B)$, $i_{B,X} \ast(b) = x$ or $a \in \pi_0(A)$, $i_{A,X} \ast(a) = x$ which becomes $b \in \pi_0(B)$, $i_{B,X} \ast(b) = x$ or $c \in \pi_0(C)$, $i_{C,X} \ast(c) = x$ when $\pi_0(C) \rightarrow \pi_0(A)$ is bijection. That is equivalent to $\pi_0(B) \rightarrow \pi_0(X)$ is bijection, which means $\pi_1(X, B) = 0$.

We only need to show that for $2 \leq q \leq m + n - 2$, $\pi_q(X; A, B) = 0$.

With $J^{q-1} := (\partial I^{q-1} \times I) \cup (I^{q-1} \times \{0\})$, we have:

$$\begin{aligned}\pi_q(P_{i_{B,X}}, P_{i_{C,A}}) &= [I^q, \partial I^q, J^{q-1}; P_{i_{B,X}}, P_{i_{C,A}}, *] \\ &= [I^q \wedge I; I^q, \partial I^q \wedge I, J^{q-1} \wedge I \rightarrow X; B, A, *]\end{aligned}$$

$$:= \text{relative homotopy classes of pointed maps } f : I^q \wedge I \rightarrow X \text{ who satisfy: } \begin{cases} f(I^q) & \subseteq B \\ f(\partial I^q \wedge I) & \subseteq A \\ f(\partial I^q) & \subseteq C \\ f(J^{q-1} \wedge I) & = * \end{cases}$$

“relative” means the homotopy h determine the classes satisfy:

$$\begin{cases} h(I^q \times I) & \subseteq B \\ h((\partial I^q \wedge I) \times I) & \subseteq A \\ h(\partial I^q \times I) & \subseteq C \\ h((J^{q-1} \wedge I) \times I) & = * \end{cases}$$

(notice that $\partial I^q \wedge I \cap I^q = \partial I^q$, therefore $f(\partial I^q) \subseteq A \cap B = C$)

(this is called (relative) homotopy class of maps of tetrads)

$$\begin{aligned}&= [(I^q \times I)/K; I^q \times \{1\}, (\partial I^q \times I)/K, (J^{q-1} \times I)/K \rightarrow X; B, A, *] \\ &\quad (K := I^q \times \{0\} \cup \{i_0\} \times I)\end{aligned}$$

$$= [I^{q+1}; (I^q \times \{1\}) \cup K, (\partial I^q \times I) \cup K, J^{q-1} \times I \cup K \rightarrow X; B, A, *]$$

$$= [I^{q+1}; I^q \times \{1\}, I^{q-1} \times \{1\} \times I, J^{q-1} \times I \cup I^q \times \{0\} \rightarrow X; B, A, *]$$

(notice that $\partial I^q = \partial I^{q-1} \times I \cup I^{q-1} \times \{0, 1\}$)

We can assume that (A, C) have no relative $q < m$ -cells and (B, C) have no relative $q < n$ -cells. And we can assume that X has finite many cells since I^q is compact.

For subcomplexes $C \subseteq A' \subseteq A$, where $A = e^m \cup A'$ (attaching one cell from A').

Let $X' := A' \cup_C B$, if the results hold for $(X'; A', B)$ and $(X; A, X')$, then it hold for $(X; A, B)$ since we have map between exact homotopy sequences of triples (A, A', C) and (X, X', B) :

$$\begin{array}{ccccccccc} \pi_{q+1}(A, A') & \longrightarrow & \pi_q(A', C) & \longrightarrow & \pi_q(A, C) & \longrightarrow & \pi_q(A, A') & \longrightarrow & \pi_{q-1}(A', C) \\ \downarrow i_{2,q+1} & & \downarrow i_{1,q} & & \downarrow i_{3,q} & & \downarrow i_{2,q} & & \downarrow i_{1,q-1} \\ \pi_{q+1}(X, X') & \longrightarrow & \pi_q(X', B) & \longrightarrow & \pi_q(X, B) & \longrightarrow & \pi_q(X, X') & \longrightarrow & \pi_{q-1}(X', B) \end{array}$$

induced by inclusion $(A, A', C) \hookrightarrow (X, X', B)$. If the result hold for $(X'; A', B)$ and $(X; A, X')$, maps $i_{1,q}$, $i_{2,q}$ are isomorphisms when $1 \geq q \geq m + n - 3$, are epimorphisms when $q = m + n - 2$. Notice the 5-lemma says that

if $i_{1,q}$ and $i_{2,q}$ are epimorphisms, $i_{1,q-1}$ are monomorphism, then $i_{3,q}$ is epimorphism.

if $i_{1,q}$ and $i_{2,q}$ are monomorphisms, $i_{2,q+1}$ are epimorphism, then $i_{3,q}$ is monomorphism.

We also have if $C \subseteq B' \subseteq B$ with $B = B' \cup e^n$, the result hold for CW-triads $(X'; A, B')$ and $(X; X', B)$ where $X' = A \cup_C B'$, since $(A, C) \hookrightarrow (X, B)$ factors as $(A, C) \hookrightarrow (X', B') \hookrightarrow (X, B)$.

Now we can assume that $A = C \cup D^m$ and $B = C \cup D^n$.

The current goal of proof is to prove any

$$f : (I^{q+1}; I^q \times \{1\}, I^{q-1} \times \{1\} \times I, J^{q-1} \times I \cup I^q \times \{0\}) \rightarrow (X; B, A, *)$$

is nullhomotopic for any $q + 1$ with $2 \leq q + 1 \leq m + n - 2$.

For $a \in \mathring{D}^m$, $b \in \mathring{D}^n$ We have inclusions of based triads:

$$(A; A, A - \{a\}) \hookrightarrow (X - \{b\}; X - \{b\}, X - \{a, b\}) \hookrightarrow (X; X - \{b\}, X - \{a\}) \hookleftarrow (X; A, B)$$

The first and the third induces isomorphisms on homotopy groups of triads since B is a strong deformation retract of $X - \{a\}$ in X and A is a strong deformation retract of $X - \{b\}$ in X . $\pi_*(A; A, A - \{a\}) = 0$ since $\pi_*(A, A - \{a\}) \rightarrow \pi_*(A, A \cap (A - \{a\}))$ are isomorphisms.

Current goal : choose good a, b to show f regarded as a pointed triad map to $(X; X - \{b\}, X - \{a\})$ is homotopic to a map

$$f' : (I^{q+1}; I^{q-1} \times \{1\} \times I, I^q \times \{1\}, J^{q-1} \times I \cup I^q \times \{0\}) \rightarrow (X - \{b\}; X - \{b\}, X - \{a, b\}, *)$$

if $2 \leq q + 1 \leq m + n - 2$.

Note. We want to homotopically remove some point $f^{-1}(b)$, first we may want to construct some Urysohn function u separating $f^{-1}(a) \cup J^{q-1} \times I \cup I^q \times \{0\}$ and $f^{-1}(b)$ and construct homotopy of cube $h^+ : (r, s, t) \mapsto (r, (1 - u(r, s)t)s)$ wishing that $f(h^+(r, s, 1))$ would miss b . The problem in this method is that points $f^{-1}(b)$ in the cube would be homotopically replaced by other points. Since our desire homotopy does not change the first q coordinates of the cube, we want to separate $p^{-1}(p(f^{-1}(a))) \cup J^{q-1} \times I$ and $p^{-1}(p(f^{-1}(b)))$ (where $p : I^q \times I \rightarrow I^q$). Our problem is to find suitable a, b such that $p(f^{-1}(a)) \cap p(f^{-1}(b)) = \emptyset$.

We use manifold structure on D^m and D^n to achieve it, now we homotopically approximate f by a map g which smooth on $f^{-1}(D_{<1/2}^m \cup D_{<1/2}^n)$.

Let $U_{<r} := f^{-1}(D_{<r}^m \cup D_{<r}^n)$, Use smooth deformation theorem to construct smooth map (for any $0 < \epsilon$) $g' : U_{<3/4} \rightarrow D_{<3/4}^m \cup D_{<3/4}^n$ with homotopy $h_1 : g' \simeq f|_{U_{<3/4}}$ (and bound $|g'(x) - f(x)| < \epsilon$ for any $x \in U_{<1}$) and take partition of unity $\{\rho, \rho'\}$ with subcoordinates $\{I^{q+1} - U_{<1/2}, U_{<3/4}\}$, we have:

$$\begin{aligned} g &:= \rho f + \rho' g' \\ h_2 : g &\simeq f \text{ rel } (I^{q+1} - U_{<3/4}) \\ h_2 : I^{q+1} \times I &\rightarrow X \\ (x, t) &\mapsto \rho(x)f(x) + \rho'(x)h_1(x, t) \end{aligned}$$

with scalar multiplication and addition is already defined on smooth structure on $D_{<3/4}^m \cup D_{<3/4}^n$. We could assume that $g(I^{q-1} \times \{1\} \times I) \cap D_{<1/2}^n = \emptyset$ (which implies g is a map of tetrads to $(X; X - \{b\}, X - \{a\}, *)$) and $g(I^q \times \{1\}) \cap D_{<1/2}^m = \emptyset$ since $f(I^{q-1} \times \{1\} \times I) \subseteq A$ and $f(I^q \times \{1\}) \subseteq B$ and we can always tighten the bound ϵ , (Similar argument also hold for h_2 , then we have $h_2 : g \simeq f$ as homotopy between maps of tetrads.)

Use the manifold structure to find good (a, b) :
 $V := g^{-1}(D_{<1/2}^m) \times g^{-1}(D_{<1/2}^n)$ is a sub-manifold of $I^{2(q+1)}$. Consider $W := \{(v, v') \in V \mid p(v) = p(v')\}$, which is the zero set of smooth submersion $(v, v') \mapsto p(v) - p(v')$. W is smooth manifold with codimension q . Therefore the map $(g, g) : W \rightarrow D_{<1/2}^m \times D_{<1/2}^n$ is smooth map between manifolds of dimension $q + 2$ and $m + n$. The map is not surjection since $q + 2 < m + n$. Then we have $(a, b) \notin (g, g)(W)$ (that is, $p(g^{-1}(a)) \cap p(g^{-1}(b)) = \emptyset$).

Since $g(I^{q-1} \times \{1\} \times I) \cap D_{<1/2}^n = \emptyset$ and $g(J^{q-1} \times I) \cap D_{<1/2}^n = \emptyset$, we have $g(\partial I^q \times I) \cap D_{<1/2}^n = \emptyset$. By Urysohn's lemma, we have $u : I^q \rightarrow I$ separating $p(g^{-1}(a)) \cup \partial I^q$ and $p(g^{-1}(b))$. Finally we have:

$$\begin{aligned} h' : I^q \times I \times I &\rightarrow I^q \times I \\ (r, s, t) &\mapsto (r, (1 - u(r)t)s) \end{aligned}$$

and $h := g \circ h'$, $f' := h(-, 1)$. $f'(I^{q+1}) \cap \{b\} = \emptyset$ since if $\exists(r, s) \in I^q \times I$, $f'(r, s) = b$, then $b = g(r, (1 - u(r))s) = g(r, 0) = *$ leads to contradiction.

Last step is to check that h is a homotopy between maps

$$(I^{q+1}; I^{q-1} \times \{1\} \times I, I^q \times \{1\}, J^{q-1} \times I \cup I^q \times \{0\}) \rightarrow (X; X - \{b\}, X - \{a\}, *)$$

Since g is, $g \circ h'$ is too. □

Corollary. Suppose that $Y_0 \hookrightarrow Y$ is cofibration, (Y, Y_0) is $(r-1)$ -connected and Y_0 is $(s-1)$ -connected, then $(Y, Y_0) \rightarrow (Y/Y_0, *)$ is $(r+s-1)$ -equivalence. ($r \geq 2, s \geq 1$)

Proof. $Y_0 \hookrightarrow CY_0$ is cofibration and (CY_0, Y_0) is s -connected. Use homotopy excision theorem (with $X = Y \cup CY_0, A = Y, B = CY_0, C = Y_0$) to see $(Y, Y_0) \rightarrow (Y \cup CY_0, CY_0)$ is $(r+s-1)$ -equivalence. And $(Y \cup CY_0, CY_0) \rightarrow (Y/Y_0, *)$ is homotopy equivalence since $Y_0 \hookrightarrow Y$ is cofibration. □

Corollary. For $n \geq 2$, $f : X \rightarrow Y$ is $(n-1)$ -equivalence between $(s-1)$ -connected spaces, then $(M_f, X) \rightarrow (C_f^+, *)$ is $(n+s-1)$ -equivalence. Where $C_f^+ := Y \cup_f C^+X, C^+X := (X \times I)/(X \times \{1\})$. is the unreduced mapping cone and the unreduced cone.

Proof. f is $(n-1)$ -equivalence implies (M_f, X) is $(n-1)$ -connected. Use corollary above. □

Corollary. For $n \geq 2$, if $f : X \rightarrow Y$ is pointed map between $(n-1)$ -connected well-pointed spaces (that is, pointed space whose inclusion of the base point is cofibration). Then C_f is $(n-1)$ -connected and $\pi_n(M_f, X) \rightarrow \pi_n(C_f, *)$ is isomorphism.

Proof. Use homotopy extension property to extend to unreduced case. f is map between $(n-1)$ -connected space implies f is at least a $(n-1)$ -equivalence. Therefore $(M_f, X) \rightarrow (C_f, *)$ is $(2n-1)$ -equivalence, Since we have $n < 2n-1$ for any $n \geq 2$, $\pi_n(M_f, X) \rightarrow \pi_n(C_f, *)$ is isomorphism. □

Theorem 3.2. (Freudenthal Suspension Theorem) If X is well-pointed and $(n-1)$ -connected ($n \geq 1$), then the map:

$$\begin{aligned} \sigma : \pi_q(X) &\rightarrow \pi_{q+1}(\Sigma X) \\ f &\mapsto \Sigma f \end{aligned}$$

is isomorphism if $q < 2n-1$ and epimorphism if $q = 2n-1$.

Proof. If we have $f : (I^q, \partial I^q) \rightarrow (X, *)$ then $f \times \text{id}_I : I^{q+1} \rightarrow X \times I$ will give a map $\overline{f \times \text{id}_I} : (I^{q+1}, \partial I^{q+1}, \partial I^q \times I \cup \partial I \times \{1\}) \rightarrow (CX, X, *)$ since $J^q = \partial I^q \times I \cup \partial I \times \{0\}$, it does not give a map in $\pi_{q+1}(CX, X)$. we should change $\overline{f \times \text{id}_I}$ into $\overline{f \times -\text{id}_I}$. we have commutative diagram:

$$\begin{array}{ccc} \pi_{q+1}(CX, X) & \xrightarrow{p_*} & \pi_{q+1}(CX/X, *) & \xrightarrow{\quad} & \overline{[f \times -\text{id}_I]} & \xrightarrow{\quad} & [p \circ \overline{[f \times -\text{id}_I]}] \\ \partial \left(\begin{array}{c} \uparrow \\ \downarrow \end{array} \right) i & & \parallel & & \left(\begin{array}{c} \uparrow \\ \downarrow \end{array} \right) & & \parallel \\ \pi_q(X) & \xrightarrow{-\sigma} & \pi_{q+1}(\Sigma X) & & [f] & \xrightarrow{\quad} & [-\Sigma f] \end{array}$$

Where $p : (CX, X) \rightarrow (CX/X, *)$ is the canonical quotient map and $i : [f] \rightarrow \overline{[f \times -\text{id}_I]}$ makes $\pi_{q+1}(CX) \rightarrow \pi_{q+1}(CX, X) \rightarrow \pi_q(X) \rightarrow \pi_q(CX)$ split in middle (that is, i is inverse of the connecting homomorphism ∂). We verify the commutativity:

$$\begin{aligned} -\Sigma f : (I^{q+1}, \partial I^{q+1}) &\rightarrow (CX/X, *) \\ (s, t) &\mapsto f(s) \wedge (1-t) \\ p \circ \overline{[f \times -\text{id}_I]} : (I^{q+1}, \partial I^{q+1}) &\rightarrow (CX/X, *) \\ (s, t) &\mapsto f(s) \wedge (1-t) \end{aligned}$$

Since $X \hookrightarrow CX$ is cofibration and n -equivalence between $(n-1)$ -connected spaces, p is an $2n$ -equivalence. Therefore, $q+1 < 2n$ implies $-\sigma$ is isomorphism, $q+1 = 2n$ implies $-\sigma$ is epimorphism, and we have $-\sigma$ is iff σ is. □

Definition 3.1. We now define the q -th stable homotopy group:

$$\pi_k^s(X) := \text{Colim}_r \pi_{k+r}(\Sigma^r X) \cong \pi_{2k+2}(\Sigma^{k+2} X) \cong \pi_{k+n}(\Sigma^n X) \quad (n-1 > k)$$

The relation right side is directly from $\Sigma^n X$ is $(n-1)$ -connected.

Note. We'll see later that $\{\pi_n^s\}_{n \in \mathbb{N}}$ defines a generalized homology theory.