

Statistical Operations and Matrices (I)

Predictive Modeling & Statistical Learning

Gaston Sanchez

CC BY-SA 4.0

Vector-Matrix Notation for Statistical Operations

Motivation

I want to discuss how we can use vector-matrix notation to represent some basic statistical operations and summaries.

First we need to quickly review some concepts around inner products.

Inner Product

The **inner product** of two vectors \mathbf{x} and \mathbf{y} —of the same size—is defined as:

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y} = \sum_i x_i y_i$$

basically the inner product consists of the element-by-element product of \mathbf{x} and \mathbf{y} , and then adding everything up. The **result** is not another vector but a single number, a **scalar**.

Inner Product

We can also write the inner product $\langle \mathbf{x}, \mathbf{y} \rangle$ in vector notation as:

$$\mathbf{x}^T \mathbf{y} = (x_1 \dots x_n) \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} = \sum_{i=1}^n x_i y_i$$

Inner Product

Keep in mind that inner products can be generalized to any type of metric **M** matrix:

$$\langle \mathbf{x}, \mathbf{y} \rangle_M = \mathbf{x}^T \mathbf{M} \mathbf{y}$$

Inner Product

Having an inner space endowed with an inner product, we can derive other concepts such as:

- ▶ **Length** of a vector (and *norms* in general)
- ▶ **Distance** between points
- ▶ **Angle** between vectors
- ▶ **Projection** of vectors

Length

Another important usage of the inner product is that it allows us to define the **length** of a vector \mathbf{x} denoted by:

$$\|\mathbf{x}\| = \sqrt{\mathbf{x}^T \mathbf{x}}$$

which is typically known as the (Euclidean) **norm** of a vector.
(There are actually other types of norms)

Length

The inner product of a vector with itself is equal to its squared norm: $\mathbf{x}^T \mathbf{x} = \|\mathbf{x}\|^2$

$$\|\mathbf{x}\|^2 = \sum_{i=1}^n x_i^2$$

Distance

The square (Euclidean) distance between two vectors \mathbf{x} and \mathbf{y} can be obtained as:

$$d^2(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|^2 = (\mathbf{x} - \mathbf{y})^\top (\mathbf{x} - \mathbf{y})$$

Distance

The square (Eculidean) distance between two vectors \mathbf{x} and \mathbf{y} can be obtained as:

$$d^2(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|^2 = (\mathbf{x} - \mathbf{y})^\top (\mathbf{x} - \mathbf{y})$$

The distance between two vectors \mathbf{x} and \mathbf{y} can be obtained as:

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\| = \sqrt{(\mathbf{x} - \mathbf{y})^\top (\mathbf{x} - \mathbf{y})}$$

Angle between Vectors

The **angle** θ between two nonzero vectors \mathbf{x}, \mathbf{y} can also be expressed using inner products:

$$\cos(\theta) = \frac{\mathbf{x}^T \mathbf{y}}{\sqrt{\mathbf{x}^T \mathbf{x}} \sqrt{\mathbf{y}^T \mathbf{y}}} = \frac{\mathbf{x}^T \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

Angle

Equivalently, we can reexpress the formula of the inner product using

$$\mathbf{x}^T \mathbf{y} = \|\mathbf{x}\| \|\mathbf{y}\| \cos(\theta)$$

Orthogonality

Besides calculating lengths of vectors and angles between vectors, an inner product allows us to know whether two vectors are orthogonal.

In a two dimensional space, orthogonality is equivalent to perpendicularity; so if two vectors are perpendicular to each other—the angle between them is a 90 degree angle—they are orthogonal.

Two vectors \mathbf{x} and \mathbf{y} are orthogonal if their inner product is zero:

$$\mathbf{x}^T \mathbf{y} = 0 \quad \Longleftrightarrow \quad \mathbf{x} \perp \mathbf{y}$$

Projection

The last aspect I want to touch related with the inner product is the so-called projections. More specifically: orthogonal projection of a vector y onto another vector x .

The basic notion of projection requires two ingredients: two vectors, x and y . To obtain the projection of y onto x , we need to express x in unit norm.

Unit Vector

Recall that a **unit vector** is a vector of length 1, sometimes also called a *direction* or *unitary* vector.

In other words, if \mathbf{v} is a unit vector, this means that $\|\mathbf{v}\| = 1$.

Given a non-zero vector \mathbf{v} , how do we get a unit vector?

Unit Vector

Given a non-zero vector \mathbf{v} , you can get a unit vector \mathbf{u} by dividing \mathbf{v} by its norm, that is:

$$\mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{\mathbf{v}}{\sqrt{\mathbf{v}^T \mathbf{v}}}$$

Some say that we “normalize \mathbf{v} ”

Projection

Having two nonzero vectors \mathbf{x} and \mathbf{y} , we can project \mathbf{y} on \mathbf{x} , denoted by $\hat{\mathbf{y}}$ as:

$$\hat{\mathbf{y}} = \mathbf{x} \left(\frac{\mathbf{y}^T \mathbf{x}}{\mathbf{x}^T \mathbf{x}} \right)$$

Note that the term in parenthesis is just a scalar.

Projection

Note that


$$\hat{\mathbf{y}} = \mathbf{x} \left(\frac{\mathbf{y}^T \mathbf{x}}{\mathbf{x}^T \mathbf{x}} \right) = \mathbf{x} \left(\frac{\mathbf{x}^T \mathbf{y}}{\mathbf{x}^T \mathbf{x}} \right)$$

can also be written as:

$$\hat{\mathbf{y}} = \mathbf{x}(\mathbf{x}^T \mathbf{x})^{-1} \mathbf{x}^T \mathbf{y}$$

Does it look familiar?

Orthogonal Projection



Projection

We can actually express $\hat{\mathbf{y}}$ as $a\mathbf{x}$. This means that a projection implies multiplying \mathbf{x} by some number a , such that $\hat{\mathbf{y}} = a\mathbf{x}$ is a stretched or shrinked version of \mathbf{x} .

Inner Products ... So what?

We'll see in a moment how many **statistical summaries** can be represented with inner products.

Review of Some Statistical Operations

Statistical Summaries

I want to review basic statistical summaries and see how we can express them in vector-matrix notation:

- ▶ Sum of values
- ▶ Sum of squared values
- ▶ Mean
- ▶ Variance

Sum of Values

Consider a variable $X \in \mathbb{R}^n$ represented with a vector $\mathbf{x} = (x_1, x_2, \dots, x_n)$

A common operation consists of adding the elements of the vector

$$\sum_{i=1}^n x_i$$

Sum of Values

Consider a variable $X \in \mathbb{R}^n$ represented with a vector $\mathbf{x} = (x_1, x_2, \dots, x_n)$

A common operation consists of adding the elements of the vector

$$\sum_{i=1}^n x_i$$

This sum can be expressed in vector notation as:

$$\sum_{i=1}^n x_i = \mathbf{x}^\top \mathbf{1} = \mathbf{1}^\top \mathbf{x}$$

where $\mathbf{1}$ is a vector of n elements equal to 1

Mean

The (arithmetic) mean of X is

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

Mean

The (arithmetic) mean of X is

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

Using vector notation the mean is expressed as:

$$\bar{x} = \frac{1}{n} \mathbf{x}^T \mathbf{1}$$

Sum of Square dValues

Another common operation is the sum of squared values

$$\sum_{i=1}^n x_i^2$$

Sum of Square dValues

Another common operation is the sum of squared values

$$\sum_{i=1}^n x_i^2$$

Using vector notation this sum can be written as an inner product:

$$\sum_{i=1}^n x_i^2 = \mathbf{x}^T \mathbf{x} = \|\mathbf{x}\|^2$$

Variance

The variance of X (in its population version)

$$\text{var}(X) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

- ▶ Let $\bar{\mathbf{x}}$ be an n -element vector constant of \bar{x} values
- ▶ Let $\tilde{\mathbf{x}} = \mathbf{x} - \bar{\mathbf{x}}$

Variance

The variance of X (in its population version)

$$\text{var}(X) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$$

- ▶ Let $\bar{\mathbf{x}}$ be an n -element vector constant of \bar{x} values
- ▶ Let $\tilde{\mathbf{x}} = \mathbf{x} - \bar{\mathbf{x}}$

The variance can be expressed as:

$$\text{var}(X) = \frac{1}{n} \tilde{\mathbf{x}}^T \tilde{\mathbf{x}}$$

Variance

If you consider the “sample” variance

$$\text{var}(X) = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

Then:

$$\text{var}(X) = \frac{1}{n-1} \tilde{\mathbf{x}}^T \tilde{\mathbf{x}}$$

Variance

- ▶ I will sometimes consider \mathbf{x} to be mean-centered.
- ▶ This means that \bar{x} has already been subtracted from each element x_i

In this case the **variance** can be compactly expressed as:

$$\text{var}(X) = \frac{1}{n} \mathbf{x}^\top \mathbf{x}$$

Covariance

Consider two variables X and Y . The covariance $cov(X, Y)$

$$cov(X, Y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$$

Covariance

Consider two variables X and Y . The covariance $cov(X, Y)$

$$cov(X, Y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$$

Assuming that X and Y are mean-centered, and represented by \mathbf{x} and \mathbf{y} , respectively, the covariance can be expressed as:

$$cov(\mathbf{x}, \mathbf{y}) = \frac{1}{n} \mathbf{x}^T \mathbf{y}$$

Correlation

Consider two variables X and Y . The correlation $cor(X, Y)$

$$cor(X, Y) = \frac{cov(X, Y)}{\sqrt{var(X)}\sqrt{var(Y)}}$$

Correlation

Consider two variables X and Y . The correlation $cor(X, Y)$

$$cor(X, Y) = \frac{cov(X, Y)}{\sqrt{var(X)}\sqrt{var(Y)}}$$

Assuming that X and Y are mean-centered, and represented by \mathbf{x} and \mathbf{y} , respectively, the covariance can be expressed as:

$$cor(\mathbf{x}, \mathbf{y}) = \frac{\mathbf{x}^\top \mathbf{y}}{\sqrt{\mathbf{x}^\top \mathbf{x}} \sqrt{\mathbf{y}^\top \mathbf{y}}} = \frac{\mathbf{x}^\top \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

Correlation: Geometric Interpretation

What does it mean geometrically:

$$\text{cor}(\mathbf{x}, \mathbf{y}) = \frac{\mathbf{x}^\top \mathbf{y}}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

Hint: recall the angle between two vectors.

You can think of the correlation between two variables as the cosine of the angle between two vectors.