Computational biology

Co-evolution to predict protein structures

Clovis Galiez

Grenoble
Statistiques pour les sciences du Vivant et de l'Homme

December 11, 2019

Goal

- Get an overview of computational biology topics
  - Topics (genomics, metagenomics, proteomics, etc.)
  - Know some important databases
  - Know standard tools (Blast, PyMol) and libraries (BioPython)
- Have a basic culture of order of magnitude in computational biology
  - Quantity of data
  - Size of genomes
  - Size of organisms
- Toward autonomy for design and implementation of methods
  - Case study of SNP detection
  - Case study of protein structure prediction

Today's outline: from gene sequence to protein structure

- Sequence-structure-function paradigm
  - Genomes, genes, proteins
  - Databases
- Evolution
  - Selective pressure
  - Multiple sequence alignment
  - Co-evolution

From genome to function, the very big picture

ACGATGTATTCAGCGATTACGATAAAGCTACGTAGTGGCA

On a genome (∼5Mbp), specific motifs define begining and end of a gene

From genome to function, the very big picture



Transcription + *translation*, to form a chain of amino acids (∼300-3000AA)

From genome to function, the very big picture



Protein folding under pysico-chemical interactions, diameter $\sim$ few nanometers

From genome to function, the very big picture



Protein endowed with a function (biochemical reactions, transport, etc.)

Data at every steps

Nucleic seq. | Amino acid seq. | Protein | Function

..ATTGTCGATGAC..





Data at every steps

Nucleic seq.	Amino acid seq.	Protein	Function
..ATTGTCGATGAC..			
NCBI	UniProt	RCSB PDB PROTEIN DATA BANK	-
`ncbi.nlm.nih.gov`	`uniprot.org`	`rcsb.org`	-

Protein evolution through mutations



Protein evolution through mutations



Sequence conservation

Aligning the sequences (MSA, multiple sequence alignment):

Tools	Database
ClustalW [Larkin et al. 07]	Pfam `pfam.xfam.org`

Sequence conservation

Aligning the sequences (MSA, multiple sequence alignment):

Tools	Database
ClustalW [Larkin et al. 07]	Pfam `pfam.xfam.org`

Why some positions are conserved, some other aren't?

Structure is determined by amino acid interactions

Preserving the function: coevolution of residues

As protein function is vital, **evolution selects mutations preserving structures**.
Leading to **compensatory** mutations:



Computers and protein structure prediction

(cheap)

↓

ACGATGTATTCAGCGATTACGATAAAGCTACGTAGTGGCA



← (expensive)

Structure determined by X-Rays
through a cristal of proteins



A simple approach for protein structure prediction



- Build or get multiple manio acid sequence alignments (**e.g.** in Pfam database)
- Quantify coevolution between positions in the sequence
- Infer what are the position in contact

What measure for co-evolution? Correlation would work?

Conservation vs. co-evolution

Conserved position carries no information in terms of co-evolution (entropy is zero).

Conservation vs. co-evolution

Conserved position carries no information in terms of co-evolution (entropy is zero). A standard approach is to measure it through Mutual Information:

$$MI(i,j) = \sum_{a,b} p(x_i = a, x_j = b) \log \frac{p(x_i=a, x_j=b)}{p(x_i=a)\, p(x_j=b)}$$

Where

- x_i is the amino acid at position i
- $p(x_i = a)$ is estimated in the MSA by $\frac{\#\text{sequences having "a" at position } i}{N}$
- N the number of sequences in the MSA
- $p(x_i = a, x_j = b)$ is estimated in the MSA by $\frac{\#\text{sequence having "a" at } i \text{ and "b" at } j}{N}$

In paractice you need $N > 1,000$ to have reasonable estimation of $p(x_i = a, x_j = b)$.

Over-repdiction at entropic position

When applying the rule

$$MI(i,j) > \tau \Rightarrow \text{contact between } i \text{ and } j$$

some positions predict too many contacts, often position with high entropy. Several corrections can be applied[1].

In your project

You can try using the simple correction:

$$MI'(i,j) = MI(i,j) - \frac{1}{N}\sum_k (MI(k,j) + MI(i,k))$$

and fix a τ to predict a contact as soon as:

$$MI'(i,j) > \tau$$

[1] See `https://doi.org/10.1093/bioinformatics/bti671`

Hints concerning your Salmonella project

- The mutated gene you identified in the resistant bacteria have a large multiple sequence alignment in Pfam, search for its name in the Pfam browser.
- WP1.T2 consists in creating a tool taking as input an MSA in Fasta format and outputing a **contact matrix**[2].
- You can model the structure using FT-comar[3] software, and compare to the native structure [4] using RasMol or PyMol software:



[2] see the *cheatsheet* for details about the file format

[3] `clovisg.github.io/teaching/protein-structure-prediction/ft-comar.tgz`

[4] `clovisg.github.io/teaching/protein-structure-prediction/target.pdb`

Summary

Check what you've learn:

- What is a genome, a gene, a protein, its structure
- How real sequencing data look like
- What is a SNP, what can be the impact
- Main tools and databases in computational biology
- Potential application of computational biology for public health studies

The project involved basic skills from different area:

- biology
- statistics (Poisson distribution)
- algorithmics (linear time algorithms required)

Projects

Remember that your project should be like professional answers to the call:

- Clarity
- Fulfillment of the call
- Trustworthiness in the description of the approach

You should send:

- a report, including:
  - description of the strategy
  - approximations and choices
  - application to the project data (what gene is impacted by the SNP)
- your code
- a step-by-step guide to reproduce the results of the report

The TATFAR
waits for
interesting answers to its call!