Packing Problems & Number Theory

> Clyde Kertzer

### Packing Problems & Number Theory

Clyde Kertzer

University of Colorado Boulder

May 7, 2025

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

> > $\ \ \, \text{A guessing jar:}$

Packing Problems & Number Theory

> Clyde Kertzer

> > A guessing jar: How many marbles?

Packing Problems & Number Theory

> Clyde Kertzer

> > A guessing jar: How many marbles? 223!

Packing Problems & Number Theory

> Clyde Kertzer

> > A guessing jar: How many marbles? 223!

1. What's the largest number of marbles that can fit in the container?

Packing Problems & Number Theory

> Clyde Kertzer

> > A guessing jar: How many marbles? 223!

- 1. What's the largest number of marbles that can fit in the container?
- 2. Given 223 marbles, what's the smallest container holding them all?

Packing Problems & Number Theory


> Clyde Kertzer

> > A guessing jar: How many marbles? 223!

- 1. What's the largest number of marbles that can fit in the container?
- 2. Given 223 marbles, what's the smallest container holding them all? We need to simplify the problem...

Packing Problems & Number Theory

Packing Problems & Number Theory



Packing Problems & Number Theory

Packing Problems & Number Theory


> Clyde Kertzer

What's the smallest square we can fit 4 squares inside of?

Packing Problems & Number Theory

> Clyde Kertzer


What's the smallest square we can fit 4 squares inside of?



Packing Problems & Number Theory

> Clyde Kertzer

What's the smallest square we can fit 4 squares inside of?



Side length: 2

Packing Problems & Number Theory

Packing Problems & Number Theory


> Clyde Kertzer

What's the smallest square we can fit 9 squares inside of?

Packing Problems & Number Theory

> Clyde Kertzer


What's the smallest square we can fit 9 squares inside of?



Packing Problems & Number Theory

> Clyde Kertzer

What's the smallest square we can fit 9 squares inside of?



Side length: 3


Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer


Packing Problems & Number Theory

> Clyde Kertzer



Packing Problems & Number Theory

> Clyde Kertzer




Side length:

Packing Problems & Number Theory


> Clyde Kertzer

#### What about 5 squares?



Side length:  $\approx 2.707...$ 


Clyde Kertzer



Side length: 
$$\approx 2.707... = 2 + \frac{\sqrt{2}}{2}$$

Clyde Kertzer

#### What about 5 squares?




Side length: 
$$\approx 2.707... = 2 + \frac{\sqrt{2}}{2}$$


Can we use this packing to find the optimal packing of 10 squares?

Packing Problems & Number Theory

Packing Problems & Number Theory



Packing Problems & Number Theory



Side length: 
$$pprox 3.707... = 3 + \frac{\sqrt{2}}{2}$$


Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer


Packing Problems & Number Theory

> Clyde Kertzer



Packing Problems & Number Theory

> Clyde Kertzer




Side length:  $\approx 3.877\dots$ 

Packing Problems & Number Theory

> Clyde Kertzer

#### What about 11 squares?




Side length:  $\approx 3.877\dots$ 

Is this the best possible packing?

Packing Problems & Number Theory

> Clyde Kertzer

#### What about 11 squares?



Side length:  $\approx 3.877...$ 


Is this the best possible packing? Mathematicians still don't know...

# Circle Packing


Packing Problems & Number Theory

# Circle Packing

Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory

> Clyde Kertzer




Try it yourself: Can you find an arrangement of 4 circles such that each one is tangent to the other?

Packing Problems & Number Theory



Packing Problems & Number Theory


> Clyde Kertzer




What if the circles aren't all the same size...

Packing Problems & Number Theory


Packing Problems & Number Theory



Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory

> Clyde Kertzer



#### Definition


A set of four mutually tangent circles is called a Descartes Quadruple

Packing Problems & Number Theory


Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory Clyde

Kertzer




With your set can you make a circle packing that is not symmetric?

Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory


Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory



Packing Problems & Number Theory



### Summary: The Descartes Equation

Packing Problems & Number Theory

### Summary: The Descartes Equation

Packing Problems & Number Theory

Definition

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition


The *curvature* of a circle with radius r is defined to be 1/r.

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition

The *curvature* of a circle with radius r is defined to be 1/r.




Packing Problems & Number Theory

> Clyde Kertzer

#### Definition

The *curvature* of a circle with radius r is defined to be 1/r.




Circle with infinite radius (Curvature 0)

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition

The *curvature* of a circle with radius r is defined to be 1/r.



Circle with infinite radius (Curvature 0)


### Definition

Packing Problems & Number Theory

> Clyde Kertzer

### Definition

The *curvature* of a circle with radius r is defined to be 1/r.



Circle with infinite radius (Curvature 0)

#### Definition


Four mutually tangent circles are called a Descartes Quadruple.

Packing Problems & Number Theory

> Clyde Kertzer

### Definition

The *curvature* of a circle with radius r is defined to be 1/r.



Circle with infinite radius (Curvature 0)

#### Definition

Four mutually tangent circles are called a Descartes Quadruple.

### **Descartes Equation**

Packing Problems & Number Theory

> Clyde Kertzer

### Definition

The *curvature* of a circle with radius r is defined to be 1/r.



Circle with infinite radius (Curvature 0)

#### Definition

Four mutually tangent circles are called a Descartes Quadruple.

### **Descartes Equation**


If four mutually tangent circles have curvatures a, b, c, d then

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition

The *curvature* of a circle with radius r is defined to be 1/r.



Circle with infinite radius (Curvature 0)

#### Definition

Four mutually tangent circles are called a Descartes Quadruple.


## Descartes Equation

If four mutually tangent circles have curvatures a, b, c, d then

$$(a+b+c+d)^2 = 2(a^2+b^2+c^2+d^2).$$


Packing Problems & Number Theory

Packing Problems & Number Theory




Packing Problems & Number Theory

Kertzer



### Corollary


If three mutually tangent circles have curvatures a, b, and c, then the two circles of Apollonius, d and d' have curvatures



### Corollary

If three mutually tangent circles have curvatures a, b, and c, then the two circles of Apollonius, d and d' have curvatures

$$d = a + b + c + 2\sqrt{ab + ac + bc}$$
$$d' = a + b + c - 2\sqrt{ab + ac + bc}$$




### Corollary

If three mutually tangent circles have curvatures a, b, and c, then the two circles of Apollonius, d and d' have curvatures

$$d = a + b + c + 2\sqrt{ab + ac + bc}$$
$$d' = a + b + c - 2\sqrt{ab + ac + bc}$$

#### The Key Relation



### Corollary

If three mutually tangent circles have curvatures a, b, and c, then the two circles of Apollonius, d and d' have curvatures


$$d = a + b + c + 2\sqrt{ab + ac + bc}$$
$$d' = a + b + c - 2\sqrt{ab + ac + bc}$$

### The Key Relation

$$d+d'=2(a+b+c)$$

Packing Problems & Number Theory

Kertzer



### Corollary


If three mutually tangent circles have curvatures a, b, and c, then the two circles of Apollonius, d and d' have curvatures

$$d = a + b + c + 2\sqrt{ab + ac + bc}$$
$$d' = a + b + c - 2\sqrt{ab + ac + bc}$$


### The Key Relation

$$d + d' = 2(a + b + c) \implies d' = 2(a + b + c) - d$$


Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory




Packing Problems & Number Theory



[-6, 11, 14, 23]


Packing Problems & Number Theory



[-6, 11, 14, 23] reduces to [-6, 11, 14, 15]

Packing Problems & Number Theory

> Clyde Kertzer



[-6, 11, 14, 15]

Packing Problems & Number Theory

Kertzer

[-6, 11, 14, 15]

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

Definition

Packing Problems & Number Theory

> Clyde Kertzer

### Definition

A positive integer a has a packing

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition

A positive integer a has a packing if there exists a primitive reduced Descartes quadruple [-a, b, c, d].

Packing Problems & Number Theory

> Clyde Kertzer

#### Definition


A positive integer a has a packing if there exists a primitive reduced Descartes quadruple [-a, b, c, d].

How many other packings does  $\it a=7$  have? How many of them are symmetric?

### Definition

A positive integer a has a packing if there exists a primitive reduced Descartes quadruple [-a, b, c, d].

How many other packings does a=7 have? How many of them are symmetric?




[-7, 12, 17, 20],

#### Definition

A positive integer a has a packing if there exists a primitive reduced Descartes quadruple [-a, b, c, d].

How many other packings does a = 7 have? How many of them are symmetric?




Clyde Kertzer

#### Definition


A positive integer a has a packing if there exists a primitive reduced Descartes quadruple [-a, b, c, d].

How many other packings does a=7 have? How many of them are symmetric?




Packing Problems & Number Theory

Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory


Clyde Kertzer




Definition

Packing Problems & Number Theory Clyde

Kertzer







#### Definition

A sum-symmetric

Packing Problems & Number Theory Clyde

Kertzer







#### Definition

A *sum-symmetric* quadruple is a primitive reduced Descartes quadruple satisfying 2(a+b+c)-d=d.

Packing Problems & Number Theory

Kertzer






#### Definition


A *sum-symmetric* quadruple is a primitive reduced Descartes quadruple satisfying 2(a + b + c) - d = d.

$$2(a+b+c)-d=d$$

Packing Problems & Number Theory

Kertzer






#### Definition


A sum-symmetric quadruple is a primitive reduced Descartes quadruple satisfying 2(a+b+c)-d=d.

$$2(a+b+c)-d=d \implies 2(a+b+c)=2d$$

Packing Problems & Number Theory

Kertzer






#### Definition

A sum-symmetric quadruple is a primitive reduced Descartes quadruple satisfying 2(a+b+c)-d=d.


$$2(a+b+c)-d=d \implies 2(a+b+c)=2d \implies a+b+c=d$$

Packing Problems & Number Theory


Packing Problems & Number Theory




Packing Problems & Number Theory



Packing Problems & Number Theory




[-4, 8, 9, 9]




Definition

Packing Problems & Number Theory Clyde

Kertzer







#### Definition

A twin-symmetric quadruple

Packing Problems & Number Theory Clyde

Kertzer






#### Definition


A *twin-symmetric* quadruple is a primitive reduced Descartes quadruple with c=d or c=b.

Packing Problems & Number Theory

Packing Problems & Number Theory




Packing Problems & Number Theory




The strip packing:  $\left[0,0,1,1\right]$ 

Packing Problems & Number Theory

Packing Problems & Number Theory



Packing Problems & Number Theory



The bug-eye packing:  $\left[-1,2,2,3\right]$ 

Packing Problems & Number Theory

> Clyde Kertzer

> > Proposition (Kertzer, 2024)

Packing Problems & Number Theory

> Clyde Kertzer

#### Proposition (Kertzer, 2024)

A symmetric packing is either sum-symmetric or twin-symmetric.

Packing Problems & Number Theory

> Clyde Kertzer

#### Proposition (Kertzer, 2024)

 $\label{lem:approx} A \ \textit{symmetric packing is either sum-symmetric or twin-symmetric}.$ 

Proposition (Kertzer, 2024)

Packing Problems & Number Theory

> Clyde Kertzer

#### Proposition (Kertzer, 2024)

A symmetric packing is either sum-symmetric or twin-symmetric.

#### Proposition (Kertzer, 2024)

Only the strip and bug-eye packing are both sum-symmetric and twin-symmetric.

Packing Problems & Number Theory

Packing Problems & Number Theory

$$[-a,b,c,d]$$
  $d-c$   $d-b$   $d+a$ 

Packing Problems & Number Theory

$$\frac{ [-a,b,c,d] \quad | \ d-c \ | \ d-b \ | \ d+a }{ [-6,10,15,19] \ | }$$

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	4	9	25

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	4	9	25
[-12, 21, 28, 37]	9	16	49

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	4	9	25
[-12, 21, 28, 37]	9	16	49
[-18, 22, 99, 103]	4	81	121

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	4	9	25
[-12, 21, 28, 37]	9	16	49
[-18, 22, 99, 103]	4	81	121
[-20, 36, 45, 61]	16	25	81

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	4	9	25
[-12, 21, 28, 37]	9	16	49
[-18, 22, 99, 103]	4	81	121
[-20, 36, 45, 61]	16	25	81
[-21, 30, 70, 79]	9	49	100

Packing Problems & Number Theory

[-a,b,c,d]	d – c	d-b	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	$2^2$	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	5 <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	7 <sup>2</sup>	10 <sup>2</sup>

Packing Problems & Number Theory Clyde

Kertzer

[-a,b,c,d]	d-c	b+a	d-b	c + a	d + a
[-6, 10, 15, 19]	$2^{2}$		3 <sup>2</sup>		5 <sup>2</sup>
[-12, 21, 28, 37]	$3^2$		<b>4</b> <sup>2</sup>		<b>7</b> <sup>2</sup>
[-18, 22, 99, 103]	$2^2$		9 <sup>2</sup>		$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>		5 <sup>2</sup>		9 <sup>2</sup>
[-21, 30, 70, 79]	$3^2$		7 <sup>2</sup>		10 <sup>2</sup>

Packing Problems & Number Theory

[-a,b,c,d]	d-c	b+a	d-b	c + a	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	2 <sup>2</sup>	3 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	2 <sup>2</sup>	9 <sup>2</sup>	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	5 <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	3 <sup>2</sup>	<b>7</b> <sup>2</sup>	7 <sup>2</sup>	10 <sup>2</sup>

Packing Problems & Number Theory Clyde

Kertzer

[-a,b,c,d]	d-c	b + a	d-b	c + a	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	2 <sup>2</sup>	3 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>7</b> <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	2 <sup>2</sup>	9 <sup>2</sup>	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	3 <sup>2</sup>	7 <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

Given the factorization of a, we can find the entire quadruple!

Packing Problems & Number Theory

Kertzer

[-a,b,c,d]	d-c	b+a	d-b	c + a	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	2 <sup>2</sup>	3 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	2 <sup>2</sup>	9 <sup>2</sup>	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	5 <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	3 <sup>2</sup>	7 <sup>2</sup>	7 <sup>2</sup>	10 <sup>2</sup>

Given the factorization of a, we can find the entire quadruple!

$$\left[\underbrace{-(2\cdot 3)}_{-6},\,\underbrace{2^2+2\cdot 3}_{10},\,\underbrace{3^2+2\cdot 3}_{15},\,\underbrace{(2+3)^2-2\cdot 3}_{19}\right]$$

Packing Problems & Number Theory

Kertzer

[-a,b,c,d]	d-c	b+a	d-b	c + a	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	2 <sup>2</sup>	3 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	2 <sup>2</sup>	9 <sup>2</sup>	$9^{2}$	$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	5 <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	3 <sup>2</sup>	7 <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

Given the factorization of a, we can find the entire quadruple!

$$\left[\underbrace{-(2\cdot 3)}_{-6}, \underbrace{2^2+2\cdot 3}_{10}, \underbrace{3^2+2\cdot 3}_{15}, \underbrace{(2+3)^2-2\cdot 3}_{19}\right]$$

$$[-xy, x^2 + xy, y^2 + xy, (x + y)^2 - xy]$$

Packing Problems & Number Theory

Kertzer

[-a,b,c,d]	d-c	b+a	d-b	c + a	d + a
[-6, 10, 15, 19]	2 <sup>2</sup>	2 <sup>2</sup>	3 <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	3 <sup>2</sup>	3 <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	2 <sup>2</sup>	9 <sup>2</sup>	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	5 <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	3 <sup>2</sup>	3 <sup>2</sup>	<b>7</b> <sup>2</sup>	7 <sup>2</sup>	10 <sup>2</sup>

Given the factorization of a, we can find the entire quadruple!

$$\left[\underbrace{-(2\cdot 3)}_{-6}, \underbrace{2^2+2\cdot 3}_{10}, \underbrace{3^2+2\cdot 3}_{15}, \underbrace{(2+3)^2-2\cdot 3}_{19}\right]$$

$$[-xy, x^2 + xy, y^2 + xy, (x + y)^2 - xy]$$

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy]$$

Packing Problems & Number Theory

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	<b>2</b> <sup>2</sup>	<mark>9</mark> 2	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

Packing Problems & Number Theory

Clyde Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>7</b> <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	<mark>9</mark> 2	$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<mark>7</mark> 2	10 <sup>2</sup>

Packing Problems & Number Theory

Clyde Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>7</b> <sup>2</sup>
[-18, 22, 99, 103]	<b>2</b> <sup>2</sup>	9 <sup>2</sup>	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy] =$$

Packing Problems & Number Theory

Clyde Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	3 <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	<mark>9</mark> 2	11 <sup>2</sup>
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy] =$$

$$\left[-2\cdot 6,\ 2(2+6),\ 6(2+6),\ (2+6)^2-2\cdot 6\right]=$$

Packing Problems & Number Theory

Clyde Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	<b>7</b> <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	<mark>9</mark> 2	$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy] =$$

$$[-2 \cdot 6, 2(2+6), 6(2+6), (2+6)^2 - 2 \cdot 6] =$$

$$[-12, 16, 48, 52]$$

Packing Problems & Number Theory

Clyde Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	7 <sup>2</sup>
[-18, 22, 99, 103]	2 <sup>2</sup>	<mark>9</mark> 2	$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	10 <sup>2</sup>

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy] =$$

$$[-2\cdot 6, 2(2+6), 6(2+6), (2+6)^2 - 2\cdot 6] =$$

$$[-12, 16, 48, 52] = [-3, 4, 12, 13]$$

Packing Problems & Number Theory

Kertzer

[-a,b,c,d]	d-c	d-b	d + a
[-6, 10, 15, 19]	<b>2</b> <sup>2</sup>	<b>3</b> <sup>2</sup>	5 <sup>2</sup>
[-12, 21, 28, 37]	<b>3</b> <sup>2</sup>	<b>4</b> <sup>2</sup>	$7^{2}$
[-18, 22, 99, 103]	<b>2</b> <sup>2</sup>	<mark>9</mark> 2	$11^{2}$
[-20, 36, 45, 61]	<b>4</b> <sup>2</sup>	<b>5</b> <sup>2</sup>	9 <sup>2</sup>
[-21, 30, 70, 79]	<b>3</b> <sup>2</sup>	<b>7</b> <sup>2</sup>	$10^{2}$

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy] =$$

$$[-2 \cdot 6, 2(2+6), 6(2+6), (2+6)^2 - 2 \cdot 6] =$$

$$[-12, 16, 48, 52] = [-3, 4, 12, 13]$$
  $(x = 3, y = 1)$ 

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

> > Theorem (Kertzer, 2024)

Packing Problems & Number Theory

> Clyde Kertzer

#### Theorem (Kertzer, 2024)

A sum-symmetric quadruple [a,b,c,d] is of the form

Packing Problems & Number Theory

> Clyde Kertzer

#### Theorem (Kertzer, 2024)

A sum-symmetric quadruple [a, b, c, d] is of the form

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy]$$

Packing Problems & Number Theory

> Clyde Kertzer

#### Theorem (Kertzer, 2024)

A sum-symmetric quadruple [a, b, c, d] is of the form

$$[-xy, x(x+y), y(x+y), (x+y)^2 - xy]$$

with gcd(x, y) = 1, and  $x, y \geqslant 0$ .

Packing Problems & Number Theory

Packing Problems & Number Theory



Packing Problems & Number Theory

> Clyde Kertzer

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

Packing Problems & Number Theory

> Clyde Kertzer

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Packing Problems & Number Theory

> Clyde Kertzer

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Because n = -xy determines the sum-symmetric packing for coprime x and y, write  $n = p_1^{e_1} p_2^{e_2} \cdots p_{\nu}^{e_k}$ , so  $\omega(n) = k$ .

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Because n=-xy determines the sum-symmetric packing for coprime x and y, write  $n=p_1^{e_1}p_2^{e_2}\cdots p_k^{e_k}$ , so  $\omega(n)=k$ . For each prime power we can choose to put it as a factor of x or y,

Clyde Kertzer

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Because n=-xy determines the sum-symmetric packing for coprime x and y, write  $n=p_1^{e_1}p_2^{e_2}\cdots p_k^{e_k}$ , so  $\omega(n)=k$ . For each prime power we can choose to put it as a factor of x or y, so there  $2^k$  total factor pairs xy

Clyde Kertzer

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Because n=-xy determines the sum-symmetric packing for coprime x and y, write  $n=p_1^{e_1}p_2^{e_2}\cdots p_k^{e_k}$ , so  $\omega(n)=k$ . For each prime power we can choose to put it as a factor of x or y, so there  $2^k$  total factor pairs xy but we divide by two to account for symmetry.

#### Corollary

A natural number n has  $2^{\omega(n)-1}$  sum-symmetric packings, where  $\omega(n)$  is the number of distinct prime divisors of n.

#### Proof.

Because n=-xy determines the sum-symmetric packing for coprime x and y, write  $n=p_1^{e_1}p_2^{e_2}\cdots p_k^{e_k}$ , so  $\omega(n)=k$ . For each prime power we can choose to put it as a factor of x or y, so there  $2^k$  total factor pairs xy but we divide by two to account for symmetry. Thus, n has  $2^k/2=2^{k-1}=2^{\omega(n)-1}$  sum-symmetric packings.  $\square$ 

Packing Problems & Number Theory

Packing Problems & Number Theory

Write 
$$60 = 2^2 \cdot 3 \cdot 5$$
,

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5,$  so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings.

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs (1,60),

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs (1,60), (4,15),

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs (1,60), (4,15), (3,20),

Packing Problems & Number Theory

> Clyde Kertzer

> > Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs (1,60), (4,15), (3,20), (5,12).

Clyde Kertzer

Write  $60=2^2\cdot 3\cdot 5$ , so 60 has  $2^{3-1}=2^2=4$  sum-symmetric packings. These correspond to the coprime factor pairs (1,60), (4,15), (3,20), (5,12). They are

$$(1,60) \implies [-60,61,3660,3661]$$

$$(4,15) \implies [-60,76,285,301]$$

$$(3,20) \implies [-60,69,460,469]$$

$$(5,12) \implies [-60,85,204,229]$$

Packing Problems & Number Theory

Packing Problems & Number Theory


> Clyde Kertzer

Packings where one of the numbers is the same:

Packing Problems & Number Theory

> Clyde Kertzer

Packings where one of the numbers is the same:



[-4, 8, 9, 9]

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

-2 none

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
<b>-7</b>	[-7, 9, 32, 32]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
-7	[-7, 9, 32, 32]
-8	[-8, 12, 25, 25]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
<b>-7</b>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<b>-9</b>	[-9, 11, 50, 50]

Packing Problems & Number Theory

none
[-3, 5, 8, 8]
[-4, 8, 9, 9]
[-5, 7, 18, 18]
none
[-7, 9, 32, 32]
[-8, 12, 25, 25]
[-9, 11, 50, 50]
none

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
<b>-7</b>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
-9	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
<b>-7</b>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<u>-9</u>	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
<del>-6</del>	none
<del>-7</del>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
-9	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]
-13	[-13, 15, 98, 98]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
<del>-6</del>	none
<b>-7</b>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<u>-9</u>	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]
-13	[-13, 15, 98, 98]
-14	none

Packing Problems & Number Theory

-2	none
<del>-3</del>	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
<del>-6</del>	none
<del>-7</del>	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<b>-9</b>	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]
-13	[-13, 15, 98, 98]
-14	none
-15	[-15, 17, 128, 128], [-15, 32, 32, 33]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
-7	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<u>-9</u>	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]
-13	[-13, 15, 98, 98]
-14	none
-15	[-15, 17, 128, 128], [-15, 32, 32, 33]
-16	[-16, 20, 81, 81]

Packing Problems & Number Theory

-2	none
-3	[-3, 5, 8, 8]
-4	[-4, 8, 9, 9]
-5	[-5, 7, 18, 18]
-6	none
-7	[-7, 9, 32, 32]
<del>-8</del>	[-8, 12, 25, 25]
<u>-9</u>	[-9, 11, 50, 50]
-10	none
-11	[-11, 13, 72, 72]
-12	[-12, 16, 49, 49], [-12, 25, 25, 28]
-13	[-13, 15, 98, 98]
-14	none
-15	[-15, 17, 128, 128], [-15, 32, 32, 33]
-16	[-16, 20, 81, 81]

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

Over the summer:

Packing Problems & Number Theory

> Clyde Kertzer

Over the summer:



Kertzer

Over the summer:

#### **Theorem**

All primitive ACPs with c = d are given by

$$\left[-x, x+y^2, \left(\frac{2x+y^2}{2y}\right)^2, \left(\frac{2x+y^2}{2y}\right)^2\right] \quad \text{y even}$$

$$\left[ -x, \ x + 2y^2, \ 2\left(\frac{x+y^2}{2y}\right)^2, \ 2\left(\frac{x+y^2}{2y}\right)^2 \right]$$
 y odd

Kertzer

Over the summer:

#### Theorem

All primitive ACPs with c = d are given by

$$\begin{bmatrix} -x, x+y^2, \left(\frac{2x+y^2}{2y}\right)^2, \left(\frac{2x+y^2}{2y}\right)^2 \end{bmatrix} \quad \text{y even}$$

$$\begin{bmatrix} -x, x+2y^2, 2\left(\frac{x+y^2}{2y}\right)^2, 2\left(\frac{x+y^2}{2y}\right)^2 \end{bmatrix} \quad \text{y odd}$$

Not ideal, not in terms of factorization.

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

Improved to:

Packing Problems & Number Theory

> Clyde Kertzer

 $Improved\ to:$ 



Improved to:

#### **Theorem**

A twin-symmetric quadruple is of the form

$$\left\{ \left[ -xy, \, xy + 2y^2, \, \frac{(x+y)^2}{2}, \, \frac{(x+y)^2}{2} \right] \, x \, odd, \, y \, odd \, x > y \right. \right\}$$

Kertzer

Improved to:

#### **Theorem**

A twin-symmetric quadruple is of the form

$$\left\{ \begin{bmatrix}
-xy, & xy + 2y^2, & \frac{(x+y)^2}{2}, & \frac{(x+y)^2}{2} \\
-xy, & xy + 4y^2, & \left(\frac{x}{2} + y\right)^2, & \left(\frac{x}{2} + y\right)^2
\end{bmatrix} \quad x \text{ odd, } y \text{ odd} \quad x > y$$

Kertzer

Improved to:

#### **Theorem**

A twin-symmetric quadruple is of the form

$$\begin{cases}
 -xy, xy + 2y^2, \frac{(x+y)^2}{2}, \frac{(x+y)^2}{2} \\
 -xy, xy + 4y^2, \left(\frac{x}{2} + y\right)^2, \left(\frac{x}{2} + y\right)^2 \\
 -xy, xy + x^2, \left(\frac{x}{2} + y\right)^2, \left(\frac{x}{2} + y\right)^2 \\
 -xy, xy + x^2, \left(\frac{x}{2} + y\right)^2, \left(\frac{x}{2} + y\right)^2 \\
 -xy, xy + x^2, \left(\frac{x}{2} + y\right)^2, \left(\frac{x}{2} + y\right)^2
\end{cases}$$

$$4 \mid x, \quad x < 2y$$

with gcd(x, y) = 1.

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

Further improved to:

Kertzer

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Kertzer

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Ex: 
$$x = 3$$
,  $y = 2$ 

Problems & Clvde Kertzer

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Ex: 
$$x = 3$$
,  $y = 2$ :  $[-12, 12 + 16, 5^2, 5^2] \implies [-12, 28, 25, 25]$ 

Kertzer

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Ex: 
$$x = 3$$
,  $y = 2$ :  $[-12, 12 + 16, 5^2, 5^2] \implies [-12, 28, 25, 25]$ 

Why won't x = 1, y = 3 work?

Kertzer

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Ex: 
$$x = 3$$
,  $y = 2$ :  $[-12, 12 + 16, 5^2, 5^2] \implies [-12, 28, 25, 25]$ 

Why won't x = 1, y = 3 work? Let's try:

Further improved to:

### Theorem (Kertzer, 2024)

A twin-symmetric quadruple is one of following two forms

$$\begin{cases} \left[ -xy, \ xy + 2y^2, \ \frac{1}{2}(x+y)^2, \ \frac{1}{2}(x+y)^2 \right] & x, y \ odd & x > y \\ \left[ -2xy, \ 2xy + 4y^2, \ (x+y)^2, \ (x+y)^2 \right] & xy \ even & x > y \end{cases}$$

with gcd(x, y) = 1 and  $x, y \ge 0$ .

Ex: 
$$x = 3$$
,  $y = 2$ :


$$[-12, 12+16, 5^2, 5^2] \implies [-12, 28, 25, 25]$$

Why won't x = 1, y = 3 work? Let's try:

$$[-3, 3+2(3)^2, 5^2, 5^2] \implies [-12, 48, 25, 25]$$

Packing Problems & Number Theory


Packing Problems & Number Theory



[-12, 48, 25, 25]

Packing Problems & Number Theory

> Clyde Kertzer



 $[-12,48,25,25] \implies [-12,28,25,25]$ 

Packing Problems & Number Theory

Packing Problems & Number Theory

> Clyde Kertzer

> > We define  $\delta_n$  as

Packing Problems & Number Theory

> Clyde Kertzer

> > We define  $\delta_n$  as

$$\delta_n = \begin{cases} 1 & n \equiv 2 \bmod 4 \\ 0 & \text{otherwise.} \end{cases}$$

Packing Problems & Number Theory

> Clyde Kertzer

> > We define  $\delta_n$  as

$$\delta_n = \begin{cases} 1 & n \equiv 2 \bmod 4 \\ 0 & \text{otherwise.} \end{cases}$$

### Corollary

Kertzer

We define  $\delta_n$  as

$$\delta_n = \begin{cases} 1 & n \equiv 2 \bmod 4 \\ 0 & \text{otherwise.} \end{cases}$$


### Corollary

A natural number n has  $(1 - \delta_n) \cdot 2^{\omega(n)-1}$  twin-symmetric packings where  $\omega(n)$  is the number of distinct prime divisors of n.

### Thank You!

Packing Problems & Number Theory

> Clyde Kertzer



Images generated using James Rickards' Code.

### Circles checklist:

Packing Problems & Number Theory

> Clyde Kertzer

> > -7 (outer ring), 8, 9, 10, 11, 12, 15, 18, 19, 20, 22, 24, 25, 28, 32 (x2), 33, 35, 38, 41, 44, 48, 56 (x2), 57, 64, 72, 80, 88, 96, 105 (x3)

### References

Packing Problems & Number Theory

Clyde Kertzer

Tabes Bridges, Warren Tai, and Karol Koziol. A classification of integral apollonian circle packings. https://www.math.columbia.edu/-staff/RTGPapers/IntegralClassification.pdf, 2011.

Duncan Buell. Binary Quadratic Forms. Springer-Verlag, Berlin, 1989.

David A. Cox. Primes of the form  $x^2 + ny^2$ . Pure and Applied Mathematics (Hoboken). John Wiley & Sons, Inc., Hoboken, N.J, second edition, 2013. Fermat, class field theory, and complex multiplication.

Ronald L. Graham, Jeffrey C. Lagarias, Colin L. Mallows, Allan R. Wilks, and Catherine H. Yan. Apollonian Circle Packings: Number Theory. J. Number Theory, 100(1):1-45, 2003.

James Rickards. Apollonian.

https://github.com/JamesRickards-Canada/Apollonian, 2023.