

**MODERN ALGEBRA II SPRING 2013:
FOURTH PROBLEM SET**

1. Let R be a ring with $R \neq \{0\}$. Show that R is a field \iff every ideal of R is either $\{0\}$ or R . (Note: we have already showed \implies . To see \impliedby , let $r \in R$, $r \neq 0$. What can you say about the principal ideal (r) ?)
2. Let F be a field and let $\rho: F \rightarrow R$ be a ring homomorphism. Show that either ρ is injective or $R = \{0\}$ and hence $\rho(a) = 0$ for all $a \in F$.
3. Define the *ideal sum* $I + J$ to be the subset $\{r + s : r \in I, s \in J\}$. In other words, $I + J$ contains all sums of two elements, one of which is in I and the other is in J . Show that $I + J$ is an ideal containing I and J , and in fact every ideal of R containing both I and J must contain $I + J$.
4. Define the *ideal product* $I \cdot J$ to be the subset

$$\left\{ \sum_{i=1}^n r_i s_i : r_i \in I, s_i \in J \right\}.$$

In other words, $I \cdot J$ contains all finite sums of products of two elements, the first of which is in I and the second of which is in J . (There is no condition on the number n of such products we add up; it could be arbitrarily large.) Show that $I \cdot J$ is an ideal contained in $I \cap J$. (As we shall see, it is not always the case that $I \cdot J = I \cap J$.)

5. In the ring \mathbb{Z} , show that $r \in (n)$ if and only if n divides r . Describe concretely the ideals $(n) + (m)$; $(n) \cap (m)$; $(n) \cdot (m)$. In particular, give an example of two ideals I and J in \mathbb{Z} such that $I \cdot J \neq I \cap J$.
6. By Part (i) of Problem 6 from the last problem set, we know that, if R is a subring of S and J is an ideal in S , then $I = R \cap J$ is an ideal in R . Show that, in this case, there is an injective homomorphism f from R/I to S/J , defined by $f(r + I) = r + J$. (Define f first as a homomorphism from R to S/J and then determine its kernel and use the fundamental theorem for homomorphisms.) Show that f is surjective \iff for every $s \in S$, there exists an $r \in R$ such that $s \equiv r \pmod{J}$, i.e. $s - r \in J$.
7. Let R be the subring $\mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\}$ of \mathbb{C} . Let $I = (2 + 3i)$ be the principal ideal in $\mathbb{Z}[i]$ generated by $2 + 3i$.

- (i) Show that I contains $2 + 3i$ and $-3 + 2i$, and in fact that the additive subgroup $(I, +)$ of the group $(\mathbb{Z}[i], +)$ is generated by $2 + 3i$ and $-3 + 2i$.
- (ii) Show that $i \equiv -5 \pmod{I}$, i.e. that $i + 5 \in I$.
- (iii) Show that the homomorphism $f: \mathbb{Z} \rightarrow \mathbb{Z}[i]/I$ defined by $f(n) = n + I$ is surjective.
- (iv) Show that $13 \in \mathbb{Z} \cap I$.
- (v) Show that $\mathbb{Z} \cap I = 13\mathbb{Z}$. (This may take some calculation.)
- (vi) Conclude by the previous problem that $\mathbb{Z}[i]/I \cong \mathbb{Z}/13\mathbb{Z}$ as rings. Is I a maximal ideal? A prime ideal?

(Note: for those who took Modern Algebra I last semester, $\mathbb{Z}[i] \cong \mathbb{Z} \times \mathbb{Z}$ as (additive) groups and the ideal I , viewed as an additive subgroup of $\mathbb{Z} \times \mathbb{Z}$, is the subgroup generated by $(2, 3)$ and $(-3, 2)$. A homework problem from last semester then shows that the additive group $\mathbb{Z}[i]/I$ is isomorphic to $\mathbb{Z}/(\det A)\mathbb{Z}$, where A is the matrix $\begin{pmatrix} 2 & -3 \\ 3 & 2 \end{pmatrix}$ and hence $\det A = 13$.)