

**MODERN ALGEBRA II SPRING 2013:
ELEVENTH PROBLEM SET**

1. Let F be a field. Show that F is algebraically closed in $F(t)$, the field of rational functions with coefficients in F , where t is an indeterminate (variable). In other words, show that, if $r(t) \in F(t)$ and $r(t)$ is algebraic over F , then $r(t) \in F$. (Hint: suppose that $r = p(t)/q(t)$ is a rational function in lowest terms, i.e. $p(t)$ and $q(t)$ are relatively prime polynomials in t , and that $f(r) = 0$ for some monic $f(x) \in F[x]$. The case $r = 0$ is clear, so we may assume $r \neq 0$ and hence that the constant term of f is nonzero. Viewing $F[x]$ as a subring of $F[t][x]$ and hence of $F(t)[x]$, conclude by the rational roots test that the numerator and denominator of r are constants, hence that $r \in F$.)
2. Let R be a UFD and let $f(x), g(x) \in R[x]$ be two nonzero polynomials. Show that $c(fg) = c(f)c(g)$.
3. Let E be a finite extension field of a field F , and let $\sigma: E \rightarrow E$ be a homomorphism such that $\sigma(a) = a$ for all $a \in F$. Show that the image of σ , $\sigma(E)$, is equal to E , i.e. that σ is surjective (and hence σ is an automorphism of E and thus an element of $\text{Gal}(E/F)$). (Hint: using the fact that σ is F -linear, show that the image $\sigma(E)$ is an F -vector subspace of E . Now use the fact that σ is injective.)
4. Consider the field $\mathbb{Q}(\sqrt{2}, \sqrt{3})$, with \mathbb{Q} -basis $1, \sqrt{2}, \sqrt{3}, \sqrt{6}$. We have seen in class that $\text{Gal}(\mathbb{Q}(\sqrt{2}, \sqrt{3})/\mathbb{Q}) = \{1, \sigma_1, \sigma_2, \sigma_3\}$, where $\sigma_1(\sqrt{2}) = -\sqrt{2}$, $\sigma_1(\sqrt{3}) = \sqrt{3}$, $\sigma_2(\sqrt{2}) = \sqrt{2}$, $\sigma_2(\sqrt{3}) = -\sqrt{3}$, and $\sigma_3 = \sigma_1\sigma_2$. Find the fixed fields $\mathbb{Q}(\sqrt{2}, \sqrt{3})^{\langle \sigma_1 \rangle}$, $\mathbb{Q}(\sqrt{2}, \sqrt{3})^{\langle \sigma_2 \rangle}$, $\mathbb{Q}(\sqrt{2}, \sqrt{3})^{\langle \sigma_3 \rangle}$.
5. Let σ be complex conjugation acting on the field $\mathbb{Q}(\sqrt[3]{2}, \omega)$, where $\omega = e^{2\pi i/3} = \frac{1}{2}(-1 + \sqrt{-3}) = \frac{1}{2}(-1 + i\sqrt{3})$ is a cube root of unity, and hence is a root of $x^2 + x + 1$. Show that $1, \omega$ is a basis for $\mathbb{Q}(\sqrt[3]{2}, \omega)$ over $\mathbb{Q}(\sqrt[3]{2})$, i.e. is a basis for the $\mathbb{Q}(\sqrt[3]{2})$ -vector space $\mathbb{Q}(\sqrt[3]{2}, \omega)$. Find the fixed field $\mathbb{Q}(\sqrt[3]{2}, \omega)^{\langle \sigma \rangle}$, where σ is complex conjugation.
6. We have seen that the polynomial $\Phi_5(x) = \frac{x^5 - 1}{x - 1}$ is an irreducible polynomial in $\mathbb{Q}[x]$ of degree 4, and hence that $[\mathbb{Q}(\zeta) : \mathbb{Q}] = 4$, where $\zeta = e^{2\pi i/5}$ is a root of $\Phi_5(x)$.
 - (a) Show that the four roots of $\Phi_5(x)$ in \mathbb{C} are $\zeta_1 = \zeta$, $\zeta_2 = \zeta^2$, $\zeta_3 = \zeta^3$, and $\zeta_4 = \zeta^4$. Conclude that there is an injective homo-

morphism from $\text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})$ to S_4 , viewed as the set of permutations of the set $\{\zeta_1, \zeta_2, \zeta_3, \zeta_4\}$.

- (b) Given $\sigma \in \text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})$, show that σ is determined by its value on ζ , and that there are at most 4 possibilities for $\sigma(\zeta)$. Hence $\#(\text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})) \leq 4$. (In fact, $\#(\text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})) = 4$ and that $\text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q})$ is cyclic; we shall see this later.)

7. Show that there is no automorphism $\sigma: \mathbb{R} \rightarrow \mathbb{R}$ such that $\sigma(\sqrt{2}) = -\sqrt{2}$. (What is $\sigma(\sqrt[4]{2})$?)

(Note: in fact, one can show that $\text{Aut } \mathbb{R} = \{\text{Id}\}$ as follows: let $\sigma \in \text{Aut } \mathbb{R}$. Then $t \in \mathbb{R}$ is a square $\iff \sigma(t)$ is a square, hence $t > 0 \iff \sigma(t) > 0$. Thus $t_1 < t_2 \implies \sigma(t_1) < \sigma(t_2)$. Now use the fact that, if $r \in \mathbb{Q}$, $\sigma(r) = r$ and that, for every real number t and every $\varepsilon > 0$, we can find rational numbers $r_1 < r_2$ with $r_1 < t < r_2$ and $r_2 - r_1 < \varepsilon$, to conclude that $r_1 < \sigma(t) < r_2$ and that $|t - \sigma(t)| < \varepsilon$. Since this is true for all $\varepsilon > 0$, $\sigma(t) = t$ for every $t \in \mathbb{R}$, i.e. $\sigma = \text{Id}$.)