

Riemann Surfaces: Lecture Notes

Nilay Kumar

Last updated: October 3, 2013

Class 10

Recall that we are studying function theory on the torus \mathbb{C}/Λ , where $\Lambda = \{m\omega_1 + n\omega_2; m, n \in \mathbb{Z}\}$. We had produced a candidate

$$\sigma(z) = z \prod_{\omega \in \Lambda^\times} \left(1 + \frac{z}{\omega}\right) e^{-\frac{z}{\omega} + \frac{1}{2} \frac{z^2}{\omega^2}}.$$

Let us check that this product converges by examining its logarithm:

$$\begin{aligned} \log \{\cdots\} &= \log\left(1 + \frac{z}{\omega}\right) - \frac{z}{\omega} + \frac{z^2}{2\omega^2} \\ &= \left(\frac{z}{\omega} - \frac{1}{2} \frac{z^2}{\omega^2} + \cdots\right) - \frac{z}{\omega} + \frac{1}{2} \frac{z^2}{\omega^2}, \end{aligned}$$

which clearly converges. Hence $\sigma(z)$ is holomorphic for $z \in \mathbb{C}$. Recall that $\sigma'(z)/\sigma(z) = \zeta(z)$ and so $\partial_z \log \sigma(z + \omega_a) = \zeta(z + \omega_a)$. Thus we have (from before) that

$$\eta_a = \zeta(z + \omega_a) - \zeta(z) = \partial_z \log \sigma(z + \omega_a) - \partial_z \log \sigma(z),$$

which gives us periodicity information. Integrating and exponentiating, we see that

$$\sigma(z + \omega_a) = \sigma(z) e^{\eta_a z + c_a}$$

where c_a is the constant of integration, and taking $z = -\omega_a/2$, we find that

$$\sigma(\omega_a/2) = \sigma(-\omega_a/2) e^{-\eta_a \frac{\omega_a}{2} + c_a}.$$

It is easy to check, however, that σ is odd, and hence we find that

$$\sigma(z + \omega_a) = -\sigma(z) e^{\eta_a(z + \frac{\omega_a}{2})}.$$

So we have found that $\sigma(z)$ is holomorphic on \mathbb{C} and that $\sigma(z) = 0$ if and only if $z = 0 \pmod{\Lambda}$. Now that we have constructed such a σ , let us give another proof of Abel's theorem. First recall our previous statement of Abel's theorem.

Theorem 1 (Abel's theorem). *Let $P_1, \dots, P_M, Q_1, \dots, Q_N$ be points in \mathbb{C} . Then there exists a meromorphic f with zeroes at P_i and poles at Q_i if and only if $M = N$ and $\sum_{i=1}^M A(P_i) = \sum_{i=1}^N A(Q_i)$.*

Recall that the Abel map takes $\mathbb{C}/\Lambda \ni p \mapsto A(p) = \int_{p_0}^p \omega$ where the value of the integral is taken modulo the lattice generated by $\oint_A \omega, \oint_B \omega$. Take $p_0 = 0$ and $\omega = dz$, which is a well-defined form, and if we take A to align with ω_2 and B to align with ω_1 , we see that $\oint_A \omega = \oint_A dz = \omega_1$ and similarly $\oint_B \omega = \omega_2$. Hence the map simply takes p to $\int_0^p dz \mod \Lambda = p$ where p is viewed as a complex number.

Let us now restate Abel's theorem.

Theorem 2 (Abel's theorem, v.2). *Let $P_1, \dots, P_M, Q_1, \dots, Q_N$ be points in \mathbb{C} . Then there exists a meromorphic f with zeroes at P_i and poles at Q_i if and only if $M = N$ and $\sum_{i=1}^M P_i = \sum_{i=1}^N Q_i \mod \Lambda$.*

Proof. Consider the function

$$f(z) = \frac{\prod_{i=1}^M \sigma(z - P_i)}{\prod_{i=1}^N \sigma(z - Q_i)}.$$

We should be a little careful to note that σ is a function not on the torus \mathbb{C}/Λ , but a function on \mathbb{C} (it transforms under a lattice translation!). Hence we must be cognizant of the fact that P_i, Q_i here are some chosen representatives in \mathbb{C} of the equivalence classes of the points P_i, Q_i . It should be clear that $f(z)$ is meromorphic with zeroes at every representative of each P_i s and poles at every representative of each Q_i . The natural question, now, is whether this function extends to a function on the torus. To check this, let us see whether it is doubly periodic using what we know about σ :

$$\begin{aligned} f(z + \omega_a) &= f(z) \frac{\prod_{i=1}^M e^{\eta_a(z - P_i)}}{\prod_{i=1}^N e^{\eta_a(z - Q_i)}} \\ &= f(z) e^{-\eta_a(\sum_{i=1}^M P_i - \sum_{i=1}^N Q_i)}. \end{aligned}$$

Hence we wish to choose P_i, Q_i representatives such that the exponential becomes unity. By hypothesis, this can be done (by shifting one, if necessary). \square

Let us now return to Weierstrass theory. Given $\omega = dz$, we defined $\omega_0 = \mathcal{P}(z)dz$ which has a double pole at 0 and $\partial_z \log \sigma(z) = \zeta(z)$ and $\zeta'(z) = -\mathcal{P}(z)$. Now we can construct a form ω_{PQ} with residues 1, -1 at P, Q respectively, by assigning $\omega_{PQ}(z) = (\zeta(z - P) - \zeta(z - Q))\omega = \partial_z \log \frac{\sigma(z - P)}{\sigma(z - Q)} dz$. What Weierstrass theory tells us that we can write everything in terms of σ , our analog of z .

Jacobi theory: θ -functions

Consider again the torus \mathbb{C}/Λ , where we now normalize the lattice as $\Lambda = \{m + n\tau; m, n \in \mathbb{Z}\}$ with $\text{Im } \tau > 0$ (by linear independence, it cannot be real). This simply corresponds to picking $\omega_1 = 1, \omega_2/\omega_1 = \tau$. Next define the **theta-function**

$$\theta(z|\tau) = \sum_{n \in \mathbb{Z}} e^{\pi i n^2 \tau + 2\pi i n z},$$

in which the structure of the lattice is explicitly clear (unlike in the Weierstrass theory). Let us examine its main properties.

First, note that $\theta(z|\tau)$ is holomorphic in $z \in \mathbb{C}$ because the series converges for all z ; this is due to the term

$$|e^{\pi i n^2 (\tau_1 + i\tau_2)}| = |e^{\pi i n^2 \tau_1} e^{-\pi^2 n^2 \tau_2}| = e^{-\pi n^2 \tau_2}$$

for $\tau = \tau_1 + i\tau_2$, whose decay dominates due to the n^2 . Next, notice that

$$\begin{aligned}\theta(z+1|\tau) &= \theta(z|\tau) \\ \theta(z+\tau|\tau) &= e^{-\pi i\tau-2\pi iz}\theta(z|\tau),\end{aligned}$$

where the second is obtained by completing the square. Though θ is not invariant, its zeroes are.

Furthermore, we claim that $\theta(z|\tau)$ vanishes at exactly one point modulo lattice translates. It suffices to compute the integral $\oint_C \frac{\theta'(z|\tau)}{\theta(z|\tau)} dz$, as it yields $2\pi i$ times the difference in the number of zeroes and poles in a given region. We shall integrate over the curve C where C traverses the circumference of one lattice segment (i.e. the whole torus):

$$\oint_C \frac{\theta'(z|\tau)}{\theta(z|\tau)} dz = \oint_B \left(-\frac{\theta'(z|\tau)}{\theta(z|\tau)} + \frac{\theta'(z+1|\tau)}{\theta(z+1|\tau)} \right) + \oint_A \left(\frac{\theta'(z|\tau)}{\theta(z|\tau)} - \frac{\theta'(z+\tau|\tau)}{\theta(z+\tau|\tau)} \right).$$

But these are just the shifts in the logarithmic derivative, and since $\partial_z \log \theta(z+\tau|\tau) = -2\pi i + \partial_z \log \theta(z|\tau)$ using the transformation rules above, we see that our integral simplifies to

$$\oint_C \frac{\theta'(z|\tau)}{\theta(z|\tau)} dz = 2\pi i \oint_A dz = 2\pi i.$$

Of course, since θ is holomorphic, it has no poles, and hence we see that we have one zero. The zero, in fact, occurs in the center: $\theta((1+\tau)/2|\tau) = 0$. To see this, consider the following function:

$$\begin{aligned}\theta\left(z + \frac{1+\tau}{2}|\tau\right) &= \sum_{n \in \mathbb{Z}} \exp\left(\pi i n^2 \tau + 2\pi i n \left(z + \frac{1+\tau}{2}\right)\right) \\ &= i \exp\left(-\pi i \frac{\tau}{4} - \pi i z\right) \sum_{n \in \mathbb{Z}} \exp\left(\pi i \left(n + \frac{1}{2}\right)^2 \tau + 2\pi i \left(n + \frac{1}{2}\right) \left(z + \frac{1}{2}\right)\right) \\ &= i \exp\left(-\pi i \frac{\tau}{4} - \pi i z\right) \theta_1(z|\tau)\end{aligned}$$

where we have completed the square and defined the function θ_1 . We claim that θ_1 is an odd function, which would imply that θ_1 vanishes at zero, which would prove the claim about the location of the zero. Hence let us verify that θ_1 is odd; switching $z \mapsto -z$ yields in the exponent

$$\log \theta_1(z|\tau) = \pi i \left(n + \frac{1}{2}\right)^2 \tau + 2\pi i \left(n + \frac{1}{2}\right) \left(-z + \frac{1}{2}\right).$$

If we switch the indices $n \mapsto m$ such that $n + \frac{1}{2} = -(m + \frac{1}{2})$, we find that the exponent is now

$$\log \theta_1(z|\tau) = \pi i \left(m + \frac{1}{2}\right)^2 \tau + 2\pi i \left(m + \frac{1}{2}\right) \left(\left(z + \frac{1}{2}\right) - 2\pi i \left(m + \frac{1}{2}\right)\right),$$

and hence θ_1 is odd. Now we see that the function we want is in fact $\theta_1(z|\tau)$ as it is odd, holomorphic, and has one zero.

We leave it as an exercise to show that

$$\sigma(z) = \omega_1 \exp\left(\eta_1 \frac{z^2}{\omega_1}\right) \frac{\theta_1\left(\frac{z}{\omega_1}|\tau\right)}{\theta_1'(0|\tau)}$$