Opioid Pain Management Analytics

A Data-Driven Analysis of Postoperative Pain Management
at Riverbend Medical Center

Corey Reynolds
Grand Canyon University
Healthcare Data Management | 2026

Introduction

Purpose

Opioid medications remain essential in postoperative pain management, yet their use carries serious risks — including inadequate pain control, adverse reactions, and dependency. This analysis leverages a purpose-built relational database to examine opioid administration patterns, pain relief effectiveness, and adverse event trends across seven postoperative patients at Riverbend Medical Center.

Track Effectiveness

Monitor opioid impact on pain scores and recovery time

Identify Risk

Flag potential misuse, dependency, or adverse events

Inform Policy

Support providers, policymakers, and researchers

Database Design Overview

Six relational tables were designed to capture the full clinical picture of postoperative opioid management:

Patients

Demographics & medical history

Doctors

Provider & specialization

Surgeries

Procedure type, date, care plan

Opioid Administration

Drug, dosage, route, duration

Pain Records

Pain scale, time to relief

Adverse Events

Event type, severity, date

Key Findings: Opioid Usage Patterns



5

unique opioids administered

7.5

days max avg Oxycodone duration

100%

of high-pain patients received oral opioids

Key Findings: Pain Relief Effectiveness

Derived from SQL Query 2 — pain relief effectiveness by opioid type:

Opioid Type	Avg Pain Score	Avg Time to Relief	Avg Pain-Free Days
Fentanyl	8.0	15 min	2.0
Morphine	8.0	20 min	2.0
Oxycodone	8.0	52.5 min	2.0
Tramadol	6.0	40 min	4.0
Hydrocodone	4.5	27.5 min	5.5

Key Insight: Hydrocodone and Tramadol produced more pain-free days than faster-acting IV opioids, suggesting that milder oral regimens may offer better sustained relief for lower-acuity procedures. Fentanyl showed fastest relief onset but fewest pain-free days, consistent with its short-acting IV profile.

Identified Issues



Opioid Dependency Risk

Patient 4 (Lumbar Fusion, Oxycodone 10mg × 10 days) requested early refill — highest misuse indicator in the dataset.

Respiratory Depression (x2)

Patients 3 and 7 experienced respiratory depression requiring clinical intervention. Both were on IV opioids.

Inadequate Pain Control

3 of 7 patients (43%) maintained pain scores ≥ 8 despite opioid therapy, signaling undertreated postoperative pain.

Demographic Disparity

Hispanic and Black patients accounted for 4 of 7 adverse events — warranting closer monitoring for equitable care.

Christian Worldview Reflection

Ethical Pain Management, Patient Dignity, and Holistic Care

I

Human Dignity & Patient-Centered Care

A Christian worldview holds that every person is created in the image of God (imago Dei). This mandates that pain management decisions honor each patient's inherent worth — ensuring that no patient, regardless of ethnicity or background, experiences undertreated pain or is over-prescribed opioids based on bias.

II

Ethical Stewardship of Medicine

Stewardship principles call healthcare providers to use powerful medications wisely. The data showing two cases of respiratory depression and one dependency-risk patient underscore the moral obligation to prescribe opioids at the lowest effective dose, with close monitoring and clear care plans.

III

Holistic & Restorative Care

Scripture's vision of shalom — wholeness and flourishing — aligns with multimodal pain management. Integrating physical therapy, non-opioid analgesics, and psychological support alongside opioids reflects a whole-person approach that promotes genuine healing rather than dependence.

Recommendations

01 Strengthen Monitoring Protocols

Implement mandatory pain reassessment at 24h and 48h post-administration. Flag patients on high-dose opioids (≥ 10mg) or extended duration (≥ 7 days) for pharmacist review.

02 Adopt Multimodal Pain Strategies

Reduce reliance on opioids by pairing them with NSAIDs, nerve blocks, and non-pharmacological interventions like PT. Results show Tramadol + PT produced 4 pain-free days vs. 2 for IV opioids.

03 Address Demographic Disparities

Standardize prescribing guidelines across all patient demographics. Adverse event data showed disproportionate impact on minority patients — provider implicit bias training is warranted.

04 Expand the Database for Research

Scale this schema to include larger patient cohorts, longitudinal follow-up, and EHR integration (Epic/Cerner) to enable population-level opioid outcome analysis.

Conclusion

What We Found

- Opioid duration and route significantly affect pain relief outcomes.
- IV opioids provide faster onset but fewer sustained pain-free days.
- Critical and severe adverse events were exclusively tied to IV administration.
- Potential opioid misuse was identified in 1 of 7 patients (14%).
- Demographic disparities in adverse events require targeted intervention.

Significance

This database provides a scalable, evidence-based framework for healthcare providers to optimize pain management protocols, reduce opioid misuse, and ensure equitable care. When guided by ethical principles — including respect for human dignity and holistic healing — data-driven approaches can meaningfully improve postoperative patient outcomes.

References

1. Centers for Disease Control and Prevention. (2022). CDC clinical practice guideline for prescribing opioids for pain — United States, 2022. MMWR Recommendations and Reports, 71(3), 1–95. https://doi.org/10.15585/mmwr.rr7103a1

2. Institute of Medicine. (2011). Relieving pain in America: A blueprint for transforming prevention, care, education, and research. The National Academies Press.

3. Jungquist, C. R., Pasero, C., Tripoli, N. M., Gorodetsky, R., Metersky, M., & Polomano, R. C. (2014). Instituting best practice for monitoring outcomes in hospitalized patients receiving opioid analgesia. Worldviews on Evidence-Based Nursing, 11(2), 82–91.

4. Matthias, M. S., Parpart, A. L., Nyland, K. A., Huffman, M. A., Stubbs, D. L., Sargent, C., & Bair, M. J. (2010). The patient–provider relationship in chronic pain care: Providers' perspectives. Pain Medicine, 11(11), 1688–1697.

5. National Institute on Drug Abuse. (2023). Opioid overdose crisis. U.S. Department of Health and Human Services. https://nida.nih.gov/research-topics/opioids/opioid-overdose-crisis

6. Peacock, P. J. & Peacock, J. L. (2018). Evidence-based healthcare: A practical guide for therapists. Elsevier.

7. World Health Organization. (2019). WHO guidelines for the pharmacological and radiotherapeutic management of cancer pain in adults and adolescents. WHO Press.