

COVID-19 Baseline Risk Score Analysis Report

MockENSEMBLE Study

USG COVID-19 Response Biostatistics Team

August 04, 2021

Contents

1	Baseline Risk Score (Proxy for SARS-CoV-2 Exposure)	9
2	Appendix	21

List of Tables

1.1	Variables considered for risk score analysis.	9
1.2	All learner-screen combinations (14 in total) used as input to the Superlearner.	11
1.3	Weights assigned by Superlearner.	15
1.4	Predictors in learners assigned weight > 0.0 by Superlearner. . .	16
1.5	Cases per treatment arm prior to risk score analysis.	20
1.6	Cases per treatment arm post risk score analysis.	20

List of Figures

1.1	Cross-validated AUC (95% CI) of algorithms for predicting COVID-19 disease status starting 7 days after Day 29.	12
1.2	CV-estimated predicted probabilities of COVID-19 disease 7 days after Day 29 by case/control status for top 2 learners, Super-Learner and Discrete SL.	13
1.3	ROC curves based off CV-estimated predicted probabilities for the top 2 learners, Superlearner and Discrete SL.	14
1.4	Superlearner predicted probabilities of COVID-19 disease in vaccinees 7 days after Day 29 by case/control status.	18
1.5	ROC curve based off Superlearner predicted probabilities in vaccinees.	19

MOCK

Chapter 1

Baseline Risk Score (Proxy for SARS-CoV-2 Exposure)

Table 1.1: Variables considered for risk score analysis.

Variable.Name	Definition	Total.missing.values	Comments
EthnicityHispanic	Indicator ethnicity = Hispanic (1 = Hispanic, 0 = complement)	0/19501 (0.0%)	NA
EthnicityNotreported	Indicator ethnicity = Not reported (1 = Not reported, 0 = complement)	0/19501 (0.0%)	NA
EthnicityUnknown	Indicator ethnicity = Unknown (1 = Unknown, 0 = complement)	0/19501 (0.0%)	NA
Black	Indicator race = Black (1=Black, 0=complement)	0/19501 (0.0%)	NA
Asian	Indicator race = Asian (1=Asian, 0=complement)	0/19501 (0.0%)	NA
NatAmer	Indicator race = American Indian or Alaska Native (1=NatAmer, 0=complement)	0/19501 (0.0%)	NA
Multiracial	Indicator race = Multiracial (1=Multiracial, 0=complement)	0/19501 (0.0%)	NA
Notreported	Indicator race = Not reported (1=Notreported, 0=complement)	0/19501 (0.0%)	NA
Unknown	Indicator race = unknown (1=Unknown, 0=complement)	0/19501 (0.0%)	NA
URMforsubcohortsampling	Indicator of under-represented minority (1=Yes, 0=No)	0/19501 (0.0%)	NA
HighRiskInd	Baseline covariate indicating ≥ 1 Co-existing conditions (1=yes, 0=no, NA=missing)	0/19501 (0.0%)	NA
Sex	Sex assigned at birth (1=female, 0=male/undifferentiated/unknown)	0/19501 (0.0%)	NA
Age	Age at enrollment in years (integer ≥ 18 , NA=missing). Note that the randomization strata included Age 18-59 vs. Age ≥ 60 .	0/19501 (0.0%)	NA

Table 1.1: Variables considered for risk score analysis. (*continued*)

Variable.Name	Definition	Total.missing.values
BMI	BMI at enrollment (Ordered categorical 1,2, 3, 4, NA=missing); 1 = Underweight BMI < 18.5; 2 = Normal BMI 18.5 to < 25; 3 = Overweight BMI 25 to < 30; 4 = Obese BMI >= 30	0/19501 (0.0%)
Country	Country of the study site of enrollment (0=United States, 1=Argentina,2=Brazil, 3=Chile,4=Columbia, 5=Mexico, 6=Peru, 7=South Africa)	0/19501 (0.0%)
HIVinfection	Indicator HIV infected at enrollment (1=infected, 0=not infected)	0/19501 (0.0%)
CalendarDateEnrollment	Date variable (used to control for calendar time trends in COVID incidence). Coded as number of days since first person enrolled until the ppt is enrolled.	0/19501 (0.0%)

Note:

Variables with more than 5% missing values were dropped from analysis; missing values for other variables were dropped. Indicator variables not meeting the threshold, such that under the null of not a risk factor there were less than 5% in the subgroup with value 1 (or 0), were dropped from analysis.

Table 1.2: All learner-screen combinations (14 in total) used as input to the Superlearner.

Learner	Screen*
SL.mean	all
SL.glm	all glmnet univar_logistic_pval highcor_random

Note:

*Screen details:

all: includes all variables

glmnet: includes variables with non-zero coefficients in the standard implementation of SL.glmnet that optimizes the lasso tuning parameter via cross-validation

univar_logistic_pval: Wald test 2-sided p-value in a logistic regression model < 0.10

highcor_random: if pairs of quantitative variables with Spearman rank correlation > 0.90 , select one of the variables at random



Figure 1.1: Cross-validated AUC (95% CI) of algorithms for predicting COVID-19 disease status starting 7 days after Day 29.



Figure 1.2: CV-estimated predicted probabilities of COVID-19 disease 7 days after Day 29 by case/control status for top 2 learners, SuperLearner and Discrete SL.



Figure 1.3: ROC curves based off CV-estimated predicted probabilities for the top 2 learners, Superlearner and Discrete SL.

Table 1.3: Weights assigned by Superlearner.

Learner	Screen	Weight
SL.glm	screen_all	0.833
SL.glm	screen_glmnet	0.167
SL.mean	screen_all	0.000
SL.glm	screen_univariate_logistic_pval	0.000
SL.glm	screen_highcor_random	0.000

Table 1.4: Predictors in learners assigned weight > 0.0 by Superlearner.

Learner	Screen	Weight	Predictors	Coefficient	Odds.Ratio
SL.glm	screen_all	0.833	(Intercept)	-19.397	0
SL.glm	screen_all	0.833	EthnicityHispanic	-0.037	0.964
SL.glm	screen_all	0.833	EthnicityNotreported	0.013	1.013
SL.glm	screen_all	0.833	EthnicityUnknown	0.05	1.051
SL.glm	screen_all	0.833	Black	-0.16	0.852
SL.glm	screen_all	0.833	Asian	-0.066	0.936
SL.glm	screen_all	0.833	NatAmer	-0.056	0.945
SL.glm	screen_all	0.833	Multiracial	-0.039	0.961
SL.glm	screen_all	0.833	Notreported	-1.403	0.246
SL.glm	screen_all	0.833	Unknown	-0.056	0.946
SL.glm	screen_all	0.833	URMforsubcohortsampling	0.063	1.065
SL.glm	screen_all	0.833	HighRiskInd	0.379	1.461
SL.glm	screen_all	0.833	Sex	12.071	174754.678
SL.glm	screen_all	0.833	Age	0.354	1.425
SL.glm	screen_all	0.833	BMI	-0.046	0.955
SL.glm	screen_all	0.833	Country	-0.017	0.983
SL.glm	screen_all	0.833	HIVinfection	0.011	1.011
SL.glm	screen_all	0.833	CalendarDateEnrollment	0.027	1.027
SL.glm	screen_glmnet	0.167	(Intercept)	-19.395	0
SL.glm	screen_glmnet	0.167	EthnicityNotreported	0.015	1.015
SL.glm	screen_glmnet	0.167	EthnicityUnknown	0.052	1.053
SL.glm	screen_glmnet	0.167	Black	-0.136	0.873
SL.glm	screen_glmnet	0.167	Asian	-0.067	0.936
SL.glm	screen_glmnet	0.167	NatAmer	-0.055	0.947
SL.glm	screen_glmnet	0.167	Multiracial	-0.04	0.961
SL.glm	screen_glmnet	0.167	Notreported	-1.403	0.246
SL.glm	screen_glmnet	0.167	Unknown	-0.056	0.946
SL.glm	screen_glmnet	0.167	URMforsubcohortsampling	0.03	1.031
SL.glm	screen_glmnet	0.167	HighRiskInd	0.379	1.461
SL.glm	screen_glmnet	0.167	Sex	12.07	174555.444
SL.glm	screen_glmnet	0.167	Age	0.354	1.425
SL.glm	screen_glmnet	0.167	BMI	-0.046	0.955
SL.glm	screen_glmnet	0.167	Country	-0.013	0.987
SL.glm	screen_glmnet	0.167	HIVinfection	0.011	1.011
SL.glm	screen_glmnet	0.167	CalendarDateEnrollment	0.026	1.027
SL.glm	screen_glmnet	0.167	(Intercept)	-19.395	0
SL.glm	screen_glmnet	0.167	EthnicityNotreported	0.015	1.015
SL.glm	screen_glmnet	0.167	EthnicityUnknown	0.052	1.053
SL.glm	screen_glmnet	0.167	Black	-0.136	0.873
SL.glm	screen_glmnet	0.167	Asian	-0.067	0.936
SL.glm	screen_glmnet	0.167	NatAmer	-0.055	0.947
SL.glm	screen_glmnet	0.167	Multiracial	-0.04	0.961
SL.glm	screen_glmnet	0.167	Notreported	-1.403	0.246
SL.glm	screen_glmnet	0.167	Unknown	-0.056	0.946
SL.glm	screen_glmnet	0.167	URMforsubcohortsampling	0.03	1.031

Table 1.4: Predictors in learners assigned weight > 0.0 by Superlearner. *(continued)*

Learner	Screen	Weight	Predictors	Coefficient	Odds.Ratio
SL.glm	screen_glmnet	0.167	HighRiskInd	0.379	1.461
SL.glm	screen_glmnet	0.167	Sex	12.07	174555.444
SL.glm	screen_glmnet	0.167	Age	0.354	1.425
SL.glm	screen_glmnet	0.167	BMI	-0.046	0.955
SL.glm	screen_glmnet	0.167	Country	-0.013	0.987
SL.glm	screen_glmnet	0.167	HIVinfection	0.011	1.011
SL.glm	screen_glmnet	0.167	CalendarDateEnrollment	0.026	1.027



Figure 1.4: Superlearner predicted probabilities of COVID-19 disease in vaccinees 7 days after Day 29 by case/control status.



Figure 1.5: ROC curve based off Superlearner predicted probabilities in vaccinees.

Table 1.5: Cases per treatment arm prior to risk score analysis.

Study-Arm	Non-Cases	Post-Day 29-Cases
Placebo	18756	521
Vaccine	19198	170

Table 1.6: Cases per treatment arm post risk score analysis.

Study-Arm	Non-Cases	Post-Day 29-Cases
Placebo	18756	521
Vaccine	19198	170

Chapter 2

Appendix

- This report was built from the [CoVPN/correlates_reporting](https://github.com/CoVPN/correlates_reporting) repository with commit hash 807f1671fac49f7ab0cc0cd9212995b7c06fbd97. A diff of the changes introduced by that commit may be viewed at https://github.com/CoVPN/correlates_reporting/commit/807f1671fac49f7ab0cc0cd9212995b7c06fbd97
- The sha256 hash sum of the raw input file, “COVID_ENSEMBLE_practicedata.csv”:
186c6fefe0d7ee781c3b0bf5cedee3686398eb67ddc3da19dfa0c9bf3ad88fcc
- The sha256 hash sum of the processed file, “janssen_pooled_mock_data_processed.csv”:
1f8c3beacf95500be32a6d6866656b93759de1680e60d61f8a34341b2a4458a6