


North Coast (Areas 3 & 4) Recreational Angling Creel Survey 2025

FINAL REPORT



Prepared for:

North Coast - Skeena First Nations Stewardship Society
Prince Rupert, BC

Prepared by:

David Robichaud¹ and Angela Addison²

¹ LGL Limited environmental research associates
9768 Second Street, Sidney, BC, V8L 3Y8

² North Coast - Skeena First Nations Stewardship Society
309 2 Ave W #363, Prince Rupert, BC, V8J 3T1



EXECUTIVE SUMMARY

Because the North Coast (Areas 3 and 4) recreational fishery can substantially affect multiple transboundary salmon stocks, it is considered a high priority for monitoring. Collecting effort and harvest data, along with biological samples and genetic information, is essential for generating fishery impact estimates, supporting effective fisheries management, and tracking indicator stocks. The North Coast Skeena First Nations Stewardship Society (NCSFNSS) has operated this program since 2015, with support from the Pacific Salmon Commission Northern Fund. The survey consists of access point angler interviews and aerial overflights, previously for the months of June through August but also including May since 2019. Lodge-based recreational harvest and effort are monitored by DFO via a voluntary logbook program. Survey design and implementation, and analytical methods are similar to and consistent with those used in Area 3 and 4 creel survey monitoring since 2008.

From 1 May to 31 August 2025, a total of 2,961 vessel interviews (involving 9,836 anglers) were collected by staff stationed at one of three access points (located in Prince Rupert or Port Edward) during 261 survey shifts. 31 aerial effort surveys were conducted. High priority was placed on gathering biosamples and fin-clip incidence rates from Chinook (*Oncorhynchus tshawytscha*) and Coho (*O. kisutch*) salmon. When possible, length measurements were collected from Pacific Halibut (*Hippoglossus stenolepis*). Survey effort was adequate, as relatively good precision was achieved for the harvest estimates for most of the main taxa (all precision goals were met).

Total fishing effort was estimated to be 14,187 vessel trips. In total, an estimated 18,745 Chinook Salmon were caught, with 7,279 fish kept and 11,466 released. It was estimated that 41,046 Coho Salmon were caught, with 38,082 fish kept and 2,964 released. Total Pacific Halibut catch was estimated to be 21,236 individuals, with 11,923 fish kept (biomass estimated at 53.0 net tonnes) and 9,313 released. Total harvest estimates included 10,094 Lingcod (*Ophiodon elongates*; the highest since the survey began), three Yelloweye Rockfish (*Sebastes ruberrimus*), 9,104 other rockfishes (*Sebastes* spp.), and 4,303 groundfish other than Pacific Halibut and Lingcod.

Creel surveyors biosampled 448 Chinook Salmon, collecting length measurements (median 70 cm), adipose fin-clip incidence rates (17.2% were clipped), and scale samples. For Coho Salmon, 530 fish were biosampled (median fork length 64 cm; 0% adipose clipped). Heads were collected from 85 Chinook Salmon and one Coho Salmon for submission to the Salmon Head Recovery Program. Biosampled Pacific Halibut (n=524) had a median length of 78 cm, and average net weights of 4.6 kg and 4.4 kg in Areas 3 and 4, respectively.

During the month of May, fishing effort and Chinook Salmon harvest were estimated at 1,163 boat trips (8% of the May-August effort), and 1,231 fish (17% of the May-August harvest). The Department of Fisheries and Oceans Canada (DFO) is supportive of a May creel survey because genetic samples are needed to resolve stock composition and determine exploitation rates on Nass and Skeena early-run-timing stocks. During the month of May 2025, scale samples were taken from 129 Chinook Salmon, and surveyors processed 23 tagged heads.

Recreational fishing effort in Areas 3 & 4 have been increasing over the last three decades, as has harvest of Coho Salmon and Lingcod. Lingcod harvest in the last five years are the five highest observed for the available time series, despite a coast-wide reduction in the Lingcod bag limit that started in 2018. Chinook Salmon harvests have not increased in recent years and may be declining, however, Chinook Salmon releases are showing an increasing trend. Pacific Halibut harvest did not show a strong among-year trend, likely the result of effective management measures that reduced daily bag limits and imposed maximum size limits. While the increasing catches of some species highlight the need for continued monitoring of the recreational fishery into the future, there is some evidence that management measures can regulate harvest if effort does not increase dramatically.

TABLE OF CONTENTS

LIST OF TABLES	IV
LIST OF FIGURES	V
INTRODUCTION	1
SCOPE.....	1
GOAL AND OBJECTIVES.....	3
METHODS	4
Angler Activity Patterns.....	6
Catch Per Unit Effort Estimation.....	7
Angler Effort Estimation	8
Lodge Effort Census.....	8
Non-lodge Effort Surveys.....	9
Catch and Harvest Estimation.....	11
Biosampling.....	12
Biomass of Harvested Pacific Halibut	12
Statistical Analyses	12
Trailer Census	12
RESULTS.....	13
Angler Interviews.....	13
Angler Activity Patterns.....	14
Catch Per Unit Effort Estimates.....	18
Angler Effort Estimates.....	23
April Effort.....	27
Catch Estimates.....	29
Biosampling.....	39
Harvested Halibut Biomass	41
DISCUSSION.....	41
Creel Estimates	41
May Effort	42
April Effort	46
Precision of the Results	46
Accuracy of the Results.....	46
RECOMMENDATIONS.....	47
ACKNOWLEDGEMENTS.....	48
LITERATURE CITED.....	48
APPENDICES	50
Appendix 1: Data forms	50
Appendix 2: HPUE estimates by period, trip type, subarea and species.....	53
Appendix 3: Counts of actively fishing sport vessels in each subarea, by date	66
Appendix 4: Retained and released fish by month, subarea, and species.	67
Appendix 5: Statistical effects of period on harvest, by species.....	79
Appendix 6: Adipose fin-clip incidence.....	80
Appendix 7: Biosampling results, May 2025	81

LIST OF TABLES

Table 1.	Delineations of temporal strata.....	3
Table 2.	Sample sizes to estimate non-lodge angler activity patterns.....	7
Table 3.	Overflight sampling effort.....	10
Table 4.	Monthly number of shifts, and numbers of boats interviewed per landing site	15
Table 5.	Fishing locations for non-lodge parties, by month and subarea.....	16
Table 6.	Numbers of non-lodge ‘shellfish-only’ boat trips, by month and subarea.....	16
Table 7.	Numbers of guided and unguided boat trips, by landing site.....	16
Table 8.	Sample sizes for CPUE estimates in each by period and subarea.....	19
Table 9.	Subarea data used to calculate CPUE in each period and subarea,	19
Table 10.	Kruskal-Wallis tests for differences in HPUE among periods, by species	21
Table 11.	Paired t-tests for differences in HPUE between guided and non-guided boats, by species	22
Table 12.	Average number of harvested fish per boat trip, by month, from the census of Area 3 lodges.....	22
Table 13.	Lodge, non-lodge, and total effort estimates, by month and subarea	25
Table 14.	Estimated April non-lodge fishing effort in 2019, 2020, 2023, 2024, and 2025.....	28
Table 15.	Total estimated harvest, by subarea, for popular sportfish taxa	30
Table 16.	Total estimated Chinook Salmon catch, by subarea	30
Table 17.	Total estimated Coho Salmon catch, by subarea.....	31
Table 18.	Total estimated Pink Salmon catch, by subarea	31
Table 19.	Total estimated Chum Salmon catch, by subarea.....	32
Table 20.	Total estimated Sockeye Salmon catch, by subarea	32
Table 21.	Total estimated Pacific Halibut catch, by subarea.....	33
Table 22.	Total estimated Lingcod catch, by subarea	33
Table 23.	Total estimated catch of other groundfish by subarea	34
Table 24.	Total estimated rockfish catch, by subarea	34
Table 25.	Kruskal Wallis tests for differences in harvest among months, by species	36
Table 26.	Biosampling results.....	38
Table 27.	Average weight, and harvest of Pacific Halibut	39
Table 28.	Monthly recreational fishing effort from 1995 to 2025	44
Table 29.	Monthly Chinook Salmon, Coho Salmon, Pacific Halibut, and Lingcod harvest, and Chinook Salmon releases, from 1995 to 2025	45

LIST OF FIGURES

Figure 1.	Map of the North Coast study area	2
Figure 2.	Map showing planned flight path of effort survey overflights.....	9
Figure 3.	Number of interviews conducted at each of three landing sites, by month	14
Figure 4.	Angler activity pattern by day type and period	17
Figure 5.	Average number of fish harvested per non-lodge boat trip, by period and subarea.....	20
Figure 6.	Linear regressions showing the relationships between small non-lodge boats counted during overflights and trailers concurrently counted at landing sites	23
Figure 7.	Total estimated numbers of boat trips, by month and subarea	24
Figure 8.	Heat map showing the estimated total effort in each subarea	26
Figure 9.	Total monthly non-lodge fishing effort as estimated using the usual creel methods or from trailer counts.....	27
Figure 10.	Relationship between total monthly non-lodge fishing effort values as estimated using the usual creel methods or from trailer counts	28
Figure 11.	Total estimated Chinook Salmon harvest, by month and subarea.....	35
Figure 12.	Total estimated Coho Salmon harvest, by month and subarea.....	35
Figure 13.	Total estimated Pacific Halibut harvest, by month and subarea.....	37
Figure 14.	Total estimated Lingcod harvest, by month and subarea	37
Figure 15.	Length-frequency distributions of biosampled fish	40
Figure 16.	Fishing effort from 1995 to 2025.....	41
Figure 17.	June to August harvest of Chinook and Coho salmon, and Pacific Halibut from 1995 to 2025, and of Lingcod from 2008 to 2025.....	43
Figure 18.	Number of Chinook Salmon released from June to August, 2009 to 2025	44

INTRODUCTION

The North Coast recreational fishery in DFO's Pacific Region Management Areas 3 and 4, continues to attract both resident and visiting anglers and harvesters who target a diverse assemblage of species. While effort remains largely directed toward Chinook (*Oncorhynchus tshawytscha*) and Coho (*O. kisutch*) salmon, recreational users also pursue Pacific Halibut (*Hippoglossus stenolepis*), Lingcod (*Ophiodon elongates*), and a variety of other finfish or invertebrate species. Fishing activity has increased markedly over the past three decades (e.g., Robichaud and Addison 2025), raising conservation concerns, and highlighting the need to quantify harvest dynamics and associated local fishery impacts, particularly for stocks of concern. Because harvested salmon can originate from Canadian or US (enhanced and wild) stocks, there is a clear need for the creel survey program to quantify local fishery impacts.

This creel survey program helps the Pacific Salmon Commission develop improved information for resource management, including better stock assessment, data acquisition, and improved scientific understanding of factors that might limit salmon production. The creel program generates statistically robust estimates of harvest and effort, and facilitates direct field observations of catches, enabling verification of species identification, quantification of harvests, and collection of biological samples from Chinook and Coho salmon. Notably, crucial data such as CWT (Coded Wire Tag) information, integral to annual stock assessments and the management of Nass and Skeena river Chinook and Coho salmon stocks, are derived from the heads of randomly sampled adipose-clipped fish collected by program observers.

Since its establishment in 2008, the creel survey program in Areas 3 and 4 has consistently generated reliable estimates of fishing effort and species-specific catch rates (Van Tongeren and Winther 2010, Robichaud and Addison 2025). In 2015, the responsibility for implementing the survey transitioned from Fisheries and Oceans Canada (Prince Rupert) to the North Coast Skeena First Nations Stewardship Society (NCSFNSS), with support from the Pacific Salmon Commission Northern Fund. The current survey design (Robichaud et al. 2016) remains consistent with earlier efforts conducted by Fisheries and Oceans Canada, which, in turn, which were themselves adapted from the Strait of Georgia Creel Survey design, developed in 1982 by LGL Limited (English et al. 2002).

This report documents the 2025 recreational catch and effort statistics for fishing activities in Pacific Fishery Management Areas 3 and 4. It also summarizes the program activities, outlines data collection procedures, and describes the analytical approaches that were used to produce the estimates.

SCOPE

The geographic scope of the creel survey was the same as in the previous 15 years (Van Tongeren and Winther 2010, Robichaud et al. 2016, 2017, Robichaud and Addison 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025), i.e., Management Areas 3 and 4 (from Portland Inlet, through Chatham Sound to Porcher Island (Figure 1). For the purposes of data collection and analysis, the study area was divided into 12 geographic strata ('subareas'), including three subareas in Area 3 (3I-3K) and nine subareas in Area 4 (4A-4H, and 4L).




Figure 1. Map of the North Coast study area, showing the major landing sites, fishing lodges, and the boundaries of the subareas that were used as spatial strata in the creel survey design.

The temporal scope of the full study was from 1 May through 31 August 2025. Survey schedules were designed to provide sufficient data to derive statistically robust catch and efforts estimates for each month and subarea. However, mid-season it was decided to use five temporal strata, termed “periods”, which better coincided with changes in the management measures regulating the fishery than did the biologically ‘arbitrary’ calendar months. For ease of among-year comparison, the period-specific data were reconstituted into monthly summaries (Table 1) wherever possible. For this reason, the second through fourth periods were divided each into two sub-periods (Table 1): one sub-period for the days in one month (e.g., 29 -31 May), and the other sub-period for the days in the next month (e.g., 1-22 June).

Table 1. Delineations of temporal strata. The five Periods coincided with changes in the fishery regulations. Periods were further divided into eight Sub-periods to allow data to be easily grouped into four Months.

<i>Chinook Limits</i>	<i>Period</i>	<i>Sub-period</i>	<i>Month</i>
2 per day	1-28 May	1-28 May	May
1 per day	29 May - 22 June	29-31 May	
		1-22 June	June
non retention	23 June - 23 July	23-30 June	
		1-23 July	July
1 per day	24 July - 17 August	24-31 July	
		1-17 August	August
2 per day	18-31 August	18-31 August	

The study included creel analysis of all major local sportfish species, with greater emphasis placed on Chinook and Coho salmon, Pacific Halibut, and Lingcod.

GOAL AND OBJECTIVES

The goal of this study was to conduct a creel survey of the North Coast (Areas 3 and 4) recreational fishery following the design and methodology used by DFO (Van Tongeren and Winther 2010), for the purpose of producing estimates of effort and catch of all species targeted by the recreational fishery. The creel survey included aerial effort counts and access point angler interviews, conducted throughout Areas 3 and 4 from May through August 2025. The precision goal for the study was to estimate total harvests of large components (such as Chinook Salmon) within 15% of the true value 19 times out of 20. Creel survey strata included temporal separation by month (or period) and day type (weekdays vs. weekend/holiday days). Spatial stratifications were based on creel survey subareas. All local fishing areas (i.e., those accessed from major boat launches in the area) were included in our survey design.

The seven objectives for this study were:

1. Maintain software for data management and analysis comparable to DFO's CREST program;
2. Provide monthly (May-August) estimates of the catch and effort of all salmon species targeted by the recreational fishery and associated estimates of precision;
3. Provide DFO and PSC with monthly in-season catch estimates for all salmon species harvested in the North Coast recreational fishery;
4. Provide communication materials to recreational harvesters to convey the usefulness and importance of their catch data in fisheries management;
5. During angler interviews, collect biological data from a representative subsample of the landed harvest, including fin clip incidence, lengths, and scale or tissue samples for Coho and Chinook salmon;
6. Produce a final written report to be available to a technical audience; and
7. Present creel survey findings at the North Coast salmon post-season review meeting to a broad audience of commercial, recreational interests, and First Nations.

Objective 1 was addressed over the course of the study period. Objectives 2 and 6 were met by completing this report. Objective 3 was met throughout the study period by providing timely and

complete summaries of the monthly survey activities (with complete creel analyses done for each update report). Objective 4 will be met subsequent to the finalization of this report, as its contents will be based on the results presented herein. Objective 5 was met by biosampling fish at access points throughout the study period: fish were measured for length and sexed, scale or tissue samples and adipose fin clip incidence data were collected for Coho and Chinook salmon, heads were collected from adipose-clipped fish, and all biological data were submitted to DFO at the end of study period. Objective 7 was met on 4 December 2025 at the DFO IHPC Post-Season Review meeting.

METHODS

A complete census of catch and effort was available for lodge-based anglers. The census data were collected by lodge staff and shared voluntarily with Fisheries and Oceans Canada stock assessment personnel. The remainder of the recreational fishery occurred over too large an area to completely census catch and effort. Instead, statistically robust methods were used to estimate catch by multiplying estimates of *angler effort* by estimates of *catch per unit of effort*. Separate estimates were generated for each creel monitoring subarea, month, day type (weekday vs. weekend), and species.

For each creel subarea during each period and day type, fishing effort was estimated by counting boats that were actively fishing during overflight surveys (sub-area counts were reduced by the number of lodge-based boats thought to have been visible to the aircraft). Catch per unit of effort for non-lodge boats was estimated from interviews (see data forms in Appendix 1) conducted at known access points. During interviews, anglers were asked about their catch, effort, and fishing locations. They were also asked about their hourly fishing activity patterns.

Data collected during interviews included:

- Date;
- Number of anglers in party;
- Whether or not the trip was guided by a professional;
- Subarea angler activity – for each subarea, the hours during which angling activity was conducted;
- Subarea catch – the number of fish kept and the number released by species and subarea;
- Subarea gear – the amount and type of fishing gears used to catch each species in each subarea;
- Subarea targets – the number of hours spent targeting each species in each subarea (anglers that targeted only shellfish were noted);
- Biosampling – if allowed by the anglers, harvested Chinook and Coho salmon were measured (nose to fork length), sexed, checked for adipose clips and, if clipped and permission given, heads were collected for the Salmon Head Recovery Program. Scale samples were taken from Chinook Salmon, and tissue samples (fin clips) from Coho Salmon. Chinook Salmon were also examined for flesh colour. Pacific Halibut were measured (total length);
- Refusals – whether or not the anglers refused an interview, allowed catch to be observed, or allowed catch to be biosampled; and
- Weather conditions.

The analytical methods used, similar to those used in previous years (e.g., Robichaud and Addison 2025), provided a statistically unbiased estimate of catch per unit effort, assuming the anglers interviewed were representative of the entire non-lodge recreational fishery. Interviews were conducted at the three main access locations: Port Edward and Rushbrooke boat launches, and Prince Rupert Yacht Club. The locations surveyed were selected from all available access points, based on their geographical distribution and the amount of fishing activity that is typically conducted from that site, in order to maximize the number of interviews obtained. This approach was based on two important observations: 1) the variability in CPUE (catch per unit effort) among fishing parties landing at a single access point tends to be as great as the variability in CPUE among different access points within a geographic area; and 2) CPUE and effort can vary substantially both within and between days at a single site (English et al. 2002). Under these conditions it is better to obtain a large number of interviews covering all temporal strata for a small number of sites than to sample a larger number of sites and obtain fewer interviews and less complete temporal coverage for any specific site.

During an observer shift, attempts were made to interview all anglers returning to the access location on that day, but some were missed (it is assumed that a random process dictated whether a boat was missed or interviewed), and some anglers refused to participate (it is not clear how the angling behaviour of the refusals differed from that of the participants). The geographic distribution of our samples was dictated by where the anglers fished, and it was assumed that the distribution of samples taken at the access locations was random relative to the fish harvested by the fishery.

The frequency of interview shifts was planned with the target of sampling 20% of estimated effort, while attempting to have “hands on” and direct observations of (and biological samples from) as many fish as possible meant allocating adequate time to each interview. The interview schedule was designed to capture data from representative angler parties in both day types, and over all time periods of the day. Interviewing sessions were separated into AM and PM shifts, with AM shifts occurring between 7:00 and 15:00, and PM shifts from 13:00 to 21:00. In total, 261 shifts were worked.

Fishing effort was estimated from boat-counts made during overflights. Overflights were scheduled on 3 weekend days and 5 weekdays each month. Mid-season, in response to the announcement of Chinook Salmon management measures, we partitioned our temporal strata into five ‘periods’ (Table 1), rather than the typical four months, which meant that period-specific sample sizes were lower than what they would have been had we generated monthly effort estimates. Timing of the overflights was targeted for peak fishing periods (11 AM to 1 PM), and in 2025 most flights occurred at target times. Fog delayed the start of two flights (flight on 28 August ran from 12:13 PM to 1:56 PM; flight on 30 August ran from 1:10 PM to 2:50 PM); and the airline’s schedule resulted in an early start for one flight (flight on 29 August ran from 10:08 AM to 11:57 AM). Storms (on 28 June and 3 August) resulted in the flights being shortened. Analytical methods accommodate the occasional off-peak flight but can result in increased uncertainty in the effort estimate. In May only seven flights were flown because of weather cancellations (Appendix 3), coupled with the charter being unavailable to reschedule.

Angler Activity Patterns

Two weighting factors were used together with the interview-derived angling activity data to estimate the daily non-lodge fishing activity pattern (English et al. 2002). The first weighting factor, $W1$, expanded the numbers of days spent interviewing at each landing site to account for the total number of days available for sampling. That is, it was assumed that the daily activity pattern recorded during the interview shifts at landing site a , were consistent for landing site a , even during the days when no interviews occurred. A specific $W1$ was calculated for each landing site during each period and day type:

$$W1_{pda} = \frac{N_{pd}}{K_{pda}}, \quad (\text{Equation 1})$$

where N_{pd} was the total number of type d days in period p ; and K_{pda} was the number of days during which interviews occurred at landing site a , on type d days during period p .

The second weighting factor, $W2$, expanded the numbers of interviews conducted, to account for the recreational fishing boats that were *not* interviewed. Some recreational fishing boats were not interviewed because of refusals, and some boats returned during periods of high traffic volume when it was not possible to interview all returning vessels. Some returning vessels were not doing recreational fishing at all (e.g., pleasure cruisers, Food-Social-Ceremonial harvesters). For the recreational fishing vessels, it was assumed that the activity pattern recorded during the interview shifts also held for those anglers that were not interviewed. A specific $W2$ was calculated for each surveyor (u) operating on each surveying date (k) at each landing site (a):

$$W2_{aku} = \frac{LF_{aku} + LM_{aku} \left(\frac{LF_{aku}}{LF_{aku} + LN_{aku}} \right)}{LF_{aku} - ri_{aku}}. \quad (\text{Equation 2})$$

where LF_{aku} was the number of interviewed boats that were fishing (a subset of these, ri_{aku} , refused an interview, or had an incomplete interview that could not be used), LN_{aku} was the number of interviewed boats that were not fishing, and LM_{aku} was the number of boats that were not interviewed by surveyor u , during surveying date k , at landing site a . This equation incorporates the assumption that interviewed boats were representative, allowing us to estimate the portion of missed boats that were fishing.

We used the term $A_{pdsakut}$ to denote the number of non-lodge boats reporting activity during time-block t , as interviewed by surveyor u on survey date k , at landing site a , in subarea s , during period p , and on day type d (n_{pdsaku} was used to denote the total number of boats that reported activity during *any* time-block). The two correction factors were applied, and the data were summed over surveyors, survey dates and landing sites:

$$A'_{pdst} = \sum_k \sum_u \sum_a (W1_{pda} \cdot W2_{aku} \cdot A_{pdsakut}). \quad (\text{Equation 3})$$

Summing the adjusted number of anglers over the hourly time-blocks gave:

$$T'_{pds} = \sum_t A'_{pdst}. \quad (\text{Equation 4})$$

The proportion of non-lodge boats (P_{pdst}) that were actively fishing during each of the hourly time-blocks (t) was calculated for each period, day type, and subarea:

$$P_{pdst} = \frac{A'_{pdst}}{\sum_k \sum_u \sum_a (W1_{pda} \cdot W2_{aku} \cdot n_{pdsaku})} \quad . \quad (\text{Equation 5})$$

Using this method, 120 unique angler activity patterns were to be estimated (i.e., 5 periods $\times$ 2 day types $\times$ 12 subareas). To reliably describe angler activity, a relatively large number of boats (~60) needed to be interviewed in each of the 120 blocks. In the end, some blocks contained too few interviews (Table 2), so it was decided to pool activity data over subareas. The equation for non-lodge angler activity was thus:

$$P_{pdt} = \frac{\sum_s A'_{pdst}}{\sum_s \sum_k \sum_u \sum_a (W1_{pda} \cdot W2_{aku} \cdot n_{pdsaku})} \quad , \quad (\text{Equation 6})$$

and the associated variance was:

$$S^2_{P_{pdt}} = \frac{(P_{pdt})(1 - P_{pdt})}{\sum_s \sum_k \sum_u \sum_a (W1_{pda} \cdot W2_{aku} \cdot n_{pdsaku})} \quad . \quad (\text{Equation 7})$$

Table 2. The amount of data (number of complete interviews) available to estimate non-lodge angler activity patterns, for all levels of each factor, 2025.

		Subarea												TOTALS		
Period	Day Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall
1-28 May	WD	3	0	5	56	19	31	10	1	34	2	2	6	8	161	169
	WE	1	2	6	73	32	38	22	3	34	2	10	2	9	216	225
	Total	4	2	11	129	51	69	32	4	68	4	12	8	17	377	394
29 May - 22 June	WD	9	22	4	94	28	41	39	3	76	12	17	8	35	318	353
	WE	12	24	8	133	47	66	47	5	110	18	24	3	44	453	497
	Total	21	46	12	227	75	107	86	8	186	30	41	11	79	771	850
23 June - 23 July	WD	35	39	15	271	20	174	30	6	106	48	93	3	89	751	840
	WE	16	27	13	163	26	129	42	4	104	27	36	5	56	536	592
	Total	51	66	28	434	46	303	72	10	210	75	129	8	145	1,287	1,432
24 July - 17 August	WD	35	46	43	179	47	129	86	4	190	25	25	7	124	692	816
	WE	19	50	19	155	46	106	70	11	109	11	37	12	88	557	645
	Total	54	96	62	334	93	235	156	15	299	36	62	19	212	1,249	1,461
18-31 August	WD	34	22	27	74	22	47	21	3	40	11	22	11	83	251	334
	WE	10	4	25	43	12	28	10	0	35	15	15	2	39	160	199
	Total	44	26	52	117	34	75	31	3	75	26	37	13	122	411	533
GRAND TOTAL		174	236	165	1241	299	789	377	40	838	171	281	59	575	4,095	4,670

Catch Per Unit Effort Estimation

Catch per unit effort (numbers of fish kept or released per boat trip) of non-lodge boats was estimated for each species from interviews of non-lodge anglers. Catch per unit effort was calculated separately for guided and non-guided boats, and overall; but only the overall estimates were used in subsequent calculations. For each interview (i), the guided status (g), period (p), and the catch (C) of each species (r) in each subarea (s) was recorded. Given that each interview represented one boat trip, then catch per unit effort (CPUE), in terms of catch per boat trip, was calculated as:

$$CPUE_{psrgi} = \frac{C_{psrgi}}{1} \quad \text{where } g \in \{\text{all, guided, not-guided}\}. \quad (\text{Equation 8})$$

Mean $CPUE$ was calculated as the average catch, over all n_{psg} boat trips:

$$\widehat{CPUE}_{psrg} = \frac{\sum_{i=1}^{n_{psg}} C_{psrgi}}{n_{psg}}. \quad (\text{Equation 9})$$

The variance for the estimate of mean catch per unit effort was calculated as:

$$S_{CPUE_{psrg}}^2 = \frac{\sum (CPUE_{psrgi} - \widehat{CPUE}_{psrg})^2}{(n_{psg} - 1)}. \quad (\text{Equation 10})$$

In the event that a period/subarea-specific sample size was < 5 , adjacent subareas would be pooled together as follows:

- if n for 3I, 3J or 3K was too low, then it was combined with the other two subareas, along with data from ambiguous parts of Area 3 (“3?”) to give a 3?IJK regional average $CPUE$;
- if n for 4A, 4B or 4C was too low, then it was combined with the other two to give a 4ABC regional average $CPUE$;
- if n for 4D, 4E or 4F was too low, then it was combined with the other two to give a 4DEF regional average $CPUE$; and
- if n for 4G, 4H or 4L was too low, then it was combined with the other two to give a 4GHL regional average $CPUE$.

Similar methods were used to calculate the average number of fish harvested (retained) per boat trip (HPUE), and the average number of fish released per boat trip (RPUE).

Angler Effort Estimation

To obtain statistically valid estimates of non-lodge fishing effort, boats were counted during overflight surveys conducted from a fixed-wing aircraft traveling along a set path over the study area (see Figure 2). In order to calculate the non-lodge effort, the estimated counts of active lodge boats needed to be subtracted from the overflight counts for that particular day.

Lodge Effort Census

A complete census of lodge-based angling effort was collected from lodge logbooks by DFO personnel. DFO used the following procedure to estimate the number of active lodge vessels that would have been available to be observed during the flights: 1) Lodges reported the number of active vessels for each flight date. 2) No lodge-based vessels were considered available to the flights (zero visible) on change-over days (days when the lodges received new clients; on typical change-over days, departing clients leave before the flight and new clients arrive after the flight).

Change-over days were often different for each lodge. 3) Lodges reported their lunchtime activity to DFO, which ranged from virtually always going in for a mid-day break to rarely leaving the water during the day. Based on this information, it was assumed that 60% of the vessel activity was available to be counted on non-changeover flight dates. 4) The number of vessels visible were summed over the active lodges.




Figure 2. Map showing planned flight path of effort survey overflights. Letters indicate subareas. Flights originated and ended in Prince Rupert and flew in the clockwise direction. On good weather days, the flights also included a foray into subarea 4L (path not shown).

None of the lodge effort reported from Area 3 included subarea-specific location information. Thus, lodge effort was partitioned among the Area 3 subareas, see below.

Non-lodge Effort Surveys

Table 3 shows the number of overflight surveys conducted for each day type and month (or period). Each survey was designed to allow visual inspection of the subareas through which the aircraft passed. Each time the aircraft entered a subarea, the start and end times were recorded, along with the numbers of small (< 30 ft) and large ($\geq$ 30 ft) recreational vessels that were observed, tallied separately depending on whether or not they were actively fishing at the time.

Table 3. Overflight sampling effort (number of flights shifts) by day type and month (left) or period (right) in 2025.

Month	Weekday	Weekend/ Holiday	Total	Period	Weekday	Weekend/ Holiday	Total
May	4	3	7	1-28 May	3	2	5
June	5	3	8	29 May - 22 June	4	2	6
July	5	3	8	23 June - 23 July	6	4	10
August	5	3	8	24 July - 17 August	2	3	5
Total	19	12	31	18-31 August	4	1	5
				Total	19	12	31

During survey o (conducted during period p and on day type d), observers tallied the total number of recreational boats that were actively fishing at time t on their i^{th} pass over subarea s , V_{pdsoit} . For each survey, the relative proportion of visible boats within Management Area 3 was calculated, ρ_{pdsoit} , and the total number of lodge boats (λ_{pdo}) apportioned accordingly:

$$\lambda_{pdsoit} = \rho_{pdsoit} \cdot \lambda_{pdo} \quad . \quad (\text{Equation 11})$$

The number of non-lodge boats was hence:

$$v_{pdsoit} = \max \{0, V_{pdsoit} - \lambda_{pdsoit}\} \quad . \quad (\text{Equation 12})$$

Since angling occurs over the course of the entire day, the numbers of non-lodge boats were divided by the proportion of average daily number of boats active (P_{pdst}) during the time block when the observations were recorded. Since the flight path (Figure 2) brought the observers into some subareas more than once during the survey (covering different ground, thus the tallies were considered additive), the adjusted tallies were summed over all visits, to calculate the total number of active non-lodge fishing boats on the day of the survey, by subarea, B_{pdso} :

$$B_{pdso} = \sum_i \frac{v_{pdsoit}}{P_{pdst}} \quad . \quad (\text{Equation 13})$$

The variance of B_{pdso} was calculated using the standard formulas for combining the variance of products and quotients of two independent random variables (Goodman 1960):

$$\begin{aligned} \text{if } z = x/y, \text{Var}(z) &= (y^{-2})\text{Var}(x) + (x^2y^{-4})\text{Var}(y) ; \\ \text{if } z = xy, \text{Var}(z) &= (y^2)\text{Var}(x) + (x^2)\text{Var}(y) ; \text{ and} \\ \text{if } z = qx, \text{Var}(z) &= (q^2)\text{Var}(x) \quad \text{where } q \text{ is a constant.} \end{aligned} \quad (\text{Equation 14})$$

Since there was no variance associated with the boat counts, it followed that

$$S_{B_{pdso}}^2 = v_{pdsoit}^2 \cdot (P_{pdst})^{-4} \cdot S_{P_{pdst}}^2 \quad . \quad (\text{Equation 15})$$

Next, the survey-specific ‘daily effort’ estimates were averaged over the number of surveys conducted, n_{pds} , as:

$$\hat{B}_{pds} = \frac{\sum_{o=1}^{n_{pds}} B_{pdso}}{n_{pds}} \quad (\text{Equation 16})$$

with a variance of:

$$S_{\hat{B}_{pds}}^2 = (n_{pds})^{-2} \cdot S_{B_{pdso}}^2 \quad . \quad (\text{Equation 17})$$

Total fishing effort for each sub-period was calculated for each subarea by multiplying the period-specific average daily effort (e.g., $\hat{B}$ for the ‘23 June – 23 July’ period was applied to both

the ‘23-30 June’ and ‘1-23 July’ sub-periods, see Table 1) by the number days of day type d that occurred in sub-period c :

$$E_{cds} = \hat{B}_{pds} \cdot N_{cd} \quad , \quad (\text{Equation 18})$$

which was then summed over day type to give:

$$E_{cs} = \sum_d \hat{B}_{pds} \cdot N_{cd} \quad , \quad (\text{Equation 19})$$

with a variance of:

$$S_{E_{cs}}^2 = \sum_d S_{\hat{B}_{pds}}^2 \cdot N_{cd}^2 \quad . \quad (\text{Equation 20})$$

Total monthly fishing effort was calculated by summing over the appropriate sub-periods (i.e., total effort for July was the sum of effort in the ‘1-23 July’ and ‘24-31 July’ sub-periods). As always, variances for sums were additive (Goodman 1960).

Because of inclement weather, not all overflights surveyed every subarea. Especially for the more exposed subareas, it was possible to have no overflight data in a given combination of period and day type. In this event, subarea-specific effort was scaled using: 1) the proportion of effort in that subarea, relative to that in a three-subarea cluster (see pooling of adjacent subareas for CPUE estimation, above), measured during overflights for which all three subareas were surveyed; and 2) the effort in the cluster’s other two subareas, estimated for the combination of period and day type during which the data gap occurred.

Rarely, this method produced an effort estimate of zero in a stratum for which interview data suggested otherwise. As expected, when breaking the sturdy duration into subperiods, the issue was more common. In 2025, no boats were counted but interviews showed that effort had occurred during 1-28 May weekdays in Subareas 3K, 4B, 4E, 4D, 4H, 4G and 4L; during 1-28 May weekends in Subarea 4E; during 1-22 June weekends in Subareas 4G and 4L; during 24-31 July and 1-17 August weekdays in Subarea 4H; and during 18-31 August weekends in Subareas 3J, 4A, and 4B. In these cases, the interview data obtained were taken as representative of total effort.

Catch and Harvest Estimation

Total catch was calculated for each sub-period, subarea, and species by multiplying sub-period-specific angling effort by period-specific catch per unit effort, and then summing over day type:

$$C_{csr} = \sum_d (E_{cds} \cdot C\hat{P}U_{psrg}) \quad , \quad (\text{Equation 21})$$

where g = ‘all’ (see Equation 8). The standard errors for these catch estimates were:

$$S_{C_{csr}} = \sqrt{\sum_d \left(E_{cds}^2 \frac{S_{C\hat{P}U_{psrg}}^2}{n_{psg}} + C\hat{P}U_{psrg}^2 \frac{S_{E_{cds}}^2}{n_{pds}} + \frac{S_{C\hat{P}U_{psrg}}^2}{n_{psg}} \frac{S_{E_{cds}}^2}{n_{pds}} \right)} \quad , \quad (\text{Equation 22})$$

where n_{pds} and n_{psg} are the sample sizes for E and CPUE estimates, respectively. Total monthly catch was calculated by summing over the appropriate sub-periods, with additive variances.

Similar methods were used to estimate monthly numbers of fish harvested, and monthly numbers of fish released by replacing CPUE with HPUE or with RPUE in Equations 21 and 22.

Biosampling

When given permission by the anglers, surveyors collected biological data from Chinook and Coho salmon and Pacific Halibut. Biosampled species were sexed when they were intact. Chinook and Coho salmon were examined for clipped adipose fins (mark rate data used by DFO for coded wire tag analyses) and when possible, heads from clipped fish were collected. The flesh color was noted for Chinook Salmon. Scale samples were collected from Chinook Salmon, and tissue samples (fin clips) were collected from Coho Salmon (used by DFO to determine age, and to carry out genetic stock identification).

Ideally, all anglers would allow their catch to be biosampled, and the examined fish would represent an unbiased sample of all harvested fish. But Robichaud et al. (2016) theorized that the interview protocol, used until the end of 2015, may have resulted in bias. The protocol had surveyors ask anglers about adipose clips before asking permission to observe the fish. Anglers were often engaged in a conversation about the purpose of the fin-clips and how the data would be used, and it appeared that anglers with clipped fish were more likely to permit handling and biosampling of catch than when no clipped fish were present. To avoid this bias, we modified our data collection protocol in 2016 so that anglers were asked if their catch could be biosampled before any other interactions took place. Also starting 2016, no fin clip data were accepted on the word of the anglers: the only fin-clip data we used were those which came from observers when they handled and inspected the catch.

Biomass of Harvested Pacific Halibut

Halibut lengths were converted to weights using the International Pacific Halibut Commission (Clark 1992) regression equations:

$$\begin{aligned}\text{Weight(lb)}_{\text{net}} &= 6.921 \times 10^{-6} \times \text{FL(cm)}^{3.24} \\ \text{Weight(lb)}_{\text{round}} &= 9.166 \times 10^{-6} \times \text{FL(cm)}^{3.24}\end{aligned}\quad (\text{Equation 23})$$

Converting average length to average weight allowed harvested biomass to be calculated from harvested fish counts.

Statistical Analyses

For each fish taxa, differences in CPUE among periods were examined using Kruskal Wallis tests (Zar 1984), whereas differences between trip-types (guided vs. non-guided) were examined using paired t-tests (i.e., the difference between guided vs unguided catch for a given subarea and period was tested against a null of zero; Sokal and Rohlf 1995). Kruskal Wallis tests were used to examine the effects of period and day type on effort, and the effects of month on catch and harvest. When statistically significant effects were observed, *post-hoc* Kruskal-Wallis tests were used to compare between each possible pair, where the false discovery rate was controlled using the Benjamini–Hochberg procedure (Benjamini and Hochberg 1995).

Trailer Census

The total number of parked trailers was recorded daily at the landing locations during the peak of the activity timing window (i.e., between 11:00 AM and 1:30 PM; similar to the targeted timing of aerial overflights). Theoretically, trailer counts could be used as an index of fishing effort, are more economical than overflights, and are not weather dependant, though no sub-area-specific

effort data are collected. DFO has been collecting trailer count data for years and we are maintaining the time series. Linear regression was used to determine the relationship between trailer counts and fishing effort. A data point was generated each time a trailer census occurred on the same day as an overflight. For the regression analysis, the location-specific trailer counts and non-lodge boat counts were used as the independent and dependent variables, respectively.

In addition to every day during the May through August study, trailer count data were also collected throughout April. The above linear regression was used to calculate from the daily trailer counts what the daily boat counts *would have been* had there been overflights on each day. Day type-specific activity pattern data were used to expand the daily estimated 'boat counts' to calculate total daily fishery effort (early May activity data were used to expand both the April and May trailer count numbers, otherwise period-specific activity pattern data were used). Summing over all days in each month produced an *index* of fishing effort for each month. A scaler was calculated as the ratio of the April index value to the sum of the May to August indices. Next, that May to August creel-based non-lodge effort values were multiplied by the scaler to estimate April non-lodge fishing effort. The same analysis was repeated using trailer count data from four other years (i.e., 2019, 2020, 2023, and 2024) in which April trailer count data were collected (using the regression line and the activity pattern data from those years, see Robichaud and Addison 2020, 2021, 2024, 2025).

RESULTS

Angler Interviews

Over the four month study period, there were 261 observer shifts, during which 2,961 interviews were conducted, involving 9,836 non-lodge anglers (Figure 3; Table 4). July was the busiest month, with 1,096 interviews conducted (average 17.1 interviews per shift), which represented 37% of the total number of study period interviews (Table 4).

Non-lodge boat trip activity was unevenly distributed among the subareas, with 4A being the favourite (Table 5). Subareas 4F and 4C were the next most likely area for anglers to have reported activity, followed by 4D and 4B. The subareas with the lowest number of trips reporting activity were 4E and 4L. Only 147 interviews were of parties that only targeted shellfish, and those were overwhelmingly in subarea 4A (Table 6).

On average, 33% of the non-lodge boat trips were guided by a professional. The guiding rate varied by landing site. Only 82 of 753 trips (11%) interviewed at Port Edward were guided, whereas trips departing from Rushbrooke and Prince Rupert Yacht Club were guided 35% and 49% of the time, respectively (Table 7).

In all there were 139 refusals (Table 5), or 4.7% refusal rate. The highest refusal rate (5.1%) was at the Rushbrooke boat launch, where 45% of the refusals occurred. The refusal rates for the Port Edward boat launch and the Prince Rupert Yacht Club were 4.9% and 4.1% (27% and 28% of the refusals), respectively.

Angler Activity Patterns

Non-lodge angler activity pattern estimates varied little among periods (Figure 4). There was virtually no difference between the weekday and weekend/holiday activity patterns. Timing of our overflights lined up with peak activity levels for all flights, with the possible exception of the very latest weekend flight in August.




Figure 3. Number of interviews conducted at each of three landing sites, by month, 2025.

Table 4. Monthly number of shifts, and numbers of boats interviewed per landing site, 2025. Non-lodge effort is expressed in units of boat trips. Sampling efficiency is also calculated.

Landing Site	May				June				Jul			
	# ints	# anglers	# stints	ints/stint	# ints	# anglers	# stints	ints/stint	# ints	# anglers	# stints	ints/stint
Port Edward Boat Launch	58	163	22	2.6	142	401	22	6.5	319	1062	22	14.5
Prince Rupert Yacht Club	99	304	21	4.7	202	657	22	9.2	342	1229	20	17.1
Rushbrooke Boat Launch	144	424	22	6.5	287	906	22	13.0	435	1469	22	19.8
TOTAL	301	891	65	4.6	631	1,964	66	9.6	1,096	3,760	64	17.1
% of Total Interviews			10%				21%				37%	
Estimated Non-lodge Effort			995				2,410				3,979	
% of Cumulative Non-lodge Effort			8.4%				20%				34%	
% of Effort Sampled			30%				26%				28%	

Landing Site	August				TOTAL			
	# ints	# anglers	# stints	ints/stint	# ints	# anglers	# stints	ints/stint
Port Edward Boat Launch	234	727	22	10.6	753	2353	88	8.6
Prince Rupert Yacht Club	319	1164	22	14.5	962	3354	85	11.3
Rushbrooke Boat Launch	380	1330	22	17.3	1246	4129	88	14.2
TOTAL	933	3,221	66	14.1	2,961	9,836	261	11.3
% of Total Interviews			32%					
Estimated Non-lodge Effort			4,419				11,803	
% of Cumulative Non-lodge Effort			37%					
% of Effort Sampled			21%				25%	

Table 5. Fishing locations as reported by interviewed non-lodge parties, by month and subarea, 2025. For interviews that reported activity in multiple subareas, the boat trip tally was evenly partitioned among the subareas visited.

<i>Month</i>	<i>Subarea</i>															<i>TOTAL</i>
	<i>3I</i>	<i>3J</i>	<i>3K</i>	<i>4A</i>	<i>4B</i>	<i>4C</i>	<i>4D</i>	<i>4E</i>	<i>4F</i>	<i>4G</i>	<i>4H</i>	<i>4L</i>	<i>3?</i>	<i>4?</i>	<i>Refusal</i>	
<i>May</i>	3	3	9	91	38	47	27	3	46	2	7	5		2	18	301
<i>June</i>	15	38	15	142	60	81	60	5	125	23	32	5	1	1	28	631
<i>July</i>	32	39	22	239	55	234	85	8	203	43	75	9	0	5	48	1,096
<i>August</i>	52	66	68	174	54	131	92	7	156	28	43	16	1	1	45	933
<i>TOTAL</i>	102	146	113	646	207	493	265	23	530	95	157	35	2	8	139	2,961

Table 6. Numbers of non-lodge boat trips that reported only fishing for shellfish, by month and subarea, 2025.

	Subarea														
Month	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	3?	4?	TOTAL
May	0	0	0	27	0	0	0	0	0	0	0	0	0	0	27
June	0	0	0	27	3	2	0	0	0	1	0	0	0	0	33
July	0	0	0	55	1	2	0	0	1	1	0	0	0	0	59
August	0	0	0	23	3	2	0	0	0	0	0	0	0	0	28
TOTAL	0	0	0	132	7	6	0	0	1	2	0	0	0	0	147

Table 7. Numbers of guided (i.e., led by professional guides) and unguided boat trips, by landing site, 2025.

<i>Landing Site</i>	<i>Trip Type</i>		<i>% guided</i>
	<i>Guided</i>	<i>Unguided</i>	
<i>Port Edward Boat Launch</i>	82	671	11%
<i>Prince Rupert Yacht Club</i>	468	494	49%
<i>Rushbrooke Boat Launch</i>	433	813	35%
<i>TOTAL</i>	983	1,978	33%




Figure 4. Angler activity pattern (proportion recreational angling boats that were actively fishing during each hour-long time block) by day type and period, pooled over all subareas, 2025. Shaded regions bound the earliest and latest flight times for each day type (blue = weekday; orange = weekend/holiday) and period.

Catch Per Unit Effort Estimates

Sample sizes were adequate ($n \geq 5$) to produce reliable estimates of non-lodge CPUE in most combinations of periods and creel study sub-areas (Table 8), except in subareas 3I, 3J, 4E, and 4G from 1-28 May, and in subarea 4E from 18-31 August. However, when the interview data were split between guided and non-guided groups, period-specific sample sizes did not suffice for several more combinations of period and management area (Table 8). In those cases where sample sizes were limited, data from three adjacent subareas were pooled together for CPUE calculations (Table 9).

Average number of fish harvested per non-lodge boat trip (HPUE) was calculated for each species by period, subarea, and by trip type (Appendix 2).

HPUE varied significantly among periods (Figure 5, Table 10) for Chinook ($\chi^2 = 29.3$, $df = 4$, $P < 0.0001$), Coho ($\chi^2 = 37.4$, $df = 4$, $P < 0.0001$), Pink (*Oncorhynchus gorbuscha*, $\chi^2 = 32.7$, $df = 4$, $P < 0.0001$), and Sockeye salmon (*Oncorhynchus nerka*, $\chi^2 = 16.8$, $df = 4$, $P = 0.002$), and also for Yellowtail Rockfish (*Sebastes flavidus*, $\chi^2 = 11.7$, $df = 4$, $P = 0.020$). Chinook Salmon HPUE was significantly lower during the 23 June to 23 July period, when retention was illegal, as compared to all other periods (Table 10). Coho Salmon HPUE increased from over time, peaked in the periods from 23 June to 17 August, and declined thereafter (Table 10; Figure 5). HPUE for Pink Salmon was significantly lower during the early periods (1 May to 22 June), as compared to the later periods (23 June to 21 August; Table 10). For Sockeye Salmon and Yellowtail Rockfish, among-period differences in HPUE were not statistically significant after controlling for the rate of false positives (Table 10). There were no significant differences for any other species examined (including Pacific Halibut and Lingcod; Figure 5).

Guided trips typically had higher HPUE than non-guided trips, and the differences were statistically significant for a variety of species (Table 11). In particular, guided trips harvested 0.5 more Chinook Salmon ($t_{47} = 5.0$, $P < 0.0001$), and 1.3 more Coho Salmon ($t_{47} = 5.1$, $P < 0.0001$) than unguided trips. In addition, guided trips harvested 0.7 more Lingcod ($t_{47} = 4.7$, $P < 0.0001$), 0.5 more Pacific Cod (*Gadus macrocephalus*, $t_{47} = 3.2$, $P = 0.003$), and 0.5 more Pacific Halibut per trip ($t_{47} = 5.7$, $P < 0.0001$) than unguided trips. Guided trips also harvested significantly more Black (*Sebastes melanops*), Bocaccio (*S. paucispinis*), Canary (*S. pinniger*), Dusky (*S. ciliatus*), Quillback (*S. maliger*), Tiger (*S. nigrocinctus*), and Yellowtail rockfish than unguided trips (Table 11).

For lodge boat trips, HPUE (and RPUE for Chinook Salmon) values are shown in Table 12.

Table 8. Sample sizes for CPUE (catch per unit effort) estimates by period, for guided and not guided non-lodge boat trips, 2025.

		Subarea											
Period	Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
1-28 May	Guided	0	1	2	58	7	22	10	0	33	0	7	8
	Unguided	4	1	9	71	44	47	22	4	35	4	5	0
	Combined	4	2	11	129	51	69	32	4	68	4	12	8
29 May - 22 June	Guided	12	19	3	131	17	23	31	2	82	15	24	8
	Unguided	9	27	9	96	58	84	55	6	104	15	17	3
	Combined	21	46	12	227	75	107	86	8	186	30	41	11
23 June - 23 July	Guided	26	39	9	223	4	114	17	3	109	31	75	7
	Unguided	25	27	19	211	42	189	55	7	101	44	54	1
	Combined	51	66	28	434	46	303	72	10	210	75	129	8
24 July - 17 August	Guided	39	55	19	161	15	64	23	4	109	16	32	13
	Unguided	15	41	43	173	78	171	133	11	190	20	30	6
	Combined	54	96	62	334	93	235	156	15	299	36	62	19
18-31 August	Guided	32	16	24	67	5	17	3	0	27	11	20	10
	Unguided	12	10	28	50	29	58	28	3	48	15	17	3
	Combined	44	26	52	117	34	75	31	3	75	26	37	13

Table 9. Subarea data (pooled over several adjacent areas, in some cases) used to calculate CPUE in each period and subarea, for guided and not guided non-lodge boat trips, 2025.

		Subarea											
Period	Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
1-28 May	Guided	3?,I,J,K	3?,I,J,K	3?,I,J,K	4A	4B	4C	4D	4D,E,F	4F	4G,H,L	4H	4L
	Unguided	3?,I,J,K	3?,I,J,K	3K	4A	4B	4C	4D	4D,E,F	4F	4G,H,L	4H	4G,H,L
	Combined	3?,I,J,K	3?,I,J,K	3K	4A	4B	4C	4D	4D,E,F	4F	4G,H,L	4H	4L
29 May - 22 June	Guided	3I	3J	3?,I,J,K	4A	4B	4C	4D	4D,E,F	4F	4G	4H	4L
	Unguided	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4G,H,L
	Combined	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
23 June - 23 July	Guided	3I	3J	3K	4A	4A,B,C	4C	4D	4D,E,F	4F	4G	4H	4L
	Unguided	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4G,H,L
	Combined	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
24 July - 17 August	Guided	3I	3J	3K	4A	4B	4C	4D	4D,E,F	4F	4G	4H	4L
	Unguided	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
	Combined	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
18-31 August	Guided	3I	3J	3K	4A	4B	4C	4D,E,F	4D,E,F	4F	4G	4H	4L
	Unguided	3I	3J	3K	4A	4B	4C	4D	4D,E,F	4F	4G	4H	4G,H,L
	Combined	3I	3J	3K	4A	4B	4C	4D	4D,E,F	4F	4G	4H	4L




Figure 5. Average number of fish harvested per non-lodge boat trip, for six sportfish species, by period and subarea, 2025. Data are for guided and unguided trips combined. Note that Y axis scale varies among plots.

Table 10. Kruskal-Wallis tests for differences in HPUE (fish harvested per non-lodge boat trip) among periods, by species, 2025. Subareas treated as replicates. CLD: red letters indicate statistical differences within species (periods not connected by a common letter were statistically different).

Species		Kruskal-Wallis Statistics			Median HPE					CLD
		χ^2	df	P	1-28 May	29 May - 22 Jun	23 Jun - 23 Jul	24 Jul - 17 Aug	18-31 Aug	
Chinook Salmon	<i>Oncorhynchus tshawytscha</i>	29.3	4	< 0.0001	0.7	0.7	0	0.5	0.4	A-A-B-A-A
Chum Salmon	<i>Oncorhynchus keta</i>	4.0	4	0.41	0	0	0	0	0	
Coho Salmon	<i>Oncorhynchus kisutch</i>	37.4	4	< 0.0001	0	0.09	3.4	2.6	0.8	D-C-A-A-B
Pink Salmon	<i>Oncorhynchus gorbuscha</i>	32.7	4	< 0.0001	0	0	0.4	0.3	0.08	B-B-A-A-A
Sockeye Salmon	<i>Oncorhynchus nerka</i>	16.8	4	0.002	0	0	0	0.002	0	A-A-A-A-A
Arrow-tooth Flounder	<i>Atheresthes stomias</i>	6.3	4	0.18	0	0	0	0	0	
Big Skate	<i>Beringraja binoculata</i>	4.0	4	0.41	0	0	0	0	0	
Cabezon	<i>Scorpaenichthys marmoratus</i>	0.5	4	0.97	0	0	0	0	0	
English Sole	<i>Parophrys vetulus</i>	4.0	4	0.41	0	0	0	0	0	
Greenlings	<i>Hexagrammos</i> spp.	3.5	4	0.48	0	0	0.002	0.002	0	
Lingcod	<i>Ophiodon elongatus</i>	0.7	4	0.95	0.6	0.4	0.4	0.6	0.7	
Longnose Skate	<i>Caliraja rhina</i>	0.0	4	1	0	0	0	0	0	
Pacific Cod	<i>Gadus macrocephalus</i>	2.4	4	0.66	0.08	0.05	0.1	0.2	0.2	
Pacific Halibut	<i>Hippoglossus stenolepis</i>	0.9	4	0.92	0.9	0.9	1.0	0.8	1.0	
Pacific Rock Sole	<i>Lepidopsetta bilineata</i>	8.9	4	0.06	0	0	0	0	0	
Pacific Spiny Dogfish	<i>Squalus suckleyi</i>	3.9	4	0.42	0	0	0	0	0	
Sablefish	<i>Anoplopoma fimbria</i>	3.1	4	0.55	0	0	0	0	0	
Spotted Ratfish	<i>Hydrolagus colliei</i>	0.0	4	1	0	0	0	0	0	
Starry Flounder	<i>Platichthys stellatus</i>	0.0	4	1	0	0	0	0	0	
Black Rockfish	<i>Sebastes melanops</i>	1.5	4	0.83	0.2	0.05	0.1	0.1	0.2	
Boccacio Rockfish	<i>Sebastes paucispinis</i>	6.2	4	0.18	0	0	0	0	0	
Canary Rockfish	<i>Sebastes pinniger</i>	8.6	4	0.07	0	0	0	0	0	
China Rockfish	<i>Sebastes nebulosus</i>	1.9	4	0.75	0	0	0.005	0.005	0	
Copper Rockfish	<i>Sebastes caurinus</i>	0.2	4	1.00	0.005	0.01	0.007	0.008	0	
Dusky Rockfish	<i>Sebastes ciliatus</i>	7.4	4	0.12	0	0	0	0	0	
Quillback Rockfish	<i>Sebastes maliger</i>	7.0	4	0.14	0.02	0.05	0.1	0.2	0.2	
Redbanded Rockfish	<i>Sebastes babcocki</i>	8.1	4	0.09	0	0	0	0	0	
Redstripe Rockfish	<i>Sebastes proriger</i>	0.0	4	1	0	0	0	0	0	
Tiger Rockfish	<i>Sebastes nigrocinctus</i>	4.4	4	0.35	0	0	0	0.02	0.004	
Widow Rockfish	<i>Sebastes entomelas</i>	3.6	4	0.46	0	0	0	0	0	
Yelloweye Rockfish	<i>Sebastes ruberrimus</i>	8.1	4	0.09	0	0	0	0	0	
Yellowtail Rockfish	<i>Sebastes flavidus</i>	11.7	4	0.020	0	0	0.02	0.03	0.06	A-A-A-A-A

Table 11. Paired t-tests for differences in HPUE (fish harvested per non-lodge boat trip) between guided and non-guided boats, by species, 2025. Periods and subareas treated as replicates.

<i>Species</i>	<i>Paired t-test Statistics</i>			<i>Median HPE</i>		<i>Paired Mean Difference</i>
	<i>t</i>	<i>df</i>	<i>P</i>	<i>Guided</i>	<i>Not</i>	
Chinook Salmon	5.0	47	< 0.0001	0.6	0.2	0.5
Chum Salmon	-1.0	47	0.32	0	0	-0.0001
Coho Salmon	5.1	47	< 0.0001	2.1	0.9	1.3
Pink Salmon	1.2	47	0.23	0.1	0.06	0.06
Sockeye Salmon	0.3	47	0.76	0	0	0.0007
Arrow-tooth Flounder	0.2	47	0.86	0	0	0.0001
Big Skate	1.4	47	0.17	0	0	0.0004
Cabezon	1.3	47	0.20	0	0	0.004
English Sole	1.0	47	0.32	0	0	0.0007
Greenlings	0.5	47	0.62	0	0	0.005
Lingcod	4.7	47	< 0.0001	1.0	0.3	0.7
Longnose Skate	0.0	47	1	0	0	0
Pacific Cod	3.2	47	0.003	0.2	0.07	0.5
Pacific Halibut	5.7	47	< 0.0001	1.4	0.8	0.5
Pacific Rock Sole	1.5	47	0.14	0	0	0.008
Pacific Spiny Dogfish	-1.0	47	0.32	0	0	-0.0011
Sablefish	1.2	47	0.23	0	0	0.006
Spotted Ratfish	0.0	47	1	0	0	0
Starry Flounder	0.0	47	1	0	0	0
Black Rockfish	3.3	47	0.002	0.2	0.06	0.3
Boccacio Rockfish	2.1	47	0.046	0	0	0.01
Canary Rockfish	2.3	47	0.024	0	0	0.01
China Rockfish	1.8	47	0.08	0	0	0.01
Copper Rockfish	1.8	47	0.09	0	0	0.02
Dusky Rockfish	2.5	47	0.018	0	0	0.02
Quillback Rockfish	4.4	47	< 0.0001	0.2	0.07	0.2
Redbanded Rockfish	0.0	47	1	0	0	0
Redstripe Rockfish	0.0	47	1	0	0	0
Tiger Rockfish	3.9	47	0.0003	0	0	0.03
Widow Rockfish	-2.0	47	0.06	0	0	-0.0022
Yelloweye Rockfish	0.0	47	1	0	0	0
Yellowtail Rockfish	2.9	47	0.006	0.03	0	0.05

Table 12. Average number of harvested fish (and for Chinook Salmon, released) per boat trip, by month, from the census of Area 3 lodges, 2025.

<i>Species</i>	<i>May</i>	<i>June</i>	<i>July</i>	<i>August</i>	<i>Total</i>
<i>Chinook Salmon:</i>					
<i>retained</i>	0.9	0.5	0.2	0.2	0.3
<i>released</i>	0.3	0.09	0.5	0.2	0.3
<i>Coho Salmon</i>	0.02	0.4	2.9	2.5	1.8
<i>Pink Salmon</i>	0	0.0068	0.005	0.01	0.008
<i>Chum Salmon</i>	0	0.001	0	0	0.0004
<i>Sockeye Salmon</i>	0	0	0	0.004	0.001
<i>Pacific Halibut</i>	0.3	0.3	0.4	0.3	0.3
<i>Lingcod</i>	1.0	0.8	0.9	0.8	0.8
<i>Other groundfish</i>	0	0	0	0	0
<i>Yelloweye Rockfish</i>	0	0	0	0	0
<i>Other rockfish</i>	0.2	0.3	0.3	0.4	0.3

Angler Effort Estimates

Over the four month study period, a total of 31 effort surveys were conducted, including 19 weekday surveys and 12 conducted on weekends/holidays (Table 3, Appendix 3). The numbers of lodge boats that were visible during the overflights (Appendix 3) were partitioned among the subareas in Management Area 3, and subtracted from the overflight counts of small boats. The remaining small boat counts (“the non-lodge small boat counts”) were significantly correlated to trailer counts at both survey launch sites (Rushbrooke: $F_{1,29} = 42.6$, $P < 0.0001$; Port Edwards: $F_{1,29} = 51.8$, $P < 0.0001$), and when the two launch sites were combined ($F_{1,29} = 61.7$, $P < 0.0001$). The fit of the combined-site regression ($R^2 = 0.67$; Figure 6) was lower in 2025 than it has been in prior years (e.g., 0.81, Robichaud and Addison 2025), mainly as a result of two dates (23 and 29 July) when considerably more boats were counted than would have been expected given the trailer counts – flight crews on neither date indicated that anything was unusual, each time only noting sunny weather and calm seas.




Figure 6. Linear regressions showing the relationships between small non-lodge boats counted during overflights and trailers concurrently counted at landing sites. Shaded areas show the 95% confidence interval of the predicted values. Note that X axis scale varies among plots.

Monthly angling effort for each subarea in 2025 is shown in Figure 7 and in Table 13, and for all months combined in Figure 8. Total lodge effort in 2025 was 2,384 boat trips (168 in May, 740 in June, 747 in July, and 729 in August; Table 13). To calculate total (lodge + non-lodge) fishing effort by subarea, lodge boat trips were distributed among the three subareas in Management Area 3 according to the relative non-lodge effort in each (Table 13; Figure 7). In all, there were 1,163 (SE = 12) boat trips in May, 3,150 (SE = 12) boat trips in June, 4,726 (SE = 17) boat trips in July, and 5,148 (SE = 20) boat trips in August. The full-season (May-August) Area 3-4 effort estimate for 2025 was 14,187 boat trips (SE = 30). For comparison with our previous study years, which did not include surveys in May, the June-August effort total was 13,024 boat trips (SE = 29).

Median daily non-lodge effort on weekdays (3.9 trips per day per subarea) was lower than on weekends/holidays (6.0 trips per day per subarea) but the effect was not statistically significant ($\chi^2 = 1.8$, $df = 1$, $P = 0.18$). Effort varied significantly among periods ($\chi^2 = 23.1$, $df = 4$, $P = 0.00012$). Effort in 1-28 May (median 2.0 trips per day per subarea) was significantly lower than that in each of the other four periods (medians: 29 May to 22 June, 5.2; 23 June to 23 July, 6.2; 24 July to 17 August, 12.8; 18 to 31 August, 5.5 trips per day per subarea), and also Period Two (29 May to 22 June) had significantly less effort per day than did Period Four (24 July to 17 August).




Figure 7. Total estimated numbers of boat trips, by month and subarea, 2025.

Table 13. Lodge, non-lodge, and total effort estimates (boat trips per month), by month and subarea, 2025.

Month		Subarea												TOTALS			SE of the Totals		
		3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4I	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May	Non-lodge Estimate	85.0	45.7	63.8	86.0	194.1	140.4	71.9	14.4	206.5	27.6	44.3	15.9	195	801	995	3	12	12
	Lodge Report	73.4	39.5	55.1										168	0	168			
	TOTAL MAY	158.5	85.2	118.8	86.0	194.1	140.4	71.9	14.4	206.5	27.6	44.3	15.9	363	801	1,163	3	12	12
	SE	2.0	1.1	1.5	2.6	6.9	4.0	2.2	0.2	8.0	1.1	1.7	0.5						
June	Non-lodge Estimate	156.0	203.3	290.2	95.4	150.4	311.6	152.6	83.4	592.0	137.3	209.6	28.0	650	1,760	2,410	8	10	12
	Lodge Report	177.7	231.6	330.7										740	0	740			
	TOTAL JUNE	333.6	434.9	620.9	95.4	150.4	311.6	152.6	83.4	592.0	137.3	209.6	28.0	1,390	1,760	3,150	8	10	12
	SE	2.6	2.6	6.7	1.5	1.9	2.7	2.1	1.5	7.8	1.8	2.6	0.6						
July	Non-lodge Estimate	220.3	486.9	288.2	72.5	190.5	679.1	255.5	38.0	1043.5	321.7	311.2	71.2	995	2,983	3,979	6	16	17
	Lodge Report	165.3	365.4	216.3										747	0	747			
	TOTAL JULY	385.7	852.3	504.5	72.5	190.5	679.1	255.5	38.0	1043.5	321.7	311.2	71.2	1,742	2,983	4,726	6	16	17
	SE	1.9	4.6	3.0	0.7	1.7	4.7	2.7	0.5	14.0	3.1	3.7	1.8						
August	Non-lodge Estimate	344.0	588.1	492.3	137.2	164.5	758.3	435.5	36.3	879.3	328.1	201.4	54.6	1,424	2,995	4,419	11	17	20
	Lodge Report	176.1	301.0	252.0										729	0	729			
	TOTAL AUGUST	520.1	889.0	744.2	137.2	164.5	758.3	435.5	36.3	879.3	328.1	201.4	54.6	2,153	2,995	5,148	11	17	20
	SE	4.0	7.5	7.1	1.0	1.8	8.0	5.4	0.6	12.2	4.0	4.1	0.9						
FULL SEASON TOTAL		1,398	2,261	1,988	391	699	1,889	915	172	2,721	815	767	170	5,648	8,539	14,187	15	28	32
FULL SEASON SE		6	9	10	3	8	10	7	2	22	5	6	2						




Figure 8. Heat map showing the estimated total effort (numbers of boat trips, lodge + non-lodge) in each subarea in 2025. See Table 13 for details.

April Effort

In 2019, 2020, 2023, 2024, and 2025 trailer counts began in April, and ran until the end of the creel monitoring period. The annual trailer-to-boat-count regression line (see Figure 6 for 2025) was used to calculate what the daily boat counts would have been had there been overflights each day, and the proportion of active anglers (specific to year, month, day type, and time of trailer count, see Figure 4 for 2025) was used to expand the boat counts to calculate total fishery effort by day. Summing over all the days of each month produced trailer count derived monthly fishing effort indices (Figure 9), which compared well to those calculated using creel methods for non-lodge anglers (Figure 10; adjusted $r^2 = 0.88$), although the slope differed significantly from the 1:1 line ($F_{1,18} = 8.6$, $P = 0.009$). The tilt of the trendline away from the 1:1 line is largely the result of data from July 2025 when considerably more boats were counted during two overflights than would have been expected given the trailer counts. Estimates suggested that an additional 8-11% of fishing effort is occurring in April, that is non accounted for by conducting a creel survey from May to August. This amounts to fishing trips numbering from 766 to 1,303 (1,296 trips in 2025) depending on the year (Table 14).




Figure 9. Total monthly non-lodge fishing effort as estimated using the usual creel methods (blue) or from trailer counts (orange; estimated using a combination of trailers counted at landing sites, the year-specific trailer-to-boat count regression, and the year-, month- and day type-specific activity pattern data). Creel-based estimates are not available for April (thus 'na').




Figure 10. Relationship between total monthly non-lodge fishing effort values as estimated using the usual creel methods (Y axis) or from trailer counts (X axis). Blue line shows the line of best fit from a linear regression, and the thin black diagonal shows the 1:1 line.

Table 14. Estimated April non-lodge fishing effort in the Areas 3 and 4 in 2019, 2020, 2023, 2024, and 2025, along with data used in calculation steps. A: The index of effort for April as calculated using trailer count data, the count-to-boat regression, and angler activity profile data. B: The trailer count effort indices for May to August, summed. A/B: the ratio of A to B, used as a scaler. C: the creel-derived published estimates of non-lodge effort, summed over May to August. A/B(C): estimates of April non-lodge fishing effort, calculated by applying the index scale to the creel-derived effort estimates from the May to August study period.

	<i>A</i>	<i>B</i>	<i>A/B</i>	<i>C</i>	<i>A/B(C)</i>
	<i>April Index</i>	<i>May-Aug Index</i>	<i>Ratio of April to May-Aug Indices</i>	<i>May-Aug Non-lodge Creel Estimate</i>	<i>April Non-lodge Effort (scaled)</i>
2019	1,133	10,777	11%	12,397	1,303
2020	646	7,884	8%	9,353	766
2023	1,010	9,989	10%	10,062	1,017
2024	1,009	11,022	9%	11,928	1,092
2025	1,115	10,162	11%	11,803	1,296

Catch Estimates

Monthly and full-season estimates were made of the numbers of fish released and retained, by species (Tables 15-24; Appendix 4). Non-lodge estimates were generated from the product of effort and CPUE estimates. Lodge catches were extracted from census documents that were submitted to DFO by the lodges, and which were then presented by DFO to NCSFNSS with identifying information removed for Privacy Act reasons. The results presented in this section are total (lodge + non-lodge) numbers of fish retained and released.

An estimated 7,279 (SE = 152) Chinook Salmon were retained by anglers in Areas 3 or 4 during the 2025 study period (Table 16; Figure 11). The largest proportional harvest came from subareas 4F (31%), 3J (19%), and 3K (10%). Harvest was highest in August (42% of total harvest), although among-month differences were not statistically significant ($\chi^2 = 3.3$, $df = 3$, $P = 0.34$; Table 25). Harvest in both June and July were significantly curtailed by the no-retention period from 23 June to 23 July ($\chi^2 = 27.4$, $df = 4$, $P < 0.0001$; Appendix 5). The number of Chinook Salmon released during the study was estimated at 11,466 (SE = 370), or 61% of the total catch. Percent retention (calculated for non-lodge anglers) was 55% in May, 39% in June, 18% in July, and 56% in August (Appendix 4).

The salmonid that was harvested in the greatest numbers during the 2025 study period was, by far, Coho Salmon (38,082 fish retained, SE = 543; Table 17; Figure 12). The largest proportional harvest came from subarea 4F (23%), followed by subareas 3J (18%), 3K (14%), 4C (14%), and 3I (14%). Harvest differed significantly among months ($\chi^2 = 22.4$, $df = 3$, $P < 0.0001$; Table 25): harvest in May was significantly lower than all other months, and harvest in June was significantly lower than July. By period, Coho Salmon harvest was significantly higher during the two periods from 23 June to 17 August than during previous or subsequent periods ($\chi^2 = 35.5$, $df = 4$, $P < 0.0001$; Appendix 5). During the study period, an estimated 2,964 (SE = 230) Coho Salmon were released by anglers (7% of the total catch).

Of the main three salmon species caught in 2025, Pink Salmon was the least likely to be retained (Table 18). We estimate that 15,857 (SE = 416) fish were caught, of which 20% were retained (3,249 fish, SE = 135), and 80% were released (12,607 fish, SE = 393). The largest proportional harvest came from subarea 4F (28%), followed by subareas 4C (16%), 3K (16%), 3I (14%), and 3J (12%). Harvest of Pink Salmon varied significantly among months ($\chi^2 = 22.9$, $df = 3$, $P < 0.0001$; Table 25): harvest in May was significantly lower than all other months. By period, Pink Salmon harvest peaked in the period from 24 July to 17 August ($\chi^2 = 32.5$, $df = 4$, $P < 0.0001$; Appendix 5).

Very few Chum (*Oncorhynchus keta*, 4 fish, SE = 2, mostly in 4F in August) or Sockeye (56 fish, SE = 12, mainly in 4F in July, and in Area 3 or 4H in August) salmon were harvested in Area 3 and 4 during the study period (Tables 19 and 20). Some interviews included catch recorded as 'unknown salmon species', resulting in an estimated 73 (SE = 21) harvested fish.

Table 15. Total (lodge + non-lodge) estimated harvest, by subarea, for popular sportfish taxa, 2025.

Species	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
<i>Chinook Salmon</i>	660	1,356	758	5	376	686	652	46	2,239	299	157	46	2,774	4,505	7,279	97	118	152
<i>Coho Salmon</i>	5,157	6,964	5,407	0	584	5,209	2,104	160	8,597	1,482	1,911	506	17,528	20,554	38,082	363	404	543
<i>Pink Salmon</i>	455	391	507	0	31	532	222	43	896	61	102	8	1,353	1,896	3,249	105	86	135
<i>Chum Salmon</i>	0	0	0	0	0	0	0	0	3	0	0	0	1	3	4	0	2	2
<i>Sockeye Salmon</i>	12	8	6	0	0	0	4	0	21	0	5	0	27	29	56	7	11	12
<i>Unknown Salmon</i>	0	0	0	0	0	0	7	3	17	7	0	39	0	73	73	0	21	21
<i>Pacific Halibut</i>	1,009	2,156	894	2	268	1,320	533	228	3,250	684	1,186	394	4,058	7,865	11,923	157	155	221
<i>Lingcod</i>	439	3,967	619	3	162	372	254	267	2,446	874	325	366	5,025	5,070	10,094	220	207	302
<i>Other Groundfish *</i>	409	245	128	0	71	230	119	9	614	542	1,696	239	783	3,521	4,303	104	241	262
<i>Rockfish †</i>	467	2,426	531	2	246	627	505	160	2,574	703	382	483	3,424	5,683	9,107	153	206	257

* includes Big and Longnose skates, Pacific Spiny Dogfish, Pacific Cod, Sablefish, Cabezon, Red Irish Lord, greenlings, Spotted Ratfish, Arrow-tooth and Starry flounders, Pacific Sanddab, Pacific Rock and English soles, and unknown groundfish

† includes Black, Boccacio, Canary, China, Copper, Dusky, Quillback, Redband, Redstripe, Tiger, Widow, Yelloweye, Yellowtail and unknown rockfish

Table 16. Total (lodge + non-lodge) estimated Chinook Salmon catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	126.6	75.1	88.3	4.7	242.7	167.1	100.5	9.3	377.2	11.1	20.3	8.3	290	941	1,231	20	61	64
Released	130.5	78.0	132.9	1.7	260.5	111.0	85.9	3.6	182.3	8.3	10.0	0.4	341	664	1,005	68	81	106
June																		
Retained	253.5	248.4	243.4	0.7	73.2	102.9	148.9	18.6	649.2	35.4	59.5	9.3	745	1,098	1,843	60	47	76
Released	258.7	420.9	502.1	0.1	104.7	452.9	139.0	9.0	768.3	84.5	175.7	15.8	1,182	1,750	2,931	193	81	210
July																		
Retained	64.5	331.5	97.7	0	21.5	111.7	130.5	1.6	324.7	69.3	15.2	7.9	494	682	1,176	28	35	45
Released	529.7	1007.0	517.7	0.4	102.1	1105.0	203.7	26.0	1064.4	188.5	403.5	38.0	2,054	3,132	5,186	163	165	232
August																		
Retained	215.7	701.1	328.6	0	38.5	304.0	272.2	16.5	887.8	183.2	61.5	20.0	1,245	1,784	3,029	68	82	106
Released	327.9	527.6	373.9	0	21.7	261.8	144.4	5.0	518.8	88.7	70.4	2.6	1,229	1,114	2,343	131	103	167
TOTAL																		
Retained	660	1,356	758	5	376	686	652	46	2,239	299	157	46	2,774	4,505	7,279	97	118	152
Released	1,247	2,033	1,527	2	489	1,931	573	44	2,534	370	660	57	4,807	6,659	11,466	293	226	370
Catch	1,907	3,390	2,284	8	865	2,616	1,225	90	4,773	669	816	102	7,581	11,164	18,745	309	255	400

Table 17. Total (lodge + non-lodge) estimated Coho Salmon catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	5.2	9.5	12.4	0	0.8	0.8	2.5	0.0	6.8	1.0	0	0.4	27	12	39	9	2	9
Released	0	0	0	0	0	0	0.2	0	1.4	0.3	0	0	0	2	2	0	1	1
June																		
Retained	400.4	487.1	270.9	0	58.4	912.7	74.4	31.3	928.7	154.3	376.0	92.6	1,158	2,628	3,787	91	98	134
Released	77.5	2.8	9.3	0	8.3	52.2	5.8	0	80.3	10.7	7.7	0	90	165	255	43	27	51
July																		
Retained	2111.6	3075.2	1533.7	0	265.3	3088.6	777.3	100.6	4383.9	718.4	1152.0	322.3	6,721	10,809	17,529	210	316	379
Released	283.3	184.5	65.2	0	32.5	184.8	72.9	0	312.2	86.0	29.1	0	533	717	1,250	132	92	161
August																		
Retained	2639.9	3392.2	3590.2	0	259.5	1206.5	1249.5	28.5	3277.7	608.3	383.6	91.1	9,622	7,105	16,727	281	233	365
Released	200.0	395.8	281.1	0	19.1	100.6	123.6	0.5	187.7	125.6	23.2	0	877	580	1,457	134	80	156
TOTAL																		
Retained	5,157	6,964	5,407	0	584	5,209	2,104	160	8,597	1,482	1,911	506	17,528	20,554	38,082	363	404	543
Released	561	583	356	0	60	338	203	1	582	223	60	0	1,500	1,465	2,964	193	125	230
Catch	5,718	7,547	5,763	0	644	5,546	2,306	161	9,179	1,705	1,971	506	19,028	22,018	41,046	411	423	590

Table 18. Total (lodge + non-lodge) estimated Pink Salmon catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	0	0	0	0	0	0.4	0	0	0.3	0	0	0	0	1	1	0	0	0
Released	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
June																		
Retained	22.3	52.9	50.8	0	0	73.4	14.9	10.7	92.3	5.7	15.3	0	126	212	338	28	19	34
Released	90.0	136.9	83.6	0	15.6	282.8	20.7	17.9	213.1	61.4	67.4	0	311	679	989	58	59	82
July																		
Retained	155.2	204.3	209.4	0	9.2	279.9	92.0	31.2	456.7	28.6	51.7	2.0	569	951	1,520	80	63	102
Released	542.0	926.2	524.6	0	94.8	1132.9	342.8	55.2	1401.4	340.9	222.5	9.2	1,993	3,600	5,593	189	194	271
August																		
Retained	277.3	133.7	247.2	0	21.8	178.6	114.7	1.4	346.8	27.0	35.1	6.2	658	732	1,390	61	55	83
Released	780.4	1192.8	736.2	0	111.0	746.0	592.6	7.5	1417.9	318.7	99.9	22.5	2,709	3,316	6,025	211	174	273
TOTAL																		
Retained	455	391	507	0	31	532	222	43	896	61	102	8	1,353	1,896	3,249	105	86	135
Released	1,412	2,256	1,344	0	221	2,162	956	81	3,032	721	390	32	5,013	7,595	12,607	289	267	393
Catch	1,867	2,647	1,852	0	252	2,694	1,178	124	3,928	782	492	40	6,366	9,491	15,857	307	281	416

Table 19. Total (lodge + non-lodge) estimated Chum Salmon catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Released	0.7	0.3	0	0	0	0	0	0	0.5	0.3	0	0	1	1	2	1	0	1
June																		
Retained	0.3	0.3	0.3	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0
Released	5.3	6.6	0	0	0	2.0	0.4	0	7.1	2.5	0	0	12	12	24	5	4	6
July																		
Retained	0	0	0	0	0	0	0	0	1.2	0	0	0	0	1	1	0	1	1
Released	0	12.1	12.8	0	0	6.8	2.2	0	13.4	0	0	0	25	22	47	9	6	11
August																		
Retained	0	0	0	0	0	0	0	0	2.3	0	0	0	0	2	2	0	2	2
Released	3.1	0	31.5	0	0	2.3	2.4	0	6.8	0	0	0	35	11	46	12	4	12
TOTAL																		
Retained	0	0	0	0	0	0	0	0	3	0	0	0	1	3	4	0	2	2
Released	9	19	44	0	0	11	5	0	28	3	0	0	73	47	119	15	8	18
Catch	9	19	45	0	0	11	5	0	31	3	0	0	74	50	124	15	9	18

Table 20. Total (lodge + non-lodge) estimated Sockeye Salmon catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Released	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
June																		
Retained	0	0	0	0	0	0	0	0	4.2	0	0	0	0	4	4	0	3	3
Released	0	2.8	0	0	0	0	0	0	0	0	0	0	3	0	3	2	0	2
July																		
Retained	3.3	2.3	1.8	0	0	0	1.2	0	14.3	0	0.5	0	7	16	23	3	10	10
Released	0	10.4	0	0	0	1.0	3.5	0	2.4	0	0	0	10	7	17	6	4	7
August																		
Retained	9.0	5.8	4.5	0	0	0	2.4	0	2.3	0	4.5	0	19	9	28	6	4	7
Released	0	4.9	4.9	0	0	2.3	7.1	0	4.5	0	0	0	10	14	24	6	7	9
TOTAL																		
Retained	12	8	6	0	0	0	4	0	21	0	5	0	27	29	56	7	11	12
Released	0	18	5	0	0	3	11	0	7	0	0	0	23	21	44	9	8	12
Catch	12	26	11	0	0	3	14	0	28	0	5	0	50	50	100	11	13	17

Table 21. Total (lodge + non-lodge) estimated Pacific Halibut catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	117.6	54.9	27.4	1.7	10.6	67.0	33.8	14.2	267.7	41.6	84.3	33.8	200	555	755	25	40	47
Released	66.4	31.9	0	0	17.9	77.0	11.2	10.0	168.8	27.6	54.9	11.3	98	379	477	28	59	65
June																		
Retained	295.9	261.6	291.1	0.1	60.2	187.5	104.9	91.5	821.6	157.9	342.2	66.7	849	1,833	2,681	86	67	109
Released	178.5	210.0	39.5	0	30.5	119.2	35.3	63.9	478.5	193.9	341.9	82.7	428	1,346	1,774	73	94	119
July																		
Retained	258.3	760.6	318.9	0.2	120.1	455.5	146.1	68.6	1295.7	299.2	432.5	174.4	1,338	2,992	4,330	85	98	130
Released	114.2	785.5	157.4	0	19.3	290.4	29.2	35.4	774.8	473.3	627.1	229.7	1,057	2,479	3,536	121	207	240
August																		
Retained	337.6	1078.6	256.3	0	76.8	610.4	247.9	53.9	864.7	185.2	327.1	118.9	1,672	2,485	4,157	97	91	133
Released	138.6	1220.7	121.1	0	28.4	313.8	72.0	31.9	654.9	469.2	316.4	159.2	1,480	2,046	3,526	195	225	297
TOTAL																		
Retained	1,009	2,156	894	2	268	1,320	533	228	3,250	684	1,186	394	4,058	7,865	11,923	157	155	221
Released	498	2,248	318	0	96	800	148	141	2,077	1,164	1,340	483	3,064	6,249	9,313	243	325	405
Catch	1,507	4,404	1,212	2	364	2,121	680	369	5,327	1,848	2,526	877	7,122	14,114	21,236	289	360	462

Table 22. Total (lodge + non-lodge) estimated Lingcod catch, by subarea

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	139.5	119.9	47.5	0	51.8	23.9	4.9	18.5	289.1	37.3	5.2	54.7	307	485	792	44	68	81
Released	0	1.0	0	0	21.2	0	0.6	5.4	15.1	4.2	1.1	0	1	48	49	1	15	15
June																		
Retained	47.7	846.1	258.2	0	25.1	71.3	33.3	118.5	551.9	151.9	71.5	47.3	1,152	1,071	2,223	108	88	140
Released	0.9	43.9	23.2	0	0	5.1	6.6	40.8	44.6	55.8	8.9	0	68	162	230	33	39	51
July																		
Retained	89.6	1541.3	148.5	0	32.4	166.7	73.1	70.8	868.1	349.9	118.6	112.9	1,779	1,792	3,572	110	119	162
Released	2.6	137.3	62.5	0	0	20.6	9.6	10.4	122.2	126.1	4.3	0.7	202	294	496	88	79	118
August																		
Retained	162.3	1459.5	164.6	2.9	52.5	109.9	142.7	59.5	737.0	335.4	130.0	151.2	1,786	1,721	3,508	150	128	197
Released	0	82.2	0	0	1.3	19.6	13.4	2.2	64.8	135.1	5.8	1.3	82	243	326	30	92	97
TOTAL																		
Retained	439	3,967	619	3	162	372	254	267	2,446	874	325	366	5,025	5,070	10,094	220	207	302
Released	3	264	86	0	23	45	30	59	247	321	20	2	353	747	1,100	98	128	162
Catch	443	4,231	704	3	184	417	284	326	2,693	1,196	345	368	5,378	5,816	11,195	241	244	342

Table 23. Total (lodge + non-lodge) estimated catch of other groundfish by subarea Included are Big and Longnose skates, Pacific Spiny Dogfish, Pacific Cod, Sablefish, Cabezon, Red Irish Lord, greenlings, Spotted Ratfish, Arrow-tooth and Starry flounders, Pacific Sanddab, Pacific Rock and English soles, and other miscellaneous groundfishes.

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	28.6	9.1	8.8	0	7.2	2.3	1.8	0.6	39.6	44.0	143.5	0.4	46	239	286	15	56	58
Released	8.9	3.6	0	0	6.0	29.3	20.7	0.8	22.3	1.7	8.2	0	13	89	101	5	23	24
June																		
Retained	77.5	15.5	20.9	0	7.6	22.2	23.7	0	77.8	178.9	646.3	35.7	114	992	1,106	37	120	125
Released	28.4	0	0	0.4	24.0	34.6	26.5	0	47.7	9.0	85.6	0	28	228	256	15	27	31
July																		
Retained	104.5	94.9	63.5	0.1	29.4	87.5	46.0	0.8	218.5	199.4	746.8	109.9	263	1,438	1,701	55	157	167
Released	86.7	18.3	31.1	1.1	19.6	67.9	22.2	0	181.6	34.6	177.5	0	136	504	641	44	65	79
August																		
Retained	198.5	126.0	34.9	0.2	26.9	117.7	47.9	7.2	278.4	119.7	159.1	93.4	359	850	1,210	79	125	148
Released	45.7	39.3	84.3	0	6.5	27.8	45.2	2.3	182.1	15.5	69.0	0	169	348	518	44	57	72
TOTAL																		
Retained	409	245	128	0	71	230	119	9	614	542	1,696	239	783	3,521	4,303	104	241	262
Released	170	61	115	2	56	160	114	3	434	61	340	0	346	1,169	1,516	64	94	113
Catch	579	307	244	2	127	389	234	12	1,048	603	2,036	239	1,129	4,690	5,819	122	258	286

Table 24. Total (lodge + non-lodge) estimated rockfish catch, by subarea Included are Black, Boccacio, Canary, China, Copper, Dusky, Quillback, Redbanded, Redstripe, Tiger, Widow, Yelloweye, and Yellowtail rockfish, and other miscellaneous rockfishes.

Month / Catch Type	Subarea												TOTALS			SE of the Totals		
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	Area 3	Area 4	Overall	Area 3	Area 4	Overall
May																		
Retained	23.3	57.5	5.8	0	61.9	65.2	19.8	15.4	213.7	33.4	13.3	67.3	87	490	577	9	62	63
Released	5.9	8.7	0	3.5	198.2	71.6	19.0	12.8	155.2	22.0	15.6	24.7	15	523	537	4	61	61
June																		
Retained	119.2	393.0	126.2	0.2	63.6	102.5	55.5	86.5	504.9	84.7	133.3	76.0	638	1,107	1,745	55	74	92
Released	52.8	169.1	67.4	0	88.3	165.6	100.9	74.5	405.9	145.6	80.0	33.5	289	1,094	1,384	68	89	112
July																		
Retained	159.6	830.8	291.8	0.7	83.4	226.1	166.8	18.2	958.0	247.7	119.6	206.4	1,282	2,027	3,309	88	133	159
Released	75.4	436.8	248.8	0	104.4	271.5	202.5	13.6	874.1	441.4	121.2	105.9	761	2,135	2,896	173	166	240
August																		
Retained	164.9	1144.5	107.1	1.2	37.2	233.1	263.1	40.4	897.9	337.4	115.6	133.3	1,416	2,059	3,476	112	125	168
Released	146.9	420.4	387.5	0.5	29.5	338.0	254.5	42.4	808.2	397.5	190.0	62.4	955	2,123	3,078	166	167	236
TOTAL																		
Retained	467	2,426	531	2	246	627	505	160	2,574	703	382	483	3,424	5,683	9,107	153	206	257
Released	281	1,035	704	4	420	847	577	143	2,243	1,007	407	227	2,020	5,874	7,894	250	259	360
Catch	748	3,461	1,235	6	667	1,473	1,082	304	4,818	1,710	789	710	5,443	11,558	17,001	293	331	442




Figure 11. Total (lodge + non-lodge) estimated Chinook Salmon harvest, by month and subarea, 2025.




Figure 12. Total (lodge + non-lodge) estimated Coho Salmon harvest, by month and subarea, 2025.

Table 25. Kruskal Wallis tests for differences in harvest (fish caught per boat trip per subarea) among months, by species, 2025. Subareas treated as replicates. CLD: red letters indicate statistical differences within species (months not connected by a common letter were statistically different).

Species	Kruskal-Wallis Statistics			Median Monthly Harvest				CLD
	χ^2	df	P	May	June	July	August	
Chinook Salmon	3.3	3	0.34	41.4	88.1	59.4	186.5	
Chum Salmon	2.0	3	0.56	0	0	0	0	
Coho Salmon	22.4	3	< 0.0001	0.9	179.7	904.4	907.4	C-B-A-AB
Pink Salmon	22.9	3	< 0.0001	0	15.1	71.9	74.9	B-A-A-A
Sockeye Salmon	12.1	3	0.007	0	0	0.3	1.1	B-AB-A-A
Arrow-tooth Flounder	3.2	3	0.36	0	0	0	0	
Big Skate	2.0	3	0.56	0	0	0	0	
Cabezon	0.6	3	0.90	0	0	0	0	
English Sole	2.0	3	0.56	0	0	0	0	
Greenlings	2.2	3	0.53	0	0.3	1.5	0.9	
Lingcod	10.6	3	0.014	30.6	71.4	103.0	121.2	B-AB-A-A
Longnose Skate	0.0	3	1	0	0	0	0	
Pacific Cod	8.4	3	0.038	3.0	19.8	73.1	95.7	A-A-A-A
Pacific Halibut	13.3	3	0.004	37.7	172.7	222.1	233.3	B-A-A-A
Pacific Rock Sole	4.6	3	0.20	0	0	0	0	
Pacific Spiny Dogfish	2.3	3	0.51	0	0	0	0	
Sablefish	1.0	3	0.79	0	0	0	0	
Spotted Ratfish	0.0	3	1	0	0	0	0	
Starry Flounder	0.0	3	1	0	0	0	0	
Black Rockfish	5.5	3	0.14	12.7	21.4	43.8	44.8	
Boccacio Rockfish	7.3	3	0.06	0	0.4	2.6	0.2	
Canary Rockfish	4.9	3	0.18	0	0	0	0	
China Rockfish	5.3	3	0.15	0.06	1.6	6.3	3.8	
Copper Rockfish	1.8	3	0.61	0.5	3.5	3.1	5.4	
Dusky Rockfish	0.7	3	0.88	0	0	0	0	
Quillback Rockfish	13.2	3	0.004	3.5	17.5	39.9	46.3	B-AB-A-A
Redbanded Rockfish	6.1	3	0.11	0	0	0	0	
Redstripe Rockfish	0.0	3	1	0	0	0	0	
Tiger Rockfish	8.1	3	0.045	0.3	0	1.9	5.7	A-A-A-A
Widow Rockfish	0.5	3	0.91	0	0	0	0	
Yelloweye Rockfish	6.1	3	0.11	0	0	0	0	
Yellowtail Rockfish	13.2	3	0.004	0.10	3.2	9.8	12.6	B-A-A-A

An estimated 11,923 (SE = 221) Pacific Halibut were harvested in Areas 3 or 4 during the 2025 study period (Table 21; Figure 13). Most of the harvest was attributed to subareas 4F (27%), followed by 3J (18%), 4C (11%), and 4H (10%). Harvest varied significantly among months ($\chi^2 = 13.3$, df = 3, $P = 0.004$; Table 25), and was significantly lower in May than in any other month. In all, 9,313 (SE = 405) Pacific Halibut were released, representing 44% of the total catch.

From May to August 2025, an estimated 10,094 (SE = 302) Lingcod were harvested, occurring mainly from subareas 3J (39%) and 4F (24%; Table 22; Figure 14). Harvest varied significantly among months ($\chi^2 = 10.6$, df = 3, $P = 0.014$; Table 25), and was significantly lower in May than in July or August. By period, Lingcod harvest peaked in the period from 24 July to 17 August ($\chi^2 = 11.8$, df = 4, $P = 0.019$; Appendix 5). Of the captured Lingcod, 10% were released (1,100 fish, SE = 162).




Figure 13. Total (lodge + non-lodge) estimated Pacific Halibut harvest, by month and subarea, 2025.




Figure 14. Total (lodge + non-lodge) estimated Lingcod harvest, by month and subarea, 2025.

An estimated 4,303 (SE = 262) fish from other groundfish species were harvested from May to August 2025, of which 91% (i.e., 3,912 fish, SE = 257) were Pacific Cod. The remaining harvest included greenlings (*Hexagrammos* spp.; 6%), and a variety of other species (<50 individuals retained per species). Over the study period, it is estimated that 313 (SE = 57) Pacific Cod were released (7% of the total catch of Pacific Cod).

Rockfish harvest was estimated at 9,107 fish (SE = 257; Table 24), composed mostly of Black (34%), Quillback (22%), Yellowtail (5%), Tiger (5%), China (*Sebastes nebulosus*, 4%), and Copper (*S.s caurinus*, 3%), and unknown (24%) rockfish, along with seven other rockfish species. Harvests of Quillback ($\chi^2 = 13.2$, df = 3, $P = 0.004$; Table 25) and Yellowtail rockfish ($\chi^2 = 13.2$, df = 3, $P = 0.004$; Table 25) were significantly higher in the later parts of the study period. In all, 7,894 (SE = 360) were released. Of the 1,715 Yelloweye Rockfish caught, all but 3 (99.8%) were released.

Table 26. Biosampling results, 2025.

	Chinook Salmon		Coho Salmon		Pacific Halibut	
	Number	Percent	Number	Percent	Number	Percent
Number Examined	448		530		524	
Subarea						
3I	7	2%	62	12%	11	2%
3J	23	5%	37	7%	18	4%
3K	10	2%	29	5%	29	6%
4A	4	1%				
4B	60	13%	22	4%	36	7%
4C	67	15%	194	37%	139	27%
4D	85	19%	59	11%	59	11%
4E	2	0%	2	0%	7	1%
4F	180	40%	78	15%	147	29%
4G	7	2%	24	5%	24	5%
4H	3	1%	23	4%	40	8%
4L					4	1%
Sex						
Male	1	20%				
Female	4	80%				
Undeterminable						
Unknown	443		530		524	
Trip Type						
Guided	203	45%	286	54%	189	36%
Unguided	245	55%	244	46%	335	64%
Adipose Clips						
Clipped	100	23%	1	0%		
Unmarked	340	77%	515	100%		
Unknown						
Head Tagged						
Yes	85	19%	1	0.2%		
No	363	81%	529	100%		
Scale Samples			Whatman Samples			
Yes	442	99%	530	100%		
No	6	1%				
Flesh Colour						
Red	50	89%				
White	5	9%				
Marble	1	2%				
Unknown						

Biosampling

During interviews, observers were told about 2,043 retained Chinook Salmon, and were permitted to handle 448 fish. Too few fish were sexed ($n=5$) to provide a robust estimate of sex ratio (Table 26). The median fork length of biosampled Chinook Salmon ($n = 446$) was 70 cm (95% of observations were between 53 and 92 cm; Figure 15). In all, 100 adipose-clipped individuals were encountered (Table 26), including 34 from standardized sampling methods (complete samples collected from anglers who were asked if their catch could be biosampled before any other interactions took place; Appendix 6) and 66 from non-standard interactions (“non-standard interactions” were interviews for which anything other than complete access was given to all harvested fish). In all, 85 Chinook Salmon had their heads collected and labelled for submission to the Salmon Head Recovery Program (Table 26). Of the standardized samples ($n=198$), 17.2% were adipose-clipped (Appendix 6). Scale samples were obtained from 99% ($n=480$) of the biosampled Chinook Salmon (Table 26). In May, 131 Chinook Salmon were handled, including scales sampled from 129 fish, and 23 heads collected (Appendix 7).

For Coho Salmon, observers were told about 9,205 retained fish, and were permitted to handle 530 fish. None of the fish were sexed (Table 26). The median Coho Salmon fork length ($n = 520$) was 64 cm (95% of observations were between 52 and 72 cm; Figure 15). In all, 1 adipose-clipped individual was encountered (Table 26), none from standardized sampling interviews. One head was collected for submission to the Salmon Head Recovery Program (Table 26). Of the standardized samples ($n=165$), 0% were adipose-clipped (Appendix 6). Tissue samples were obtained from all ($n=530$) of the biosampled Coho Salmon (Table 26).

Of the 3,258 Pacific Halibut that anglers reported as retained, 524 were handled (Table 26). All were measured, but none were sexed. The median length was 78 cm (95% of observations were between 56 and 99 cm; Figure 15). Average net and round weights, calculated from average lengths using regression formulas, were 4.6 kg (10.1 lb) and 6.1 kg (13.4 lb) in Area 3, and 4.4 kg (9.7 lb) and 5.8 kg (12.9 lb) in Area 4, respectively (Table 27).

Table 27. Average weight, and harvest (numbers and biomass) of Pacific Halibut in Area 3 and 4, 2025.

	Stat Area 3				Stat Area 4			
	Net Weight		Round Weight		Net Weight		Round Weight	
	kg	lb	kg	lb	kg	lb	kg	lb
Creel Interview (Non-Lodge) Average Weights								
Guided	4.9	10.8	6.5	14.3	5.1	11.2	6.8	14.9
Unguided	4.3	9.5	5.7	12.6	4.0	8.9	5.3	11.8
Overall	4.6	10.1	6.1	13.4	4.4	9.7	5.8	12.9
Estimated Non-lodge Harvest (number of fish)								
Overall			3,193				7,852	
Estimated Lodge Harvest (number of fish)								
Overall			827				0	
Biomass Harvested								
Non-lodge	14,612	32,214	19,352	42,663	34,581	76,239	45,799	100,969
Lodge	3,784	8,343	5,012	11,049	0	0	0	0
Overall	18,396	40,557	24,364	53,713	34,581	76,239	45,799	100,969




Figure 15. Length-frequency distributions of biosampled Chinook (n = 446) and Coho (n = 520) salmon, and Pacific Halibut (n = 524), 2025. Note that X and Y axis scales vary among plots.

Harvested Halibut Biomass

Based on the average weights of biosampled Pacific Halibut, and the estimated harvest, it was estimated that total harvested Pacific Halibut net biomass was 53.0 tonnes (116,796 lb), including 18.4 t (40,557 lb) in Area 3 and 34.6 t (76,239 lb) in Area 4. In terms of round weight, Pacific Halibut harvest was 70.2 t (154,681 lb) including 24.4 t (53,713 lb) in Area 3 and 45.8 t (100,969 lb) in Area 4 (Table 27).

DISCUSSION

Creel Estimates

In general, recreational fishing effort and salmon catch in Areas 3 & 4 have been increasing over the last 31 years (Table 28 and Figure 16). Using only published creel data collected since 1995, we found 17 years (1995-96, 2001-02, 2008-09¹, and 2015-2025) for which effort data were collected in June, July, and August (Van Tongeren and Winther 2010, Robichaud et al. 2016, 2017, Robichaud and Addison 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025), and we have therefore limited this meta analyses to those three months. Meta-analysis revealed (Figure 16) that fishing effort has increased significantly (by 186.4 trips per year on average; $F_{1,15} = 21.6$, $P = 0.0003$) since 1995. Effort in 2024 and 2025 were close to the expected value given the long-term trend, whereas effort from 2020-2023 was lower than what would have been predicted.




Figure 16. Fishing effort from 1995 to 2025. Annual estimates only include data from June, July and August. Only years with data from all three months are included. Long-term linear regression equation is shown in bottom right of the panel.

¹ Creel data from 2010-2014 were collected and analyzed by DFO, and preliminary results were presented as part of their post-season review process. We based our meta-analysis on published accounts (written accounts showing methods, results, and discussing the caveats of the results), and hence have not included data from these years.

Over the same three-decade period, Coho Salmon harvests have increased significantly by 870 fish per year, on average ($F_{1,15} = 14.2$, $P = 0.002$, Table 29, Figure 17). The 2025 Coho Salmon harvest was very close to that expected given the long-term trend.

The number of Chinook Salmon being released by anglers has been increasing strongly since 2009 (the first year with data available; ln scale, $F_{1,10} = 35.3$, $P = 0.0001$; Figure 18). Chinook Salmon harvest had been on an increasing trend until 2019, but now appears to be declining over time. The long-term trend (39.5 additional fish per year on average) is not statistically significant ($F_{1,15} = 0.2$, $P = 0.66$), whereas the trend since 2015 (680 fewer fish per year on average) is strongly and significantly negative ($F_{1,9} = 808$, $P = 0.016$). The dome-shaped long term trend may be the result of management measures to control harvest (as evidenced by the increasing trend in Chinook salmon release numbers), or the result of declining stocks.

Pacific Halibut harvest did not show a strong among-year trend ($F_{1,13} = 0.4$, $P = 0.55$, Table 29, Figure 17). While fishing effort increased over the 31-year period, the total harvest of Pacific Halibut in recent years (2009, 2015-2025) has been very similar to that from 2001-2002; Table 29). This is very likely the result of effective management measures that were enacted between 2008 and 2009 (including the reduction in the daily bag limit from two to one), that appeared to have reduced the catch of halibut by nearly 50% (Figure 17).

The Lingcod harvest (June to August) in 2025 is the highest observed for the available time series, and that for each of the last five years (2021-2025) are the five highest observed (Figure 17). In general, the June to August Lingcod harvest from 2018 to 2025 (4,064 to 9,302) has been higher than in other recent years (e.g., 1,803 to 3,052 in 2015-2017). The long-term trend is statically significant (ln scale, $F_{1,14} = 11.4$, $P = 0.005$; Figure 17). This, despite a coast-wide reduction in the Lingcod bag limit from three to two fish starting in May 2018. The elevated and harvest values from 2021 and 2025 could potentially cause concern.

The meta-analysis results make three important points: 1) the effort and CPUE of certain species has increased tremendously over the past 30 years (even despite the COVID-19 pandemic), potentially imperilling stocks if left unchecked; 2) management decisions for Pacific Halibut and possibly Chinook Salmon appear to have been effective, indicating that tools are available to keep effort and CPUE increases from going unchecked; and 3) despite that, Lingcod harvest is increasing to a concerning degree.

May Effort

After the 2018 study year, we made the case that substantial angling activity was being missed by starting the creel survey in June each year (Robichaud and Addison 2019). Surveys in 1995 and 1996 (Van Tongeren and Winther 2010) included May, which accounted for 14% and 5% of the May-August effort, and for 8% and 1% of the May-August Chinook harvest. May surveys occurred again in 2001, 2002, and 2008, representing 5%, 10%, and 8% of the May-August effort, respectively (and 5%, 7%, and 4% of the May-August Chinook harvest). Beginning in 2019, we did start the survey in May, and found that May effort (1,163 to 1,851 boat trips) accounted for 8-15% of the May-August total, and in most years (2019 to 2021 and in 2023 and 2025) accounted for 15-21% of the May-August total Chinook Salmon harvest (May fishing in 2022 and 2024 accounted for 32% and 28% of the Chinook Salmon harvest, respectively).




Figure 17. Harvest of Chinook and Coho salmon, and Pacific Halibut from 1995 to 2025, and of Lingcod from 2008 to 2025. Annual estimates only include data from June, July and August. Only years with data from all three months are included. Long-term linear regression equations are shown in black within each panel, and the Chinook Salmon harvest trend since 2015 is in red. For Pacific Halibut, bag limits changed after 2008 from two fish per day (orange dots) to one fish per day (blue dots). Trendline for Lingcod was calculated on a logarithmic scale.




Figure 18. Number of Chinook Salmon released from June to August by anglers in Areas 3 and 4 from 2009 to 2025. Only years with data from all three months are included. Long-term linear regression equation (calculated on a logarithmic scale) is shown in the bottom right.

Table 28. Monthly recreational fishing effort (boat trips) in Area 3 and 4 from 1995 to 2025, as estimated using creel survey methods.

Fishing Effort

	Year																			
Month	1995	1996	1998	2000	2001	2002	2006	2008	2009	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
May	965	391	--	--	449	960	--	983	--	--	--	--	--	1,553	1,320	1,785	1,587	1,428	1,851	1,163
June	1,590	2,490	--	--	3,801	3,463	--	4,696	3,043	3,424	3,961	4,574	2,873	3,790	1,970	2,507	2,712	2,735	3,335	3,150
July	1,394	2,385	1,307	1,842	2,588	2,892	--	3,750	3,804	4,279	5,295	6,044	4,593	4,644	3,745	4,318	4,040	3,868	4,247	4,726
Aug	1,927	1,400	1,351	1,327	1,808	1,843	2,593	2,529	2,865	3,881	3,325	2,980	3,304	4,175	2,631	3,642	3,285	4,055	4,296	5,148
Sept	864	488	504	954	540	676	758	394	817	--	--	--	--	--	--	--	--	--	--	--
TOTAL	6,740	7,154	3,162	4,123	9,186	9,834	3,351	12,352	10,529	11,585	12,581	13,598	10,770	14,163	9,665	12,252	11,624	12,086	13,729	14,187

Table 29. Monthly Chinook Salmon, Coho Salmon, Pacific Halibut, and Lingcod harvest, and Chinook Salmon releases, in the Area 3 and 4 recreational fishery from 1995 to 2025, as estimated using creel survey methods.

Month	Year																						
	1995	1996	1998	2000	2001	2002	2006	2008	2009	2010	2011	2012	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025
Chinook Salmon Harvest																							
May	141	45	--	--	509	522	--	474	--	--	--	--	--	--	--	--	2,234	1,256	1,580	1,428	1,191	2,594	1,231
June	907	2,165	--	--	6,630	4,028	--	7,708	4,797	--	--	--	7,071	5,678	5,229	1,273	5,675	2,574	1,834	1,564	3,180	2,848	1,843
July	505	1,097	254	1,611	3,260	2,822	--	3,080	3,873	--	--	--	4,016	3,474	3,800	2,964	4,806	2,856	2,797	9	1,155	1,272	1,176
Aug	245	257	35	108	299	474	1,207	688	503	--	--	--	1,669	891	1,078	1,593	2,437	1,560	1,172	1,431	917	2,525	3,029
Sept	15	62	6	93	61	59	203	20	4	--	--	--	--	--	--	--	--	--	--	--	--	--	--
TOTAL	1,813	3,626	295	1,812	10,759	7,905	1,410	11,970	9,177				12,756	10,043	10,107	5,830	15,152	8,247	7,383	4,432	6,443	9,239	7,279
Chinook Salmon Release																							
May	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	1,145	933	497	536	980	1,357	1,005
June	--	--	--	--	--	--	--	--	808	--	--	--	1,534	975	2,800	2,121	4,735	2,088	3,652	2,297	3,790	4,303	2,931
July	--	--	--	--	--	--	--	--	705	--	--	--	594	710	1,811	2,612	2,382	1,560	3,694	6,213	3,415	3,433	5,186
Aug	--	--	--	--	--	--	--	--	175	--	--	--	103	506	697	1,264	2,866	1,061	1,799	966	627	1,511	2,343
Sept	--	--	--	--	--	--	--	--	15	--	--	--	--	--	--	--	--	--	--	--	--	--	--
TOTAL									1,703				2,231	2,190	5,308	5,997	11,129	5,642	9,643	10,012	8,812	10,605	11,466
Coho Salmon Harvest																							
May	0	0	--	--	0	0	--	0	--	--	--	--	--	--	--	--	29	0	0	3	2	27	39
June	335	474	--	--	1,771	1,332	--	3,913	3,686	--	--	--	4,436	4,389	1,839	126	1,475	1,006	3,038	1,356	2,555	2,808	3,787
July	402	3,565	1,937	568	4,736	9,798	--	15,982	17,043	--	--	--	15,101	21,761	22,871	5,758	10,341	16,304	17,685	13,929	17,360	15,643	17,529
Aug	1,907	3,771	1,500	2,125	6,488	8,292	15,647	13,791	16,459	--	--	--	12,612	11,200	14,004	4,580	11,974	8,288	17,403	16,331	22,608	10,775	16,727
Sept	1,115	1,417	79	790	1,006	1,010	3,215	733	3,136	--	--	--	--	--	--	--	--	--	--	--	--	--	--
TOTAL	3,759	9,227	3,516	3,483	14,001	20,432	18,862	34,419	40,324				32,148	37,350	38,714	10,464	23,820	25,599	38,126	31,620	42,525	29,253	38,082
Pacific Halibut Harvest																							
May	--	--	--	--	281	467	--	1,435	--	--	--	--	--	--	--	--	941	867	1,499	1,385	934	1,795	755
June	--	--	--	--	4,872	3,912	--	8,367	3,029	--	--	--	3,551	4,032	4,632	3,407	2,698	2,344	3,725	4,229	2,636	3,375	2,681
July	--	--	--	3,963	3,535	4,716	--	8,507	4,466	--	--	--	4,096	6,069	6,563	4,263	3,894	4,150	6,518	5,558	4,887	4,385	4,330
Aug	--	--	1,801	2,530	3,085	3,247	8,178	5,865	4,142	--	--	--	5,201	3,670	3,410	3,183	3,435	3,504	4,929	3,953	5,049	5,402	4,157
Sept	--	--	328	1,224	222	615	1,812	774	1,205	--	--	--	--	--	--	--	--	--	--	--	--	--	--
TOTAL			2,129	7,717	11,995	12,957	9,990	24,948	12,842				12,848	13,771	14,605	10,853	10,967	10,865	16,671	15,125	13,505	14,957	11,923
Lingcod Harvest																							
May	--	--	--	--	--	--	--	81	--	--	--	--	--	--	--	--	219	635	1,178	1,661	802	1,477	792
June	--	--	--	--	--	--	--	1,462	760	1,093	2,241	692	1,042	732	1,124	1,754	1,201	964	2,120	2,722	2,347	1,916	2,223
July	--	--	--	--	--	--	--	1,348	1,019	1,536	1,180	1,342	951	701	1,173	1,740	1,389	2,157	2,312	2,972	2,987	3,166	3,572
Aug	--	--	--	--	--	--	--	884	999	1,521	1,528	274	1,059	370	475	776	1,704	943	1,733	1,792	2,747	2,174	3,508
Sept	--	--	--	--	--	--	--	167	208	--	--	--	--	--	--	--	--	--	--	--	--	--	--
TOTAL								3,942	2,986	4,150	4,949	2,308	3,052	1,803	2,772	4,270	4,514	4,699	7,343	9,147	8,884	8,733	10,094

Much of the early season angling targets Chinook Salmon, and occurs during a period in which the Area 3 and 4 stock composition is unknown or poorly resolved. As such, the Department of Fisheries and Oceans Canada (DFO) expressed interest in a May creel survey because genetic samples collected from Chinook Salmon would help them resolve stock composition and help determine exploitation rates on early-run-timing Chinook Salmon stocks. In 2019, during the month of May, 269 Chinook Salmon were biosampled, including 230 scale samples, and 22 tagged heads. Since then, 129-194 Chinook Salmon scale samples have been collected each May (129 in 2025), representing 29-52% of the annual total (29% in 2025). Since 2020, May accounted for 11-23 of the Chinook Salmon heads submitted to the Salmon Head Recovery Program (23 in 2025), 26-56 % of the season-wide totals (27% in 2025).

April Effort

Trailer counts made during April of 2019-2020, and 2023-2025 allowed the estimation of the amount of non-lodge angling activity that is being missed by starting the creel survey in May each year. Over these five years, April effort may have varied from 766 to 1,303 trips, which represents an additional 8-11% of angling effort in addition to effort that is documented from May to August. It is not known what species these early season anglers are targeting.

Precision of the Results

Overall, our level of survey effort, both for CPUE and Effort estimation, were adequate, as evidenced by the relatively good precision of the harvest estimates for most of the taxa surveyed. In 2025 our precision goal was to estimate total harvests within 15% of the true value 19 times out of 20, for the main taxa (Chinook Salmon, Coho Salmon, and Pacific Halibut), for which harvest estimates were precise to within 3-4%. Pink Salmon, Lingcod, and rockfish estimates were precise to within 8%, 6%, and 6%, respectively. Chum and Sockeye salmon estimates, and those for other groundfish, were not precise to within 15%, though the survey was not designed to meet any level of precision for these last taxa.

The precision of our harvest estimates depended largely on the variability in the CPUE and effort data. Differing levels of angler expertise, and day-to-day changes in abundance of the target species, fishing effort, weather, or fishing conditions mean that fishing trips have a wide range of outcomes. Similarly, day-to-day differences in weather or other factors (such as leisure time, day of week, the price of gasoline, etc.) affect the number of anglers that decide to go out on the water on any given day, resulting in variability within the boat-count dataset. Sampling error is another main source of estimation imprecision; and by meeting our precision goals we have demonstrated that we are sampling an adequate proportion of the population.

Accuracy of the Results

The accuracy of the results depends on the quality of data provided by the anglers to the interviewers. This is partly why we have a target to sample 20% of estimated effort. Even so, each time an angler is relied upon for unconfirmed data, there is opportunity for results to become inaccurate. For example, in this study, 1,493 of the 2,043 Chinook Salmon (73%) and 5,989 of the 9,205 Coho Salmon (65%) reported as kept during angler interviews were recorded as observed by the interviewer, and 448 and 530 (22% and 6%) were handled for biosampling.

While not every fish can be observed², each one that goes unconfirmed is nevertheless a possible source of inaccuracy in the reported harvest totals. Angler compliance with requests to examine their catch is voluntary.

In this study, the angler effort estimates depend on the accuracy of the aerial survey boat counts and the activity patterns for each month. For example, we have no estimate of the proportion of angling boats that are missed during overflights, where missed boats would contribute to the underestimation of recreational fishing effort. Also, we do not know if all fishing areas are adequately covered by the current flightpath – perhaps there are fishing locations that are hidden behind islands (e.g., Dunira Island or the Moffat Islands), too far from the flightpath for the surveyors to see active fishing boats. On the other hand, our estimate of fishing effort may be overestimated due to our inability to survey and count boats on foggy or bad weather days. We know that during bad weather, fewer anglers go out on the water; yet these same conditions can cause flights to be cancelled, and thus the lowest effort values are sometimes not sampled. In 2026, we will begin recording weather (fog, wind speed) along with each trailer count to determine how many fewer trailers are present on bad weather days.

Another potential source of inaccuracy is the removal of lodge boats from the overflight counts. There is uncertainty around whether all of the active lodge boats were observed by the overflights (e.g., vessels outside of the survey area), as evidenced by surveys for which more lodge boats were operating than there were total boats observed in Area 3 during the flight. There is also uncertainty around the proportion of lodge vessels that were active during the overflights. These factors may influence the accuracy of the estimates of non-lodge fishing effort.

RECOMMENDATIONS

- The catch and effort estimates for the Area 3 and 4 recreational creel surveys for 2010-2014 should be compiled and included in summary tables similar to Table 28 and 29 in this report. If sufficient funding can be obtained, the results from 2010-14 surveys should be compiled into a manuscript report. This additional information will help provide context for the current results, and help clarify how rates of effort and harvest have changed in recent years.
- We recommend that future surveys continue to run for four months (May to August) rather than three (June to August). May surveys will help document and quantify angling effort and harvest during that month. The Department of Fisheries and Oceans Canada (DFO) has expressed interest in a May creel survey because genetic samples collected from Chinook Salmon will help resolve the currently unknown early-season Area 3 and 4 stock composition, which will help determine exploitation rates on early-run-timing Chinook Salmon stocks.

² Fish may not be observed when anglers refuse to provide access to their fish storage containers for various reasons. Reluctance to provide access may be due to fish already being processed and packed in ice, or boat launch traffic congestion causing tension, among other reasons.

- Trailer counts should be conducted in April 2026 for comparison with April trailer counts in 2019-2025; and should be continued throughout the creel survey season as in recent years. For every day, observers should record whether the weather was bad enough to cancel a flight, had there been one booked for that day.
- Factors that are resulting increased Lingcod harvest should be investigated to inform managers should additional measures be enacted.

ACKNOWLEDGEMENTS

We would like to thank the creel program manager, Anke de Wit, and surveyors (Arianna Bonafine, Kassidy Main-Cooper, Isabelle Pebernat, and Rory Scott) for collecting large amounts of data and bio-samples from anglers. Thanks to Pacific Salmon Commission, who provided support for this work through the Northern Fund. Thanks to Chelsea May, Ryan Whitmore, Jessica Otley, Shaun Davies, and Darren Chow for acting as liaison between our team and DFO. Thanks also go to the administrative staff at LGL and NCSFNSS, whose work made it possible for this study to be conducted. We also would like to thank Karl English for providing useful comments and edits on previous drafts of this report. Last but not least, we thank the many anglers, charter operators, and fishing lodges who participated in the survey.

LITERATURE CITED

- Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B* 57: 289-300.
- Clark, W. G. 1992. Validation of the IPHC length-weight relationship for halibut. *International Pacific Halibut Commission Report of Assessment and Research Activities* 1991: 113-116.
- English, K. K., G. F. Searing, and D. A. Nagtegaal. 2002. Review of Georgia Strait recreational creel survey, 1983-1999. *Canadian Technical Report of Fisheries and Aquatic Sciences* 2414: 81 p.
- Goodman, L. 1960. On the exact variance of products. *Journal of the American Statistical Association* 55: 708-713.
- Robichaud, D., and A. Addison. 2018. North Coast (Areas 3 & 4) recreational angling creel survey 2017. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2019. North Coast (Areas 3 & 4) recreational angling creel survey 2018. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2020. North Coast (Areas 3 & 4) recreational angling creel survey 2019. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.

- Robichaud, D., and A. Addison. 2021. North Coast (Areas 3 & 4) recreational angling creel survey 2020. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2022. North Coast (Areas 3 & 4) recreational angling creel survey 2021. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2023. North Coast (Areas 3 & 4) recreational angling creel survey 2022. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2024. North Coast (Areas 3 & 4) recreational angling creel survey 2023. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., and A. Addison. 2025. North Coast (Areas 3 & 4) recreational angling creel survey 2024. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., A. Addison, and M. Imrie. 2016. North Coast (Areas 3 & 4) recreational angling creel survey 2015. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Robichaud, D., A. Addison, and M. Imrie. 2017. North Coast (Areas 3 & 4) recreational angling creel survey 2016. Report for North Coast - Skeena First Nations Stewardship Society, Prince Rupert, BC.
- Sokal, R.R. and F.J. Rohlf. 1995. Biometry. W. H. Freeman and Company, New York.
- Van Tongeren, V., and I. Winther. 2010. 2009 North Coast (Areas 3 & 4) creel survey statistics for salmon and groundfish. *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 2907.
- Zar, J. H. 1984. Biostatistical Analysis. Prentice-Hall Inc., Englewood Cliffs, NJ.

APPENDICES

APPENDIX 1: DATA FORMS

SHIFT SUMMARY SHEET

Landing Site: _____

Interviewer Name: _____

Interviewer Code: _____

Date: _____

Shift Code: _____

Shift Times: _____

TIME BLOCKS		Fishing Conditions	Boats Missed	Not Fishing	Fishing
From - To	TB				
Before 7:00	1				
7:00-7:59	2				
8:00-8:59	3				
9:00-9:59	4				
10:00-10:59	5				
11:00-11:59	6				
12:00-12:59	7				
13:00-13:59	8				
14:00-14:59	9				
15:00-15:59	10				
16:00-16:59	11				
17:00-17:59	12				
18:00-18:59	13				
19:00-19:59	14				
20:00-20:59	15				
21:00-21:59	16				
TOTALS:					

Comments: _____

Trailer Count: _____ Time: _____

Database Tally #:	(office use only)
-------------------	-------------------

**2025 AREA 3/4 CREEL SURVEY
AERIAL SURVEY VESSEL ENUMERATION FORM**

DATE: _____

FLIGHT#: _____

SURVEYOR: _____

Note: only survey shaded subareas when conditions permit

Sub Area	RIGHT FRONT COUNT				TOTAL COUNT				FLIGHT TIME		SEA STATE (1-4)
	SPORT (Fishing)	SPORT (Moving)	LARGE (Fishing)	LARGE (Moving)	S (F)	S (M)	C (F)	C (M)	START	FINISH	
A											
C											
B											
C											
D											
E											
C											
F											
E											
F											
G											
L											
G											
J											
I											
K											
I											
H											
C											
A											

WEATHER / SEA STATE / COMMENTS: _____

SIGNATURE: _____

[illegible]

APPENDIX 2: HPUE ESTIMATES (AVERAGE NUMBER OF FISH HARVESTED PER NON-LODGE BOAT TRIP) BY PERIOD, TRIP TYPE, SUBAREA AND SPECIES.

Average number of salmon harvested per boat trip, 1-28 May 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Chinook Salmon												
Guided	n/a	n/a	n/a	0.05	3.14	2.23	2.30	1.79	1.64	0.60	0.71	0.50
Unguided	0.29	0.29	0.44	0.07	1.00	0.85	1.09	1.61	2.06			
Combined	0.65	0.65	0.82	0.06	1.29	1.29	1.47	1.68	1.85	0.38	0.42	0.50
Chum Salmon												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Coho Salmon												
Guided	n/a	n/a	n/a				0.10	0.02				
Unguided												
Combined							0.03	0.01				
Pink Salmon												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Sockeye Salmon												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Unknown Salmon												
Guided	n/a	n/a	n/a									
Unguided												
Combined												

Average number of salmon harvested per boat trip, 29 May to 22 June 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Chinook Salmon												
Guided	1.08	1.68	1.41		1.24	1.26	1.87	2.10	2.22	0.60	0.54	1.25
Unguided	1.44	0.89	0.44	0.02	0.53	0.48	0.78		1.38	0.33	0.53	0.40
Combined	1.24	1.22	0.58	0.01	0.69	0.64	1.17	0.25	1.75	0.47	0.54	0.91
Chum Salmon												
Guided												
Unguided												
Combined												
Coho Salmon												
Guided	0.25	1.05	0.68		0.24	0.09	0.10	0.16	0.18	0.27		0.50
Unguided	0.33	0.19	0.44			0.02	0.02		0.10			
Combined	0.29	0.54	0.33		0.05	0.04	0.05		0.13	0.13		0.36
Pink Salmon												
Guided												
Unguided						0.02			0.01			
Combined						0.02			0.01			
Sockeye Salmon												
Guided												
Unguided												
Combined												
Unknown Salmon												
Guided												
Unguided												
Combined												

Average number of salmon harvested per boat trip, 23 June to 23 July 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Chinook Salmon												
Guided												
Unguided					0.05	0.01	0.02					
Combined					0.04	0.01	0.01					
Chum Salmon												
Guided												
Unguided												
Combined												
Coho Salmon												
Guided	8.23	3.23	5.33		3.16	9.26	2.71	4.34	4.58	4.03	4.28	5.71
Unguided	3.64	3.44	0.42		0.71	3.88	2.60	2.86	3.35	1.16	3.15	2.23
Combined	5.98	3.32	2.00		1.11	5.90	2.63	3.50	3.99	2.35	3.81	5.00
Pink Salmon												
Guided	0.73	0.69	2.00		0.15	0.45	0.29	0.37	0.34	0.23	0.19	
Unguided	0.20	0.33	0.16			0.47	0.65	0.86	0.49		0.11	0.06
Combined	0.47	0.55	0.75			0.46	0.57	1.20	0.41	0.09	0.16	
Sockeye Salmon												
Guided												
Unguided									0.04			
Combined									0.02			
Unknown Salmon												
Guided												
Unguided								0.14		0.05		0.02
Combined								0.10		0.03		

Average number of salmon harvested per boat trip, 24 July to 17 August 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Chinook Salmon												
Guided	0.54	1.40	1.05		0.60	0.64	1.70	1.45	1.45	0.56	0.53	0.77
Unguided	0.60	0.83	0.49		0.23	0.38	0.54	0.18	0.60	0.45	0.37	0.33
Combined	0.56	1.16	0.66		0.29	0.45	0.71	0.13	0.91	0.50	0.45	0.63
Chum Salmon												
Guided												
Unguided									0.01			
Combined									0.00			
Coho Salmon												
Guided	7.92	5.87	10.00		4.80	3.14	5.48	5.42	5.61	2.38	3.53	2.69
Unguided	5.07	4.71	4.26		1.60	1.69	2.82	1.09	4.05	1.85	2.10	1.50
Combined	7.13	5.38	6.02		2.12	2.09	3.21	0.80	4.62	2.08	2.84	2.32
Pink Salmon												
Guided	1.15	0.13	0.42		0.27	0.56	0.35	0.43	0.46	0.13	0.25	
Unguided	0.73	0.44	0.79		0.15	0.24	0.26		0.51	0.05	0.27	0.50
Combined	1.04	0.26	0.68		0.17	0.33	0.28		0.49	0.08	0.26	0.16
Sockeye Salmon												
Guided	0.05	0.02	0.05					0.01	0.01		0.03	
Unguided							0.01					
Combined	0.04	0.01	0.02				0.01		0.00		0.02	
Unknown Salmon												
Guided								0.01	0.01			1.54
Unguided							0.02		0.02			
Combined							0.01		0.02			1.05

Average number of salmon harvested per boat trip, 18-31 August 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Chinook Salmon												
Guided	0.66	0.44	0.88		0.60	0.24	2.30	2.30	2.56	1.91	0.25	0.20
Unguided	0.25	0.90	0.21			0.29	0.14	0.46	0.67	0.13	0.12	0.11
Combined	0.55	0.62	0.52		0.09	0.28	0.13	0.96	1.35	0.88	0.19	0.15
Chum Salmon												
Guided												
Unguided												
Combined												
Coho Salmon												
Guided	5.56	0.94	6.79		0.80	0.71	0.73	0.73	0.81	0.27	1.90	1.50
Unguided	3.17	3.50	5.04		0.07	0.31	1.00	0.77	0.69	0.80	0.29	0.49
Combined	4.91	1.92	5.85		0.18	0.40	0.90	0.76	0.73	0.58	1.16	1.15
Pink Salmon												
Guided	0.59	0.13	0.54				0.13	0.13	0.07			0.10
Unguided			0.14		0.03	0.02	0.14	0.09	0.06	0.13	0.24	0.17
Combined	0.43	0.08	0.33		0.03	0.01	0.19	0.10	0.07	0.08	0.11	0.08
Sockeye Salmon												
Guided												
Unguided											0.06	0.03
Combined											0.03	
Unknown Salmon												
Guided												
Unguided												
Combined												

Average number of groundfish harvested per boat trip, 1-28 May 2025, by trip type and subarea. Zeros hidden.

		Subarea										
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Arrow-tooth Flounder												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Big Skate												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Cabezon												
Guided	n/a	n/a	n/a					0.02	0.03			
Unguided												
Combined								0.01	0.01			
English Sole												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Greenlings												
Guided	n/a	n/a	n/a									
Unguided												
Combined	0.18	0.18	0.27									
Lingcod												
Guided	n/a	n/a	n/a		1.86	0.55		1.53	2.00	1.87		3.50
Unguided	0.07	0.07			0.02		0.05	0.74	1.11	0.89		0.89
Combined	0.88	0.88	0.18		0.27	0.17	0.03	1.07	1.54	1.50		3.50
Longnose Skate												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Pacific Cod												
Guided	n/a	n/a	n/a		0.29	0.05		0.33	0.42	2.47	5.29	
Unguided	0.14	0.14						0.02				
Combined	0.12	0.12			0.04	0.01		0.14	0.21	1.54	3.08	
Pacific Halibut												
Guided	n/a	n/a	n/a	0.05		0.55	0.40	1.63	2.00	2.40	2.71	2.13
Unguided	0.86	0.86			0.05	0.43	0.41	0.49	0.60	0.44	0.80	0.44
Combined	0.94	0.94		0.02	0.04	0.46	0.41	0.96	1.28	1.67	1.92	2.13
Pacific Rock Sole												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Pacific Spiny Dogfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Sablefish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Spotted Ratfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Starry Flounder												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Unknown Groundfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												

Average number of groundfish harvested per boat trip, 29 May to 22 June 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Arrow-tooth Flounder												
Guided												
Unguided												
Combined												
Big Skate												
Guided								0.01	0.01			
Unguided												
Combined									0.01			
Cabezon												
Guided												
Unguided							0.02					0.03
Combined							0.01					0.09
English Sole												
Guided												
Unguided												
Combined												
Greenlings												
Guided								0.01	0.01			
Unguided							0.04					
Combined							0.02		0.01			
Lingcod												
Guided	0.17	3.89	2.41		0.53	0.26	0.39	0.89	1.10	1.67	0.29	2.50
Unguided	0.11	0.41			0.07	0.12	0.07	1.83	0.85	0.27	0.41	0.51
Combined	0.14	1.85	0.50		0.17	0.15	0.19	1.38	0.96	0.97	0.34	2.45
Longnose Skate												
Guided												
Unguided												
Combined												
Pacific Cod												
Guided	0.92		0.32				0.06	0.07	0.07	3.47	5.88	0.50
Unguided					0.02	0.04	0.07		0.08		0.24	0.11
Combined	0.52				0.01	0.03	0.07		0.08	1.73	3.54	0.36
Pacific Halibut												
Guided	2.25	1.37	1.74		0.24	0.30	1.16	1.58	1.78	1.53	2.54	1.75
Unguided	0.78	0.56	0.22		0.26	0.62	0.40	1.33	1.01	0.67	0.94	1.03
Combined	1.62	0.89	0.67		0.25	0.55	0.67	1.00	1.35	1.10	1.88	2.18
Pacific Rock Sole												
Guided												
Unguided												
Combined												
Pacific Spiny Dogfish												
Guided								0.01	0.01			
Unguided												
Combined									0.01			
Sablefish												
Guided												
Unguided												
Combined												
Spotted Ratfish												
Guided												
Unguided												
Combined												
Starry Flounder												
Guided												
Unguided												
Combined												
Unknown Groundfish												
Guided												
Unguided									0.02			
Combined									0.01			

Average number of groundfish harvested per boat trip, 23 June to 23 July 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Arrow-tooth Flounder												
Guided												
Unguided						0.01			0.01			
Combined						0.00			0.00			
Big Skate												
Guided												
Unguided												
Combined												
Cabezon												
Guided								0.01	0.01			
Unguided		0.04										
Combined		0.02							0.00			
English Sole												
Guided												
Unguided												
Combined												
Greenlings												
Guided					0.00	0.01		0.07	0.08			
Unguided		0.11	0.32				0.02		0.06	0.02		0.01
Combined		0.05	0.21			0.00	0.01		0.07	0.01		
Lingcod												
Guided	0.31	3.28	0.11		0.19	0.54	0.59	1.14	1.15	1.97	0.32	1.43
Unguided	0.04	1.00	0.32		0.07	0.17	0.31	0.86	0.61	0.80	0.37	0.56
Combined	0.18	2.35	0.25		0.15	0.31	0.38	1.80	0.89	1.28	0.34	1.25
Longnose Skate												
Guided												
Unguided												
Combined												
Pacific Cod												
Guided	0.77	0.13			0.07	0.20	0.06	0.09	0.10	1.45	3.91	1.86
Unguided	0.08		0.05		0.14	0.05	0.49		0.09	0.27	0.72	0.52
Combined	0.43	0.08	0.04		0.13	0.11	0.39		0.10	0.76	2.57	1.63
Pacific Halibut												
Guided	0.85	1.36	0.56		0.27	0.76	0.82	1.76	1.88	1.35	1.75	2.57
Unguided	0.80	0.48	1.16	0.01	0.69	0.59	0.73	1.57	0.99	1.11	0.81	0.96
Combined	0.82	1.00	0.96	0.00	0.72	0.65	0.75	1.90	1.45	1.21	1.36	2.50
Pacific Rock Sole												
Guided		0.03										0.14
Unguided						0.01						
Combined		0.02				0.00						0.13
Pacific Spiny Dogfish												
Guided												
Unguided												
Combined												
Sablefish												
Guided			0.22									
Unguided												
Combined			0.07									
Spotted Ratfish												
Guided												
Unguided												
Combined												
Starry Flounder												
Guided												
Unguided												
Combined												
Unknown Groundfish												
Guided		0.03										
Unguided									0.01			
Combined		0.02							0.00			

Average number of groundfish harvested per boat trip, 24 July to 17 August 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Arrow-tooth Flounder												
Guided								0.01	0.02			
Unguided					0.01							
Combined					0.01				0.01			
Big Skate												
Guided												
Unguided												
Combined												
Cabezon												
Guided		0.02						0.01	0.01			
Unguided									0.01			
Combined		0.01							0.01			
English Sole												
Guided											0.03	
Unguided												
Combined											0.02	
Greenlings												
Guided		0.02	0.05				0.04	0.01	0.01			
Unguided					0.01	0.01	0.08					
Combined		0.01	0.02		0.01	0.01	0.08		0.00			
Lingcod												
Guided	0.51	2.02	1.26		1.33	0.16	0.30	1.04	1.12	0.19	1.16	3.62
Unguided		0.90	0.07			0.12	0.24	1.64	0.49	1.35	0.23	2.17
Combined	0.37	1.54	0.44		0.22	0.13	0.25	2.00	0.72	0.83	0.71	3.16
Longnose Skate												
Guided												
Unguided												
Combined												
Pacific Cod												
Guided	0.67	0.09	0.11			0.25		0.42	0.51	0.94	1.47	0.85
Unguided	0.20	0.37			0.23	0.10	0.02		0.09		0.40	
Combined	0.54	0.21	0.03		0.19	0.14	0.01	0.07	0.25	0.42	0.95	0.58
Pacific Halibut												
Guided	0.90	1.84	0.63		0.53	1.20	0.96	1.35	1.37	0.44	1.91	2.38
Unguided	1.33	0.93	0.70		0.38	0.51	0.42	1.00	0.53	0.65	1.40	1.83
Combined	1.02	1.45	0.68		0.41	0.70	0.50	1.60	0.84	0.56	1.66	2.21
Pacific Rock Sole												
Guided												
Unguided												
Combined												
Pacific Spiny Dogfish												
Guided												
Unguided				0.01								
Combined				0.00								
Sablefish												
Guided			0.05									
Unguided												
Combined			0.02									
Spotted Ratfish												
Guided												
Unguided												
Combined												
Starry Flounder												
Guided												
Unguided												
Combined												
Unknown Groundfish												
Guided						0.02						
Unguided												
Combined						0.00						

Average number of groundfish harvested per boat trip, 18-31 August 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Arrow-tooth Flounder												
Guided												
Unguided												
Combined												
Big Skate												
Guided												
Unguided												
Combined												
Cabezon												
Guided		0.13					0.03	0.03	0.04			
Unguided												
Combined		0.08						0.01	0.01			
English Sole												
Guided												
Unguided												
Combined												
Greenlings												
Guided		0.13	0.04				0.13	0.13	0.15			
Unguided												
Combined		0.08	0.02					0.04	0.05			
Lingcod												
Guided	0.34	3.38			4.00	0.41	2.13	2.13	1.81	2.36	0.70	2.30
Unguided		0.60	0.07	0.12		0.12	0.32	0.67	0.92	1.87	0.47	1.29
Combined	0.25	2.31	0.04	0.05	0.59	0.19	0.77	1.07	1.24	2.08	0.59	2.46
Longnose Skate												
Guided												
Unguided												
Combined												
Pacific Cod												
Guided	0.38					0.47	0.73	0.73	0.59		0.80	2.90
Unguided	1.33		0.11			0.07	0.04	0.19	0.29	0.13	0.47	0.43
Combined	0.64		0.06			0.16	0.23	0.34	0.40	0.08	0.65	2.62
Pacific Halibut												
Guided	0.69	2.81	0.04		1.40	1.29	2.20	2.20	2.15	0.36	1.95	2.30
Unguided	0.25	0.80	0.39		0.48	0.98	0.79	0.96	1.10	0.80	1.18	1.06
Combined	0.57	2.04	0.23		0.62	1.05	0.97	1.30	1.48	0.62	1.59	2.15
Pacific Rock Sole												
Guided					0.20							
Unguided												
Combined					0.03							
Pacific Spiny Dogfish												
Guided												
Unguided								0.03	0.04			
Combined								0.02	0.03			
Sablefish												
Guided												
Unguided												
Combined												
Spotted Ratfish												
Guided												
Unguided												
Combined												
Starry Flounder												
Guided												
Unguided												
Combined												
Unknown Groundfish												
Guided												
Unguided												
Combined												

Average number of rockfish harvested per boat trip, 1-28 May 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Black Rockfish												
Guided	n/a	n/a	n/a		1.43	0.27	0.50	0.77	0.85	0.87		1.63
Unguided	0.07	0.07	0.11			0.02	0.05	0.25	0.34			
Combined	0.12	0.12	0.18		0.20	0.10	0.19	0.46	0.59	0.54		1.63
Boccacio Rockfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Canary Rockfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
China Rockfish												
Guided	n/a	n/a	n/a			0.09		0.02	0.03	0.07		0.13
Unguided								0.03	0.03			
Combined						0.03		0.03	0.03	0.04		0.13
Copper Rockfish												
Guided	n/a	n/a	n/a		0.29							
Unguided	0.07	0.07				0.06		0.02	0.03			
Combined	0.06	0.06			0.04	0.04		0.01	0.01			
Dusky Rockfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Quillback Rockfish												
Guided	n/a	n/a	n/a					0.40	0.52	0.13		0.25
Unguided					0.05	0.19		0.10	0.14			
Combined					0.04	0.13		0.22	0.32	0.08		0.25
Redbanded Rockfish												
Guided	n/a	n/a	n/a					0.02	0.03			
Unguided								0.02	0.03			
Combined								0.02	0.03			
Redstripe Rockfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Tiger Rockfish												
Guided	n/a	n/a	n/a			0.05		0.09	0.12	0.07		0.13
Unguided								0.02	0.03			
Combined						0.01		0.05	0.07	0.04		0.13
Widow Rockfish												
Guided	n/a	n/a	n/a									
Unguided												
Combined												
Yelloweye Rockfish												
Guided	n/a	n/a	n/a							0.07		0.13
Unguided												
Combined										0.04		0.13
Yellowtail Rockfish												
Guided	n/a	n/a	n/a									
Unguided					0.05			0.08	0.03			
Combined					0.04			0.05	0.01			
Unknown rockfish												
Guided	n/a	n/a	n/a			0.41	0.10	0.02		1.13		2.13
Unguided						0.06	0.09	0.15	0.09			
Combined						0.17	0.09	0.10	0.04	0.71		2.13

Average number of rockfish harvested per boat trip, 29 May to 22 June 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Black Rockfish												
Guided	0.75	1.00	0.82		0.18		0.03	0.05	0.06	0.60		
Unguided		0.33			0.02	0.04	0.04	0.50	0.17	0.13	0.12	0.11
Combined	0.43	0.61			0.05	0.03	0.03	0.38	0.12	0.37	0.05	
Boccacio Rockfish												
Guided								0.02	0.02			
Unguided					0.02							
Combined					0.01				0.01			
Canary Rockfish												
Guided								0.01	0.01			
Unguided												
Combined									0.01			
China Rockfish												
Guided							0.03	0.02	0.01			
Unguided						0.04			0.02	0.07		0.03
Combined						0.03	0.01		0.02	0.03		
Copper Rockfish												
Guided	0.08	0.05	0.06					0.03	0.04		0.04	
Unguided					0.09	0.05	0.02				0.06	0.03
Combined	0.05	0.02			0.07	0.04	0.01		0.02		0.05	
Dusky Rockfish												
Guided		0.11	0.06					0.09	0.12		0.21	
Unguided							0.02		0.06			
Combined		0.04					0.01		0.09		0.12	
Quillback Rockfish												
Guided	0.25	0.74	0.50				0.06	0.24	0.32		0.38	
Unguided		0.15			0.07	0.05	0.02	0.50	0.11			
Combined	0.14	0.39			0.05	0.04	0.03	0.38	0.20		0.22	
Redbanded Rockfish												
Guided												
Unguided												
Combined												
Redstripe Rockfish												
Guided												
Unguided												
Combined												
Tiger Rockfish												
Guided		0.05	0.03				0.06	0.09	0.10	0.13		
Unguided									0.03			
Combined		0.02					0.02		0.06	0.07		
Widow Rockfish												
Guided												
Unguided		0.04					0.02					
Combined		0.02					0.01					
Yelloweye Rockfish												
Guided												
Unguided												
Combined												
Yellowtail Rockfish												
Guided	0.17		0.06					0.02	0.02		0.08	
Unguided						0.04			0.01			
Combined	0.10					0.03			0.02		0.05	
Unknown rockfish												
Guided		0.32	0.18		0.18		0.13	0.28	0.34		0.17	2.75
Unguided		0.07			0.21	0.19	0.11	0.50	0.17	0.20	0.71	0.43
Combined		0.17			0.20	0.15	0.12	0.38	0.25	0.10	0.39	2.00

Average number of rockfish harvested per boat trip, 23 June to 23 July 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Black Rockfish												
Guided	0.58	0.74			0.04	0.12	1.29	0.74	0.66	0.19	0.09	2.57
Unguided		0.26			0.12	0.03	0.04		0.25	0.14	0.02	0.07
Combined	0.29	0.55			0.13	0.06	0.33	0.10	0.46	0.16	0.06	2.25
Boccacio Rockfish												
Guided		0.03	0.11								0.01	
Unguided									0.03	0.02		0.01
Combined		0.02	0.04						0.01	0.01	0.01	
Canary Rockfish												
Guided		0.03	0.11							0.13		
Unguided						0.01			0.01			
Combined		0.02	0.04			0.01			0.00	0.05		
China Rockfish												
Guided		0.15			0.01	0.01		0.05	0.04	0.10		
Unguided		0.07			0.02	0.01			0.08	0.02		0.01
Combined		0.12			0.04	0.01		0.20	0.06	0.05		
Copper Rockfish												
Guided		0.10			0.02	0.05		0.04	0.05		0.05	
Unguided		0.11			0.10		0.02		0.04			
Combined		0.11			0.09	0.02	0.01		0.04		0.03	
Dusky Rockfish												
Guided								0.02	0.02			
Unguided						0.02			0.02		0.02	0.01
Combined						0.01			0.02		0.01	
Quillback Rockfish												
Guided	0.19	0.90	0.33		0.04	0.09	0.41	0.26	0.25	0.29	0.24	
Unguided	0.04	0.04			0.12	0.15	0.09		0.06	0.18	0.07	0.12
Combined	0.12	0.55	0.11		0.15	0.13	0.17		0.16	0.23	0.17	
Redbanded Rockfish												
Guided												
Unguided												
Combined												
Redstripe Rockfish												
Guided												
Unguided												
Combined												
Tiger Rockfish												
Guided		0.15					0.06	0.09	0.09	0.16		
Unguided							0.02		0.08	0.05		0.02
Combined		0.09					0.03		0.09	0.09		
Widow Rockfish												
Guided												
Unguided							0.02					
Combined							0.01					
Yelloweye Rockfish												
Guided												
Unguided												
Combined												
Yellowtail Rockfish												
Guided		0.15	0.11		0.01	0.02	0.18	0.06	0.05	0.13		
Unguided				0.03	0.07	0.02	0.04		0.01			
Combined		0.09	0.04	0.01	0.07	0.02	0.07		0.03	0.05		
Unknown rockfish												
Guided		0.15	0.11		0.03	0.09	0.18	0.16	0.17	0.03	0.15	1.00
Unguided	0.24	0.11	0.58			0.11	0.29		0.04	0.02		0.01
Combined	0.12	0.14	0.43		0.02	0.10	0.26		0.10	0.03	0.09	0.88

Average number of rockfish harvested per boat trip, 24 July to 17 August 2025, by trip type and subarea.
Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Black Rockfish												
Guided	0.13	1.29	0.37			0.06	0.17	0.40	0.46		0.44	0.85
Unguided	0.13	0.41			0.03	0.05	0.12		0.28	0.60		0.17
Combined	0.13	0.92	0.11		0.02	0.05	0.13		0.34	0.33	0.23	0.63
Boccacio Rockfish												
Guided	0.10					0.03					0.03	
Unguided						0.01						
Combined	0.07					0.01					0.02	
Canary Rockfish												
Guided		0.02										
Unguided									0.01			
Combined		0.01							0.00			
China Rockfish												
Guided	0.10						0.04	0.05	0.05			
Unguided					0.01	0.02	0.01		0.04			
Combined	0.07				0.01	0.01	0.01	0.07	0.04			
Copper Rockfish												
Guided	0.03	0.11						0.02	0.03			
Unguided						0.02	0.02		0.01		0.20	
Combined	0.02	0.06				0.02	0.02		0.02		0.10	
Dusky Rockfish												
Guided			0.05									
Unguided												
Combined			0.02									
Quillback Rockfish												
Guided	0.18	0.31	0.53		0.13	0.08	0.13	0.34	0.32	0.31	0.19	0.08
Unguided	0.07	0.07	0.05	0.01	0.12	0.09	0.15		0.13	0.35	0.10	0.33
Combined	0.15	0.21	0.19	0.00	0.12	0.09	0.15	0.53	0.20	0.33	0.15	0.16
Redbanded Rockfish												
Guided												
Unguided												
Combined												
Redstripe Rockfish												
Guided												
Unguided												
Combined												
Tiger Rockfish												
Guided		0.13	0.05					0.13	0.12	0.06		0.15
Unguided		0.12				0.01	0.02		0.01	0.05	0.03	
Combined		0.13	0.02			0.01	0.02	0.27	0.05	0.06	0.02	0.11
Widow Rockfish												
Guided												
Unguided							0.03					
Combined							0.03					
Yelloweye Rockfish												
Guided												
Unguided												
Combined												
Yellowtail Rockfish												
Guided	0.13	0.02	0.11			0.22	0.48	0.14	0.07			0.08
Unguided					0.13	0.04	0.04		0.02			
Combined	0.09	0.01	0.03		0.11	0.09	0.10		0.04			0.05
Unknown rockfish												
Guided	0.05	0.07				0.02		0.16	0.20		0.09	1.31
Unguided	0.07				0.03	0.03	0.12		0.06	0.30		
Combined	0.06	0.04			0.02	0.03	0.10		0.11	0.17	0.05	0.89

Average number of rockfish harvested per boat trip, 18-31 August 2025, by trip type and subarea. Zeros hidden.

	Subarea											
Species / Trip Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L
Black Rockfish												
Guided		0.06			0.20	0.18	0.37	0.37	0.41	0.82		2.40
Unguided		0.50			0.03	0.16	0.46	0.35	0.31	0.13	0.06	0.20
Combined		0.23			0.06	0.16	0.42	0.36	0.35	0.42	0.03	2.15
Boccacio Rockfish												
Guided		0.13										0.20
Unguided				0.02								
Combined		0.08		0.01								0.15
Canary Rockfish												
Guided							0.10	0.10	0.11			
Unguided						0.02		0.03	0.04			
Combined						0.01		0.05	0.07			
China Rockfish												
Guided		0.06					0.17	0.17	0.19		0.05	
Unguided								0.10	0.17		0.06	0.03
Combined		0.04						0.12	0.17		0.05	
Copper Rockfish												
Guided							0.13	0.13	0.15		0.30	0.30
Unguided						0.02		0.03	0.04			
Combined						0.01		0.06	0.08		0.16	0.23
Dusky Rockfish												
Guided			0.04				0.23	0.23	0.26			
Unguided		0.10										
Combined		0.04	0.02					0.06	0.09			
Quillback Rockfish												
Guided	0.03	0.56				0.06	0.77	0.77	0.70	0.73	0.20	0.40
Unguided		0.10	0.07		0.03	0.12	0.14	0.28	0.33		0.24	0.11
Combined	0.02	0.38	0.04		0.03	0.11	0.26	0.41	0.47	0.31	0.22	0.31
Redband Rockfish												
Guided												
Unguided												
Combined												
Redstripe Rockfish												
Guided												
Unguided												
Combined												
Tiger Rockfish												
Guided		0.13					0.10	0.10	0.11	0.18	0.05	
Unguided				0.02				0.05	0.06	0.13	0.06	0.09
Combined		0.08		0.01				0.06	0.08	0.15	0.05	
Widow Rockfish												
Guided												
Unguided												
Combined												
Yelloweye Rockfish												
Guided												
Unguided												
Combined												
Yellowtail Rockfish												
Guided	0.03	0.44	0.04				0.07	0.07	0.07		0.10	0.10
Unguided	0.25					0.02	0.07	0.10	0.13			
Combined	0.09	0.27	0.02			0.01	0.06	0.09	0.11		0.05	0.08
Unknown rockfish												
Guided		1.19					0.17	0.17	0.19		0.05	
Unguided		0.20				0.02	0.14	0.34	0.42	1.60		0.69
Combined		0.81				0.01	0.13	0.29	0.33	0.92	0.03	

APPENDIX 3: COUNTS OF ACTIVELY FISHING SPORT VESSELS IN EACH SUBAREA, BY DATE

Counts of actively fishing sport vessels in each subarea, by date, 2025. Weekend and holiday flights are shaded. Also shown, the number of lodge boats that would have been visible in Area 3 during flight.

Month	Date	Subarea												Area 3 Lodge Boats
		3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	
May	2 May 2025	0	0	0	0	0	0	0	0	1	0	0	na	0
	9 May 2025	2	1	0	0	0	4	0	0	0	0	0	0	0
	10 May 2025	1	1	2	1	7	2	1	0	1	0	0	0	0
	15 May 2025	2	0	0	3	0	1	0	0	0	0	0	na	0
	18 May 2025	1	1	1	3	11	4	4	0	15	2	3	1	0
	29 May 2025	0	3	5	3	2	1	3	1	2	0	1	0	4
	31 May 2025	1	1	na	2	5	4	2	5	6	0	2	0	0
June	3 Jun 2025	0	3	2	0	1	6	3	1	23	3	1	1	4
	8 Jun 2025	13	2	16	4	3	7	3	2	10	0	4	0	6
	10 Jun 2025	6	6	4	2	2	0	5	1	3	0	7	0	7
	18 Jun 2025	2	10	4	1	2	5	2	1	9	5	1	0	4
	26 Jun 2025	7	1	4	0	3	3	1	0	9	3	1	na	7
	27 Jun 2025	1	na	6	0	1	7	3	0	0	1	3	na	0
	28 Jun 2025	na	na	na	4	1	4	1	na	na	na	na	na	0
July	29 Jun 2025	2	9	1	2	6	8	0	1	6	2	2	na	7
	2 Jul 2025	5	10	7	0	1	6	1	1	13	7	4	0	7
	5 Jul 2025	8	13	0	1	3	9	1	1	17	7	6	0	7
	10 Jul 2025	6	7	4	2	3	17	0	0	2	1	1	0	10
	13 Jul 2025	4	10	2	0	2	14	3	0	69	9	5	7	7
	17 Jul 2025	5	7	6	1	3	7	1	0	5	1	19	na	7
	23 Jul 2025	4	14	16	0	6	18	4	2	19	5	12	1	4
August	26 Jul 2025	11	31	0	3	7	35	5	0	29	12	7	0	7
	29 Jul 2025	7	20	7	3	3	12	13	0	43	16	0	2	4
	3 Aug 2025	na	na	na	3	0	9	24	0	5	na	na	na	0
	6 Aug 2025	5	14	11	0	4	19	14	2	15	5	0	0	3
	17 Aug 2025	8	5	14	6	6	15	1	1	3	2	5	1	0
	19 Aug 2025	7	10	28	0	1	7	3	0	1	0	5	na	4
	22 Aug 2025	8	13	18	0	2	3	4	3	10	4	4	2	7
	28 Aug 2025	4	3	3	2	1	5	1	0	3	3	3	2	4
	29 Aug 2025	6	3	6	1	3	15	1	0	5	2	2	0	0
	30 Aug 2025	6	0	6	0	0	9	3	0	15	1	5	na	4

APPENDIX 4: RETAINED AND RELEASED FISH (LODGE REPORTS AND NON-LODGE CREEL ESTIMATES) BY MONTH, SUBAREA, AND SPECIES.

Estimated numbers of Chinook Salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	64.2	38.0	44.7	4.7	242.7	167.1	100.5	9.3	377.2	11.1	20.3	8.3	1,088
Lodge Report	62.4	37.0	43.5	0	0	0	0	0	0	0	0	0	143
MAY RETAINED	126.6	75.1	88.3	4.7	242.7	167.1	100.5	9.3	377.2	11.1	20.3	8.3	1,231
SE	14.4	6.2	12.1	1.4	39.1	20.0	14.6	2.0	38.0	4.4	9.4	5.4	64
June													
Creel Estimate	136.8	134.1	131.4	0.7	73.2	102.9	148.9	18.6	649.2	35.4	59.5	9.3	1,500
Lodge Report	116.7	114.3	112.0	0	0	0	0	0	0	0	0	0	343
JUNE RETAINED	253.5	248.4	243.4	0.7	73.2	102.9	148.9	18.6	649.2	35.4	59.5	9.3	1,843
SE	24.4	18.9	51.0	0.5	8.1	11.3	18.0	13.5	34.3	10.4	14.0	3.7	76
July													
Creel Estimate	49.5	254.3	74.9	0	21.5	111.7	130.5	1.6	324.7	69.3	15.2	7.9	1,061
Lodge Report	15.0	77.2	22.7	0	0	0	0	0	0	0	0	0	115
JULY RETAINED	64.5	331.5	97.7	0	21.5	111.7	130.5	1.6	324.7	69.3	15.2	7.9	1,176
SE	9.0	24.0	10.7	0	5.6	11.9	12.3	1.4	22.9	19.3	2.8	2.7	45
August													
Creel Estimate	189.7	616.7	289.0	0	38.5	304.0	272.2	16.5	887.8	183.2	61.5	20.0	2,879
Lodge Report	26.0	84.4	39.6	0	0	0	0	0	0	0	0	0	150
AUGUST RETAINED	215.7	701.1	328.6	0	38.5	304.0	272.2	16.5	887.8	183.2	61.5	20.0	3,029
SE	25.9	52.6	34.8	0	7.5	29.2	22.8	3.9	59.8	38.6	10.6	5.3	106
TOTAL													
Creel Estimate	440	1,043	540	5	376	686	652	46	2,239	299	157	46	6,528
Lodge Report	220	313	218	0	0	0	0	0	0	0	0	0	751
TOTAL RETAINED	660	1,356	758	5	376	686	652	46	2,239	299	157	46	7,279
SE	39	61	64	1	41	39	35	14	82	45	20	9	152

Estimated numbers of Chinook Salmon released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	112.5	67.2	114.6	1.7	260.5	111.0	85.9	3.6	182.3	8.3	10.0	0.4	958
Lodge Report	18.0	10.7	18.3	0	0	0	0	0	0	0	0	0	47
MAY RELEASED	130.5	78.0	132.9	1.7	260.5	111.0	85.9	3.6	182.3	8.3	10.0	0.4	1,005
SE	47.9	19.9	44.0	0.9	72.7	17.8	19.6	0.5	23.8	3.7	4.9	0.4	106
June													
Creel Estimate	244.7	398.1	474.9	0.1	104.7	452.9	139.0	9.0	768.3	84.5	175.7	15.8	2,867
Lodge Report	14.0	22.8	27.2	0	0	0	0	0	0	0	0	0	64
JUNE RELEASED	258.7	420.9	502.1	0.1	104.7	452.9	139.0	9.0	768.3	84.5	175.7	15.8	2,931
SE	46.9	52.5	180.2	0.1	15.5	36.8	19.6	5.5	58.2	17.5	26.5	9.6	210
July													
Creel Estimate	425.8	809.5	416.2	0.4	102.1	1105.0	203.7	26.0	1064.4	188.5	403.5	38.0	4,783
Lodge Report	103.9	197.5	101.6	0	0	0	0	0	0	0	0	0	403
JULY RELEASED	529.7	1007.0	517.7	0.4	102.1	1105.0	203.7	26.0	1064.4	188.5	403.5	38.0	5,186
SE	56.6	116.5	99.0	0.3	25.4	88.2	24.4	15.5	109.9	34.2	62.1	30.5	232
August													
Creel Estimate	294.6	473.9	335.8	0	21.7	261.8	144.4	5.0	518.8	88.7	70.4	2.6	2,218
Lodge Report	33.3	53.6	38.0	0	0	0	0	0	0	0	0	0	125
AUGUST RELEASED	327.9	527.6	373.9	0	21.7	261.8	144.4	5.0	518.8	88.7	70.4	2.6	2,343
SE	61.7	98.7	61.2	0	6.0	36.8	20.9	1.4	88.2	26.7	18.0	2.0	167
TOTAL													
Creel Estimate	1,078	1,749	1,342	2	489	1,931	573	44	2,534	370	660	57	10,827
Lodge Report	169	285	185	0	0	0	0	0	0	0	0	0	639
TOTAL RELEASED	1,247	2,033	1,527	2	489	1,931	573	44	2,534	370	660	57	11,466
SE	107	163	219	1	79	104	42	16	154	47	70	32	370

Estimated numbers of Coho Salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	4.4	8.1	10.5	0	0.8	0.8	2.5	0.0	6.8	1.0	0	0.4	35
Lodge Report	0.8	1.4	1.8	0	0	0	0	0	0	0	0	0	4
MAY RETAINED	5.2	9.5	12.4	0	0.8	0.8	2.5	0.0	6.8	1.0	0	0.4	39
SE	2.0	3.0	8.4	0	0.4	0.4	1.5	0.0	1.4	1.0	0	0.4	9
June													
Creel Estimate	303.3	369.0	205.2	0	58.4	912.7	74.4	31.3	928.7	154.3	376.0	92.6	3,506
Lodge Report	97.1	118.2	65.7	0	0	0	0	0	0	0	0	0	281
JUNE RETAINED	400.4	487.1	270.9	0	58.4	912.7	74.4	31.3	928.7	154.3	376.0	92.6	3,787
SE	31.0	47.8	70.4	0	18.5	46.2	9.5	10.7	58.6	24.8	44.1	31.9	134
July													
Creel Estimate	1420.0	2068.1	1031.4	0	265.3	3088.6	777.3	100.6	4383.9	718.4	1152.0	322.3	15,328
Lodge Report	691.6	1007.2	502.3	0	0	0	0	0	0	0	0	0	2,201
JULY RETAINED	2111.6	3075.2	1533.7	0	265.3	3088.6	777.3	100.6	4383.9	718.4	1152.0	322.3	17,529
SE	87.9	146.0	122.3	0	52.2	132.5	51.4	30.6	199.9	97.0	120.2	109.6	379
August													
Creel Estimate	2145.6	2756.9	2917.8	0	259.5	1206.5	1249.5	28.5	3277.7	608.3	383.6	91.1	14,925
Lodge Report	494.4	635.3	672.3	0	0	0	0	0	0	0	0	0	1,802
AUGUST RETAINED	2639.9	3392.2	3590.2	0	259.5	1206.5	1249.5	28.5	3277.7	608.3	383.6	91.1	16,727
SE	120.8	186.8	172.3	0	28.4	85.2	83.3	10.4	139.2	124.6	55.9	32.5	365
TOTAL													
Creel Estimate	3,873	5,202	4,165	0	584	5,209	2,104	160	8,597	1,482	1,911	506	33,794
Lodge Report	1,284	1,762	1,242	0	0	0	0	0	0	0	0	0	4,288
TOTAL RETAINED	5,157	6,964	5,407	0	584	5,209	2,104	160	8,597	1,482	1,911	506	38,082
SE	153	242	223	0	62	164	98	34	251	160	140	119	543

Estimated numbers of Coho Salmon released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0.2	0	1.4	0.3	0	0	2
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0	0	0	0	0	0	0.2	0	1.4	0.3	0	0	2
SE	0	0	0	0	0	0	0.2	0	0.5	0.3	0	0	1
June													
Creel Estimate	77.5	2.8	9.3	0	8.3	52.2	5.8	0	80.3	10.7	7.7	0	255
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	77.5	2.8	9.3	0	8.3	52.2	5.8	0	80.3	10.7	7.7	0	255
SE	42.2	2.3	9.0	0	4.1	11.9	2.6	0	22.9	4.0	3.8	0	51
July													
Creel Estimate	283.3	184.5	65.2	0	32.5	184.8	72.9	0	312.2	86.0	29.1	0	1,250
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	283.3	184.5	65.2	0	32.5	184.8	72.9	0	312.2	86.0	29.1	0	1,250
SE	120.0	46.9	26.9	0	12.0	33.8	14.6	0	74.6	35.7	10.7	0	161
August													
Creel Estimate	200.0	395.8	281.1	0	19.1	100.6	123.6	0.5	187.7	125.6	23.2	0	1,457
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	200.0	395.8	281.1	0	19.1	100.6	123.6	0.5	187.7	125.6	23.2	0	1,457
SE	61.0	92.5	75.6	0	8.0	22.4	24.1	0.3	36.9	61.9	10.5	0	156
TOTAL													
Creel Estimate	561	583	356	0	60	338	203	1	582	223	60	0	2,964
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	561	583	356	0	60	338	203	1	582	223	60	0	2,964
SE	141	104	81	0	15	42	28	0	86	72	15	0	230

Estimated numbers of Pink Salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0.4	0	0	0.3	0	0	0	1
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	0	0	0	0	0	0.4	0	0	0.3	0	0	0	1
SE	0	0	0	0	0	0.2	0	0	0.2	0	0	0	0
June													
Creel Estimate	21.4	50.8	48.8	0	0	73.4	14.9	10.7	92.3	5.7	15.3	0	333
Lodge Report	0.9	2.1	2.0	0	0	0	0	0	0	0	0	0	5
JUNE RETAINED	22.3	52.9	50.8	0	0	73.4	14.9	10.7	92.3	5.7	15.3	0	338
SE	5.7	12.9	24.1	0	0	9.9	3.5	5.5	13.3	3.0	5.2	0	34
July													
Creel Estimate	154.1	202.9	208.0	0	9.2	279.9	92.0	31.2	456.7	28.6	51.7	2.0	1,516
Lodge Report	1.1	1.4	1.5	0	0	0	0	0	0	0	0	0	4
JULY RETAINED	155.2	204.3	209.4	0	9.2	279.9	92.0	31.2	456.7	28.6	51.7	2.0	1,520
SE	21.2	39.3	66.4	0	2.5	29.1	13.0	15.4	49.2	10.2	14.4	1.3	102
August													
Creel Estimate	273.1	131.7	243.4	0	21.8	178.6	114.7	1.4	346.8	27.0	35.1	6.2	1,380
Lodge Report	4.2	2.0	3.8	0	0	0	0	0	0	0	0	0	10
AUGUST RETAINED	277.3	133.7	247.2	0	21.8	178.6	114.7	1.4	346.8	27.0	35.1	6.2	1,390
SE	34.4	34.3	37.6	0	5.3	24.2	16.5	0.4	44.5	10.3	8.8	2.9	83
TOTAL													
Creel Estimate	449	385	500	0	31	532	222	43	896	61	102	8	3,230
Lodge Report	6	6	7	0	0	0	0	0	0	0	0	0	19
TOTAL RETAINED	455	391	507	0	31	532	222	43	896	61	102	8	3,249
SE	41	54	80	0	6	39	21	16	68	15	18	3	135

Estimated numbers of Pink Salmon released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0	0	0	0	0	0	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
June													
Creel Estimate	90.0	136.9	83.6	0	15.6	282.8	20.7	17.9	213.1	61.4	67.4	0	989
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	90.0	136.9	83.6	0	15.6	282.8	20.7	17.9	213.1	61.4	67.4	0	989
SE	24.8	40.2	33.8	0	7.6	39.3	7.1	9.2	32.2	21.1	14.6	0	82
July													
Creel Estimate	542.0	926.2	524.6	0	94.8	1132.9	342.8	55.2	1401.4	340.9	222.5	9.2	5,593
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	542.0	926.2	524.6	0	94.8	1132.9	342.8	55.2	1401.4	340.9	222.5	9.2	5,593
SE	83.1	135.1	102.1	0	23.5	114.4	37.3	25.8	120.9	76.2	40.1	5.8	271
August													
Creel Estimate	780.4	1192.8	736.2	0	111.0	746.0	592.6	7.5	1417.9	318.7	99.9	22.5	6,025
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	780.4	1192.8	736.2	0	111.0	746.0	592.6	7.5	1417.9	318.7	99.9	22.5	6,025
SE	107.2	154.6	95.0	0	21.6	81.1	58.7	5.0	112.2	80.7	21.5	10.7	273
TOTAL													
Creel Estimate	1,412	2,256	1,344	0	221	2,162	956	81	3,032	721	390	32	12,607
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	1,412	2,256	1,344	0	221	2,162	956	81	3,032	721	390	32	12,607
SE	138	209	143	0	33	146	70	28	168	113	48	12	393

Estimated numbers of Chum Salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	0	0	0	0	0	0	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
June													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0.3	0.3	0.3	0	0	0	0	0	0	0	0	0	1
JUNE RETAINED	0.3	0.3	0.3	0	0	0	0	0	0	0	0	0	1
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
July													
Creel Estimate	0	0	0	0	0	0	0	0	1.2	0	0	0	1
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RETAINED	0	0	0	0	0	0	0	0	1.2	0	0	0	1
SE	0	0	0	0	0	0	0	0	1.1	0	0	0	1
August													
Creel Estimate	0	0	0	0	0	0	0	0	2.3	0	0	0	2
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RETAINED	0	0	0	0	0	0	0	0	2.3	0	0	0	2
SE	0	0	0	0	0	0	0	0	1.8	0	0	0	2
TOTAL													
Creel Estimate	0	0	0	0	0	0	0	0	3	0	0	0	3
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	1
TOTAL RETAINED	0	0	0	0	0	0	0	0	3	0	0	0	4
SE	0	0	0	0	0	0	0	0	2	0	0	0	

Estimated numbers of Chum Salmon released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0.7	0.3	0	0	0	0	0	0	0.5	0.3	0	0	2
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0.7	0.3	0	0	0	0	0	0	0.5	0.3	0	0	2
SE	0.6	0.3	0	0	0	0	0	0	0.3	0.3	0	0	1
June													
Creel Estimate	5.3	6.6	0	0	0	2.0	0.4	0	7.1	2.5	0	0	24
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	5.3	6.6	0	0	0	2.0	0.4	0	7.1	2.5	0	0	24
SE	4.0	3.2	0	0	0	1.0	0.3	0	2.7	2.0	0	0	6
July													
Creel Estimate	0	12.1	12.8	0	0	6.8	2.2	0	13.4	0	0	0	47
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	0	12.1	12.8	0	0	6.8	2.2	0	13.4	0	0	0	47
SE	0	6.9	5.6	0	0	2.9	1.3	0	5.6	0	0	0	11
August													
Creel Estimate	3.1	0	31.5	0	0	2.3	2.4	0	6.8	0	0	0	46
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	3.1	0	31.5	0	0	2.3	2.4	0	6.8	0	0	0	46
SE	2.4	0	11.3	0	0	1.6	1.8	0	3.1	0	0	0	12
TOTAL													
Creel Estimate	9	19	44	0	0	11	5	0	28	3	0	0	119
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	9	19	44	0	0	11	5	0	28	3	0	0	119
SE	5	8	13	0	0	3	2	0	7	2	0	0	18

Estimated numbers of Sockeye Salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	0	0	0	0	0	0	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
June													
Creel Estimate	0	0	0	0	0	0	0	0	4.2	0	0	0	4
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RETAINED	0	0	0	0	0	0	0	0	4.2	0	0	0	4
SE	0	0	0	0	0	0	0	0	3.0	0	0	0	3
July													
Creel Estimate	3.3	2.3	1.8	0	0	0	1.2	0	14.3	0	0.5	0	23
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RETAINED	3.3	2.3	1.8	0	0	0	1.2	0	14.3	0	0.5	0	23
SE	1.7	1.8	1.5	0	0	0	1.0	0	9.5	0	0.4	0	10
August													
Creel Estimate	7.6	4.9	3.8	0	0	0	2.4	0	2.3	0	4.5	0	25
Lodge Report	1	1	1	0	0	0	0	0	0	0	0	0	3
AUGUST RETAINED	9.0	5.8	4.5	0	0	0	2.4	0	2.3	0	4.5	0	28
SE	3.8	3.5	2.8	0	0	0	1.8	0	1.8	0	2.5	0	7
TOTAL													
Creel Estimate	11	7	6	0	0	0	4	0	21	0	5	0	53
Lodge Report	1	1	1	0	0	0	0	0	0	0	0	0	3
TOTAL RETAINED	12	8	6	0	0	0	4	0	21	0	5	0	56
SE	4	4	3	0	0	0	2	0	10	0	3	0	12

Estimated numbers of Sockeye Salmon released by lodge and non-lodge anglers, by month, 2025.

Month / Catch Type	Subarea												TOTAL
	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	
May													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0	0	0	0	0	0	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
June													
Creel Estimate	0	2.8	0	0	0	0	0	0	0	0	0	0	3
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	0	2.8	0	0	0	0	0	0	0	0	0	0	3
SE	0	2.3	0	0	0	0	0	0	0	0	0	0	2
July													
Creel Estimate	0	10.4	0	0	0	1.0	3.5	0	2.4	0	0	0	17
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	0	10.4	0	0	0	1.0	3.5	0	2.4	0	0	0	17
SE	0	6.5	0	0	0	0.8	2.9	0	2.1	0	0	0	7
August													
Creel Estimate	0	4.9	4.9	0	0	2.3	7.1	0	4.5	0	0	0	24
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	0	4.9	4.9	0	0	2.3	7.1	0	4.5	0	0	0	24
SE	0	3.5	4.3	0	0	1.6	5.3	0	3.6	0	0	0	9
TOTAL													
Creel Estimate	0	18	5	0	0	3	11	0	7	0	0	0	44
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	0	18	5	0	0	3	11	0	7	0	0	0	44
SE	0	8	4	0	0	2	6	0	4	0	0	0	12

Estimated numbers of unknown salmon retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0	0	0	0	0	0	0
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	0	0	0	0	0	0	0	0	0	0	0	0	0
SE	0	0	0	0	0	0	0	0	0	0	0	0	0
June													
Creel Estimate	0	0	0	0	0	0	0	0.9	0	1.6	0	0	3
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RETAINED	0	0	0	0	0	0	0	0.9	0	1.6	0	0	3
SE	0	0	0	0	0	0	0	0.7	0	1.2	0	0	1
July													
Creel Estimate	0	0	0	0	0	0	2.3	2.6	6.0	4.9	0	13.2	29
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RETAINED	0	0	0	0	0	0	2.3	2.6	6.0	4.9	0	13.2	29
SE	0	0	0	0	0	0	1.9	1.9	4.3	3.5	0	9.3	11
August													
Creel Estimate	0	0	0	0	0	0	4.8	0	11.3	0	0	25.5	42
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RETAINED	0	0	0	0	0	0	4.8	0	11.3	0	0	25.5	42
SE	0	0	0	0	0	0	3.5	0	7.5	0	0	16.3	18
TOTAL													
Creel Estimate	0	0	0	0	0	0	7	3	17	7	0	39	73
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RETAINED	0	0	0	0	0	0	7	3	17	7	0	39	73
SE	0	0	0	0	0	0	4	2	9	4	0	19	21

Estimated numbers of unknown salmon released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	0	0	0	0	0	0.6	0	0	0	0	0	1
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0	0	0	0	0	0	0.6	0	0	0	0	0	1
SE	0	0	0	0	0	0	0.5	0	0	0	0	0	0
June													
Creel Estimate	0	0	0	0	0	1.5	4.4	0	0	0	0	0	6
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	0	0	0	0	0	1.5	4.4	0	0	0	0	0	6
SE	0	0	0	0	0	1.2	3.5	0	0	0	0	0	4
July													
Creel Estimate	0	0	3.7	0	0.6	4.3	8.2	0	0	0	0	0	17
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	0	0	3.7	0	0.6	4.3	8.2	0	0	0	0	0	17
SE	0	0	3.0	0	0.4	3.4	5.2	0	0	0	0	0	7
August													
Creel Estimate	0	0	7.6	0	1.3	0	16.6	0	0	0	0	0	26
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	0	0	7.6	0	1.3	0	16.6	0	0	0	0	0	26
SE	0	0	5.5	0	0.9	0	9.4	0	0	0	0	0	11
TOTAL													
Creel Estimate	0	0	11	0	2	6	30	0	0	0	0	0	49
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	0	0	11	0	2	6	30	0	0	0	0	0	49
SE	0	0	6	0	1	4	11	0	0	0	0	0	13

Estimated numbers of Pacific Halibut retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	90.5	42.3	21.1	1.7	10.6	67.0	33.8	14.2	267.7	41.6	84.3	33.8	709
Lodge Report	27.1	12.6	6.3	0	0	0	0	0	0	0	0	0	46
MAY RETAINED	117.6	54.9	27.4	1.7	10.6	67.0	33.8	14.2	267.7	41.6	84.3	33.8	755
SE	21.5	8.9	9.4	1.2	4.5	12.9	8.2	4.3	31.3	7.4	16.0	7.0	47
June													
Creel Estimate	216.4	191.3	212.9	0.1	60.2	187.5	104.9	91.5	821.6	157.9	342.2	66.7	2,453
Lodge Report	79.5	70.3	78.2	0	0	0	0	0	0	0	0	0	228
JUNE RETAINED	295.9	261.6	291.1	0.1	60.2	187.5	104.9	91.5	821.6	157.9	342.2	66.7	2,681
SE	43.1	26.1	70.2	0.0	10.3	14.8	14.3	31.0	41.3	22.6	27.1	8.8	109
July													
Creel Estimate	198.8	585.4	245.4	0.2	120.1	455.5	146.1	68.6	1295.7	299.2	432.5	174.4	4,022
Lodge Report	59.5	175.1	73.4	0	0	0	0	0	0	0	0	0	308
JULY RETAINED	258.3	760.6	318.9	0.2	120.1	455.5	146.1	68.6	1295.7	299.2	432.5	174.4	4,330
SE	23.7	53.5	61.5	0.1	23.2	31.5	15.1	15.2	68.3	32.4	34.4	26.7	130
August													
Creel Estimate	288.1	920.6	218.7	0	76.8	610.4	247.9	53.9	864.7	185.2	327.1	118.9	3,912
Lodge Report	49.5	158.0	37.5	0	0	0	0	0	0	0	0	0	245
AUGUST RETAINED	337.6	1078.6	256.3	0	76.8	610.4	247.9	53.9	864.7	185.2	327.1	118.9	4,157
SE	38.3	81.0	37.9	0	10.5	45.2	26.5	8.3	54.7	37.1	28.2	15.9	133
TOTAL													
Creel Estimate	794	1,740	698	2	268	1,320	533	228	3,250	684	1,186	394	11,096
Lodge Report	215	416	195	0	0	0	0	0	0	0	0	0	827
TOTAL RETAINED	1,009	2,156	894	2	268	1,320	533	228	3,250	684	1,186	394	11,923
SE	66	101	101	1	28	58	35	36	102	55	54	33	221

Estimated numbers of Pacific Halibut released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	66.4	31.9	0	0	17.9	77.0	11.2	10.0	168.8	27.6	54.9	11.3	477
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	66.4	31.9	0	0	17.9	77.0	11.2	10.0	168.8	27.6	54.9	11.3	477
SE	25.1	11.6	0	0	10.1	26.5	3.2	3.2	44.1	9.6	22.9	6.8	65
June													
Creel Estimate	178.5	210.0	39.5	0	30.5	119.2	35.3	63.9	478.5	193.9	341.9	82.7	1,774
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	178.5	210.0	39.5	0	30.5	119.2	35.3	63.9	478.5	193.9	341.9	82.7	1,774
SE	45.9	52.0	24.2	0	11.1	17.6	12.0	22.9	53.0	41.3	45.0	33.4	119
July													
Creel Estimate	114.2	785.5	157.4	0	19.3	290.4	29.2	35.4	774.8	473.3	627.1	229.7	3,536
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	114.2	785.5	157.4	0	19.3	290.4	29.2	35.4	774.8	473.3	627.1	229.7	3,536
SE	27.0	96.1	68.4	0	6.0	41.5	7.7	16.4	81.7	114.6	91.5	112.4	240
August													
Creel Estimate	138.6	1220.7	121.1	0	28.4	313.8	72.0	31.9	654.9	469.2	316.4	159.2	3,526
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	138.6	1220.7	121.1	0	28.4	313.8	72.0	31.9	654.9	469.2	316.4	159.2	3,526
SE	33.1	187.4	42.5	0	7.1	43.9	15.1	10.3	90.1	185.2	62.9	41.5	297
TOTAL													
Creel Estimate	498	2,248	318	0	96	800	148	141	2,077	1,164	1,340	483	9,313
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	498	2,248	318	0	96	800	148	141	2,077	1,164	1,340	483	9,313
SE	68	217	84	0	18	68	21	30	140	222	122	125	405

Estimated numbers of Lingcod retained by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	63.6	54.7	21.7	0	51.8	23.9	4.9	18.5	289.1	37.3	5.2	54.7	625
Lodge Report	75.9	65.2	25.8	0	0	0	0	0	0	0	0	0	167
MAY RETAINED	139.5	119.9	47.5	0	51.8	23.9	4.9	18.5	289.1	37.3	5.2	54.7	792
SE	38.6	17.1	13.5	0	31.2	11.6	2.0	8.4	55.4	11.3	1.8	13.5	81
June													
Creel Estimate	23.8	422.3	128.9	0	25.1	71.3	33.3	118.5	551.9	151.9	71.5	47.3	1,646
Lodge Report	23.9	423.8	129.3	0	0	0	0	0	0	0	0	0	577
JUNE RETAINED	47.7	846.1	258.2	0	25.1	71.3	33.3	118.5	551.9	151.9	71.5	47.3	2,223
SE	9.5	60.9	89.0	0	10.0	12.8	10.8	60.1	48.7	31.6	16.7	13.3	140
July													
Creel Estimate	56.2	966.1	93.1	0	32.4	166.7	73.1	70.8	868.1	349.9	118.6	112.9	2,908
Lodge Report	33.4	575.2	55.4	0	0	0	0	0	0	0	0	0	664
JULY RETAINED	89.6	1541.3	148.5	0	32.4	166.7	73.1	70.8	868.1	349.9	118.6	112.9	3,572
SE	17.0	102.5	34.9	0	13.3	28.4	14.5	25.5	80.2	61.0	28.3	36.8	162
August													
Creel Estimate	110.8	996.2	112.4	2.9	52.5	109.9	142.7	59.5	737.0	335.4	130.0	151.2	2,941
Lodge Report	51.5	463.2	52.3	0	0	0	0	0	0	0	0	0	567
AUGUST RETAINED	162.3	1459.5	164.6	2.9	52.5	109.9	142.7	59.5	737.0	335.4	130.0	151.2	3,508
SE	36.4	139.1	42.5	2.3	20.6	21.9	31.9	11.6	71.3	84.8	31.8	32.9	197
TOTAL													
Creel Estimate	254	2,439	356	3	162	372	254	267	2,446	874	325	366	8,119
Lodge Report	185	1,527	263	0	0	0	0	0	0	0	0	0	1,975
TOTAL RETAINED	439	3,967	619	3	162	372	254	267	2,446	874	325	366	10,094
SE	57	184	106	2	41	40	37	67	130	110	46	53	302

Estimated numbers of Lingcod released by lodge and non-lodge anglers, by month, 2025.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	0	1.0	0	0	21.2	0	0.6	5.4	15.1	4.2	1.1	0	49
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	0	1.0	0	0	21.2	0	0.6	5.4	15.1	4.2	1.1	0	49
SE	0	0.8	0	0	11.7	0	0.5	3.8	8.9	2.3	0.6	0	15
June													
Creel Estimate	0.9	43.9	23.2	0	0	5.1	6.6	40.8	44.6	55.8	8.9	0	230
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	0.9	43.9	23.2	0	0	5.1	6.6	40.8	44.6	55.8	8.9	0	230
SE	0.7	23.6	22.6	0	0	2.1	3.6	27.1	16.4	21.3	4.5	0	51
July													
Creel Estimate	2.6	137.3	62.5	0	0	20.6	9.6	10.4	122.2	126.1	4.3	0.7	496
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	2.6	137.3	62.5	0	0	20.6	9.6	10.4	122.2	126.1	4.3	0.7	496
SE	1.9	64.0	60.3	0	0	6.2	3.5	7.7	50.1	60.1	2.4	0.6	118
August													
Creel Estimate	0	82.2	0	0	1.3	19.6	13.4	2.2	64.8	135.1	5.8	1.3	326
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	0	82.2	0	0	1.3	19.6	13.4	2.2	64.8	135.1	5.8	1.3	326
SE	0	29.9	0	0	1.1	6.7	5.5	1.4	22.4	88.4	5.0	1.0	97
TOTAL													
Creel Estimate	3	264	86	0	23	45	30	59	247	321	20	2	1,100
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	3	264	86	0	23	45	30	59	247	321	20	2	1,100
SE	2	74	64	0	12	9	7	28	58	109	7	1	162

Estimated numbers of other groundfish retained by lodge and non-lodge anglers, by month, 2025. Included are Big and Longnose skates, Pacific Spiny Dogfish, Pacific Cod, Sablefish, Cabezon, Red Irish Lord, greenlings, Spotted Ratfish, Arrow-tooth and Starry flounders, Pacific Sanddab, Pacific Rock and English soles, and other miscellaneous groundfishes.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	28.6	9.1	8.8	0	7.2	2.3	1.8	0.6	39.6	44.0	143.5	0.4	286
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	28.6	9.1	8.8	0	7.2	2.3	1.8	0.6	39.6	44.0	143.5	0.4	286
SE	12.4	4.7	7.5	0	6.4	1.3	0.6	0.4	25.6	17.7	45.8	0.4	58
June													
Creel Estimate	77.5	15.5	20.9	0	7.6	22.2	23.7	0	77.8	178.9	646.3	35.7	1,106
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RETAINED	77.5	15.5	20.9	0	7.6	22.2	23.7	0	77.8	178.9	646.3	35.7	1,106
SE	33.2	5.6	14.5	0	3.0	5.5	8.6	0	14.8	71.0	92.3	21.4	125
July													
Creel Estimate	104.5	94.9	63.5	0.1	29.4	87.5	46.0	0.8	218.5	199.4	746.8	109.9	1,701
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RETAINED	104.5	94.9	63.5	0.1	29.4	87.5	46.0	0.8	218.5	199.4	746.8	109.9	1,701
SE	29.7	25.4	38.8	0.1	11.1	16.5	20.3	0.7	45.1	79.1	101.9	72.8	167
August													
Creel Estimate	198.5	126.0	34.9	0.2	26.9	117.7	47.9	7.2	278.4	119.7	159.1	93.4	1,210
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RETAINED	198.5	126.0	34.9	0.2	26.9	117.7	47.9	7.2	278.4	119.7	159.1	93.4	1,210
SE	65.8	41.2	15.1	0.2	16.4	24.2	12.4	2.5	69.3	86.9	36.7	30.5	148
TOTAL													
Creel Estimate	409	245	128	0	71	230	119	9	614	542	1,696	239	4,303
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RETAINED	409	245	128	0	71	230	119	9	614	542	1,696	239	4,303
SE	80	49	45	0	21	30	25	3	88	138	150	82	262

Estimated numbers of other groundfish released by lodge and non-lodge anglers, by month, 2025. Included are Big and Longnose skates, Pacific Spiny Dogfish, Pacific Cod, Sablefish, Cabezon, Red Irish Lord, greenlings, Spotted Ratfish, Arrow-tooth and Starry flounders, Pacific Sanddab, Pacific Rock and English soles, and other miscellaneous groundfishes.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	8.9	3.6	0	0	6.0	29.3	20.7	0.8	22.3	1.7	8.2	0	101
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	8.9	3.6	0	0	6.0	29.3	20.7	0.8	22.3	1.7	8.2	0	101
SE	4.6	1.8	0	0	3.3	12.7	14.4	0.3	11.8	1.5	4.7	0	24
June													
Creel Estimate	28.4	0	0	0.4	24.0	34.6	26.5	0	47.7	9.0	85.6	0	256
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	28.4	0	0	0.4	24.0	34.6	26.5	0	47.7	9.0	85.6	0	256
SE	14.6	0	0	0.2	6.6	5.7	12.2	0	9.5	4.4	19.8	0	125
July													
Creel Estimate	86.7	18.3	31.1	1.1	19.6	67.9	22.2	0	181.6	34.6	177.5	0	641
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	86.7	18.3	31.1	1.1	19.6	67.9	22.2	0	181.6	34.6	177.5	0	641
SE	41.3	9.5	11.9	0.6	6.5	12.7	7.3	0	38.1	14.4	48.1	0	79
August													
Creel Estimate	45.7	39.3	84.3	0	6.5	27.8	45.2	2.3	182.1	15.5	69.0	0	518
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	45.7	39.3	84.3	0	6.5	27.8	45.2	2.3	182.1	15.5	69.0	0	518
SE	29.8	18.5	26.1	0	2.9	7.2	13.4	1.5	51.5	11.6	15.1	0	72
TOTAL													
Creel Estimate	170	61	115	2	56	160	114	3	434	61	340	0	1,516
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	170	61	115	2	56	160	114	3	434	61	340	0	1,516
SE	53	21	29	1	10	20	24	2	66	19	54	0	113

Estimated numbers of other groundfish retained by lodge and non-lodge anglers, by taxon, 2025.

Month / Catch Type	Taxon															TOTAL
	Arrow-tooth Flounder	Big Skate	Cabezon	English Sole	Greenlings	Longnose Skate	Pacific Cod	Pacific Sanddab	Pacific Rock Sole	Spiny Dogfish	Red Irish Lord	Sablefish	Spotted Ratfish	Starry Flounder	Unknown / Other	
May																
Creel Estimate	0	0.3	2.6	0	27.2	0	255.0	0	0	0.3	0	0	0	0	0.5	286
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RETAINED	0	0.3	2.6	0	27.2	0	255.0	0	0	0.3	0	0	0	0	0.5	286
SE	0	0.2	2.1	0	12.8	0	56.3	0	0	0.2	0	0	0	0	0.4	58
June																
Creel Estimate	1.6	2.0	4.9	0	40.6	0	1040.0	0	4.1	2.0	0	4.6	0	0	6.5	1,106
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RETAINED	1.6	2.0	4.9	0	40.6	0	1040.0	0	4.1	2.0	0	4.6	0	0	6.5	1,106
SE	0.9	1.5	1.9	0	15.3	0	124.2	0	2.1	1.5	0	4.5	0	0	3.4	125
July																
Creel Estimate	7.7	0	12.0	0.5	125.5	0	1520.0	0	12.8	0.1	0	14.3	0	0	8.3	1,701
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RETAINED	7.7	0	12.0	0.5	125.5	0	1520.0	0	12.8	0.1	0	14.3	0	0	8.3	1,701
SE	3.4	0	4.5	0.4	41.4	0	160.8	0	6.9	0.1	0	12.2	0	0	4.0	167
August																
Creel Estimate	5.8	0	21.3	1.4	70.5	0	1097.5	0	1.3	5.9	0	3.8	0	0	2.3	1,210
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RETAINED	5.8	0	21.3	1.4	70.5	0	1097.5	0	1.3	5.9	0	3.8	0	0	2.3	1,210
SE	3.7	0	9.9	1.2	14.8	0	146.7	0	1.1	3.8	0	2.8	0	0	1.6	148
TOTAL																
Creel Estimate	15	2	41	2	264	0	3,912	0	18	8	0	23	0	0	18	4,303
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RETAINED	15	2	41	2	264	0	3,912	0	18	8	0	23	0	0	18	4,303
SE	5	2	11	1	48	0	257	0	7	4	0	13	0	0	5	262

* excludes Lingcod and Pacific Halibut

Estimated numbers of other groundfish released by lodge and non-lodge anglers, by taxon, 2025.

Month / Catch Type	Taxon															TOTAL
	Arrow-tooth Flounder	Big Skate	Cabezon	English Sole	Greenlings	Longnose Skate	Pacific Cod	Pacific Sanddab	Pacific Rock Sole	Spiny Dogfish	Red Irish Lord	Sablefish	Spotted Ratfish	Starry Flounder	Unknown / Other	
May																
Creel Estimate	10.8	2.2	0	0.0	47.5	0	30.2	0	0	7.1	0	0	2.7	0.4	0.5	101
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	10.8	2.2	0	0.0	47.5	0	30.2	0	0	7.1	0	0	2.7	0.4	0.5	101
SE	3.9	0.9	0	0.0	21.6	0	8.7	0	0	2.2	0	0	2.1	0.2	0.3	24
June																
Creel Estimate	45.0	22.1	3.1	0.8	12.0	0	73.4	0	4.0	76.2	0	0	9.6	6.6	3.5	256
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	45.0	22.1	3.1	0.8	12.0	0	73.4	0	4.0	76.2	0	0	9.6	6.6	3.5	256
SE	15.6	6.9	1.4	0.6	3.6	0	19.0	0	1.7	15.9	0	0	2.7	2.1	1.9	31
July																
Creel Estimate	152.1	29.2	24.1	12.5	23.9	0	174.2	0	18.9	164.8	0	0	28.3	10.6	2.1	641
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	152.1	29.2	24.1	12.5	23.9	0	174.2	0	18.9	164.8	0	0	28.3	10.6	2.1	641
SE	47.8	6.3	10.0	7.6	6.7	0	51.5	0	6.6	29.6	0	0	7.1	4.1	1.1	79
August																
Creel Estimate	106.3	27.1	31.7	21.4	43.3	0	34.8	0	24.0	181.8	0	0	42.6	0	4.6	518
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	106.3	27.1	31.7	21.4	43.3	0	34.8	0	24.0	181.8	0	0	42.6	0	4.6	518
SE	45.9	6.9	18.0	13.9	16.0	0	11.6	0	9.8	43.5	0	0	10.7	0	2.3	72
TOTAL																
Creel Estimate	314	81	59	35	127	0	313	0	47	430	0	0	83	18	11	1,516
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	314	81	59	35	127	0	313	0	47	430	0	0	83	18	11	1,516
SE	68	12	21	16	28	0	57	0	12	55	0	0	13	5	3	113

* excludes Lingcod and Pacific Halibut

Estimated numbers of rockfish retained by lodge and non-lodge anglers, by month, 2025. Included are Black, Boccacio, Canary, China, Copper, Dusky, Quillback, Redbanded, Redstripe, Tiger, Widow, Yelloweye, Yellowtail, and other miscellaneous rockfishes.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	23.3	24.5	5.8	0	61.9	65.2	19.8	15.4	213.7	33.4	13.3	67.3	544
Lodge Report	0	33.0	0	0	0	0	0	0	0	0	0	0	33
MAY RETAINED	23.3	57.5	5.8	0	61.9	65.2	19.8	15.4	213.7	33.4	13.3	67.3	577
SE	7.7	4.5	3.4	0	33.7	16.8	6.2	5.0	41.2	13.3	4.0	22.8	63
June													
Creel Estimate	103.0	296.5	41.8	0.2	63.6	102.5	55.5	86.5	504.9	84.7	133.3	76.0	1,548
Lodge Report	16.2	96.4	84.4	0	0	0	0	0	0	0	0	0	197
JUNE RETAINED	119.2	393.0	126.2	0.2	63.6	102.5	55.5	86.5	504.9	84.7	133.3	76.0	1,745
SE	39.8	34.8	16.7	0.1	10.8	13.8	8.7	35.1	42.6	21.1	30.1	25.2	92
July													
Creel Estimate	122.3	747.4	154.5	0.7	83.4	226.1	166.8	18.2	958.0	247.7	119.6	206.4	3,051
Lodge Report	37.3	83.4	137.3	0	0	0	0	0	0	0	0	0	258
JULY RETAINED	159.6	830.8	291.8	0.7	83.4	226.1	166.8	18.2	958.0	247.7	119.6	206.4	3,309
SE	33.0	67.4	46.1	0.4	17.7	26.1	23.4	6.5	82.8	43.9	19.6	82.5	159
August													
Creel Estimate	137.6	872.7	107.1	1.2	37.2	233.1	263.1	40.4	897.9	337.4	115.6	133.3	3,177
Lodge Report	27	272	0	0	0	0	0	0	0	0	0	0	299
AUGUST RETAINED	164.9	1144.5	107.1	1.2	37.2	233.1	263.1	40.4	897.9	337.4	115.6	133.3	3,476
SE	26.6	106.3	24.9	0.6	7.9	29.1	36.2	9.0	71.8	82.4	21.7	30.1	168
TOTAL													
Creel Estimate	386	1,941	309	2	246	627	505	160	2,574	703	382	483	8,320
Lodge Report	81	485	222	0	0	0	0	0	0	0	0	0	787
TOTAL RETAINED	467	2,426	531	2	246	627	505	160	2,574	703	382	483	9,107
SE	59	131	55	1	40	45	44	37	125	97	42	94	257

Estimated numbers of rockfish released by lodge and non-lodge anglers, by month, 2025. Included are Black, Boccacio, Canary, China, Copper, Dusky, Quillback, Redbanded, Redstripe, Tiger, Widow, Yelloweye, Yellowtail, and other miscellaneous rockfishes.

	Subarea												
Month / Catch Type	3I	3J	3K	4A	4B	4C	4D	4E	4F	4G	4H	4L	TOTAL
May													
Creel Estimate	5.9	8.7	0	3.5	198.2	71.6	19.0	12.8	155.2	22.0	15.6	24.7	537
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	5.9	8.7	0	3.5	198.2	71.6	19.0	12.8	155.2	22.0	15.6	24.7	537
SE	2.2	3.0	0	1.8	52.5	10.8	4.3	7.6	22.4	7.4	6.8	11.7	61
June													
Creel Estimate	52.8	169.1	67.4	0	88.3	165.6	100.9	74.5	405.9	145.6	80.0	33.5	1,384
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	52.8	169.1	67.4	0	88.3	165.6	100.9	74.5	405.9	145.6	80.0	33.5	1,384
SE	16.4	44.1	48.9	0	16.6	24.2	15.3	54.0	41.5	36.4	25.4	16.6	112
July													
Creel Estimate	75.4	436.8	248.8	0	104.4	271.5	202.5	13.6	874.1	441.4	121.2	105.9	2,896
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	75.4	436.8	248.8	0	104.4	271.5	202.5	13.6	874.1	441.4	121.2	105.9	2,896
SE	20.6	108.8	133.3	0	30.5	34.0	28.3	5.6	82.8	107.7	54.7	56.3	240
August													
Creel Estimate	146.9	420.4	387.5	0.5	29.5	338.0	254.5	42.4	808.2	397.5	190.0	62.4	3,078
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	146.9	420.4	387.5	0.5	29.5	338.0	254.5	42.4	808.2	397.5	190.0	62.4	3,078
SE	38.9	77.6	141.9	0.4	8.0	46.0	38.1	10.4	99.1	108.6	45.9	22.6	236
TOTAL													
Creel Estimate	281	1,035	704	4	420	847	577	143	2,243	1,007	407	227	7,894
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	281	1,035	704	4	420	847	577	143	2,243	1,007	407	227	7,894
SE	47	141	201	2	64	63	50	56	138	157	76	64	360

Estimated numbers of rockfish retained by lodge and non-lodge anglers, by taxon, 2025.

	Taxon														
Month / Catch Type	Black Rockfish	Boccacio Rockfish	Canary Rockfish	China Rockfish	Copper Rockfish	Dusky Rockfish	Quillback Rockfish	Redbande d Rockfish	Redstripe Rockfish	Tiger Rockfish	Widow Rockfish	Yelloweye Rockfish	Yellowtail Rockfish	Unknown rockfish	TOTAL
May															
Creel Estimate	235.2	0.7	0.3	12.7	25.0	7.0	106.7	4.7	0	20.3	0.5	2.7	13.2	114.5	544
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	33.0	33
MAY RETAINED	235.2	0.7	0.3	12.7	25.0	7.0	106.7	4.7	0	20.3	0.5	2.7	13.2	147.5	577
SE	53.0	0.3	0.2	4.2	8.5	2.6	21.5	3.0	0	5.9	0.3	1.5	6.8	22.9	63
June															
Creel Estimate	483.2	13.8	11.1	47.0	63.2	58.2	360.3	0	0	66.2	4.2	0	54.4	386.9	1,548
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	197.0	197
JUNE RETAINED	483.2	13.8	11.1	47.0	63.2	58.2	360.3	0	0	66.2	4.2	0	54.4	583.9	1,745
SE	62.3	3.9	3.6	7.3	10.6	19.2	36.0	0	0	9.7	2.4	0	10.2	50.5	92
July															
Creel Estimate	1199.7	34.9	29.7	124.6	120.2	21.4	730.4	0	0	167.8	5.7	0	155.7	461.0	3,051
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	258.0	258
JULY RETAINED	1199.7	34.9	29.7	124.6	120.2	21.4	730.4	0	0	167.8	5.7	0	155.7	719.0	3,309
SE	124.6	10.3	9.4	18.7	22.8	7.8	52.0	0	0	19.3	3.1	0	23.1	71.0	159
August															
Creel Estimate	1180.9	37.7	24.2	103.9	114.5	32.9	794.5	0	0	174.0	9.5	0	243.8	460.9	3,177
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	299	299
AUGUST RETAINED	1180.9	37.7	24.2	103.9	114.5	32.9	794.5	0	0	174.0	9.5	0	243.8	759.9	3,476
SE	116.0	14.6	7.3	16.2	20.1	13.4	58.1	0	0	24.0	5.5	0	39.4	90.2	168
TOTAL															
Creel Estimate	3,099	87	65	288	323	120	1,992	5	0	428	20	3	467	1,423	8,320
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	787	787
TOTAL RETAINED	3,099	87	65	288	323	120	1,992	5	0	428	20	3	467	2,210	9,107
SE	189	18	12	26	33	25	89	3	0	33	7	2	47	128	257

Estimated numbers of rockfish released by lodge and non-lodge anglers, by taxon, 2025.

	Taxon														
Month / Catch Type	Black Rockfish	Boccacio Rockfish	Canary Rockfish	China Rockfish	Copper Rockfish	Dusky Rockfish	Quillback Rockfish	Redbande d Rockfish	Redstripe Rockfish	Tiger Rockfish	Widow Rockfish	Yelloweye Rockfish	Yellowtail Rockfish	Unknown rockfish	TOTAL
May															
Creel Estimate	84.4	0	0.3	14.8	11.8	0.8	71.3	0	0.5	3.3	0	78.5	19.7	251.8	537
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MAY RELEASED	84.4	0	0.3	14.8	11.8	0.8	71.3	0	0.5	3.3	0	78.5	19.7	251.8	537
SE	31.9	0	0.2	7.7	2.1	0.5	15.0	0	0.4	2.2	0	17.5	9.9	44.7	61
June															
Creel Estimate	105.0	0	2.4	53.8	42.3	6.6	144.1	0	4.0	9.5	0	224.6	22.5	768.8	1,384
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JUNE RELEASED	105.0	0	2.4	53.8	42.3	6.6	144.1	0	4.0	9.5	0	224.6	22.5	768.8	1,384
SE	24.0	0	1.6	12.8	11.8	3.7	23.6	0	3.1	5.0	0	35.0	6.9	99.1	112
July															
Creel Estimate	188.2	2.3	2.2	126.2	24.1	2.0	284.5	0	0	10.7	0	657.3	64.3	1533.7	2,896
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
JULY RELEASED	188.2	2.3	2.2	126.2	24.1	2.0	284.5	0	0	10.7	0	657.3	64.3	1533.7	2,896
SE	53.4	1.8	1.3	27.6	7.3	1.6	42.3	0	0	5.2	0	95.8	18.0	206.4	240
August															
Creel Estimate	118.8	4.9	2.4	75.2	19.6	0	336.5	0	0	14.1	0	751.7	59.7	1694.8	3,078
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
AUGUST RELEASED	118.8	4.9	2.4	75.2	19.6	0	336.5	0	0	14.1	0	751.7	59.7	1694.8	3,078
SE	26.2	3.5	1.8	21.7	7.5	0	51.1	0	0	5.6	0	122.8	14.7	191.0	236
TOTAL															
Creel Estimate	496	7	7	270	98	9	836	0	5	38	0	1,712	166	4,249	7,894
Lodge Report	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
TOTAL RELEASED	496	7	7	270	98	9	836	0	5	38	0	1,712	166	4,249	7,894
SE	72	4	3	38	16	4	72	0	3	9	0	161	26	301	360

APPENDIX 5: STATISTICAL EFFECTS OF PERIOD ON HARVEST, BY SPECIES.

Kruskal Wallis tests for differences in harvest (fish caught per boat trip per subarea) among periods, by species, 2025. Subareas treated as replicates.
CLD: red letters indicate statistical differences within species (periods not connected by a common letter were statistically different).

Species	Kruskal-Wallis Statistics			Median Harvest by Period					CLD
	χ^2	df	P	1-28 May	29 May - 22 Jun	23 Jun - 23 Jul	24 Jul - 17 Aug	18-31 Aug	
<i>Chinook Salmon</i>	27.4	4	< 0.0001	23.1	98.5	0	186.1	32.7	A-A-B-A-A
<i>Chum Salmon</i>	4.0	4	0.41	0	0	0	0	0	
<i>Coho Salmon</i>	35.5	4	< 0.0001	0	6.7	527.0	1244.3	73.6	D-C-A-A-B
<i>Pink Salmon</i>	32.5	4	< 0.0001	0	0	57.5	93.6	6.4	C-C-AB-A-B
<i>Sockeye Salmon</i>	17.5	4	0.002	0	0	0	1.0	0	A-A-A-A-A
<i>Arrow-tooth Flounder</i>	6.3	4	0.18	0	0	0	0	0	
<i>Big Skate</i>	4.0	4	0.41	0	0	0	0	0	
<i>Cabezon</i>	0.5	4	0.97	0	0	0	0	0	
<i>English Sole</i>	4.0	4	0.41	0	0	0	0	0	
<i>Greenlings</i>	4.9	4	0.30	0	0	0.7	0.9	0	
<i>Lingcod</i>	11.8	4	0.019	23.9	35.2	79.2	112.7	45.7	B-AB-AB-A-AB
<i>Longnose Skate</i>	0.0	4	1	0	0	0	0	0	
<i>Pacific Cod</i>	9.7	4	0.046	2.7	4.6	50.6	71.2	14.7	A-A-A-A-A
<i>Pacific Halibut</i>	17.3	4	0.002	30.4	98.0	211.3	233.8	63.9	B-AB-A-A-AB
<i>Pacific Rock Sole</i>	9.2	4	0.056	0	0	0	0	0	
<i>Pacific Spiny Dogfish</i>	3.7	4	0.45	0	0	0	0	0	
<i>Sablefish</i>	3.1	4	0.55	0	0	0	0	0	
<i>Spotted Ratfish</i>	0.0	4	1	0	0	0	0	0	
<i>Starry Flounder</i>	0.0	4	1	0	0	0	0	0	
<i>Black Rockfish</i>	6.1	4	0.19	9.2	6.2	35.2	38.8	13.1	
<i>Boccacio Rockfish</i>	6.7	4	0.15	0	0	0	0	0	
<i>Canary Rockfish</i>	8.8	4	0.07	0	0	0	0	0	
<i>China Rockfish</i>	3.7	4	0.45	0	0	2.9	0.9	0	
<i>Copper Rockfish</i>	1.7	4	0.79	0.02	2.2	0.7	2.7	0	
<i>Dusky Rockfish</i>	6.6	4	0.16	0	0	0	0	0	
<i>Quillback Rockfish</i>	15.5	4	0.004	0.4	6.5	26.9	54.9	12.6	B-AB-AB-A-AB
<i>Redbanded Rockfish</i>	8.1	4	0.09	0	0	0	0	0	
<i>Redstripe Rockfish</i>	0.0	4	1	0	0	0	0	0	
<i>Tiger Rockfish</i>	7.6	4	0.11	0	0	0	6.1	0.2	
<i>Widow Rockfish</i>	3.7	4	0.45	0	0	0	0	0	
<i>Yelloweye Rockfish</i>	8.1	4	0.09	0	0	0	0	0	
<i>Yellowtail Rockfish</i>	12.3	4	0.015	0	0	7.7	9.2	3.6	B-AB-A-A-A

APPENDIX 6: ADIPOSE FIN-CLIP INCIDENCE

Incidence of fin clipping observed in biosamples, by species, month, and subarea, in 2025. Excludes any interviews for which observers were not able to confirm adipose status, or for which biosampling was not a complete sample of fish kept.

Species		Subarea														TOTAL
Month	Adipose	3I	3J	3K	3?	4A	4B	4C	4D	4E	4F	4G	4H	4L	4?	
Chinook Salmon																
May	Clipped	0	0	0	0	0	2	2	4	0	8	0	0	0	0	16
	Not Clipped	0	0	3	0	3	18	14	13	2	17	0	0	0	0	70
	Percent Clipped	-	-	0%	-	0%	10%	13%	24%	0%	32%	-	-	-	-	19%
June	Clipped	0	1	0	0	0	1	3	5	0	4	0	0	0	0	14
	Not Clipped	1	2	0	0	0	17	12	13	0	24	0	2	0	0	71
	Percent Clipped	0%	33%	-	-	-	6%	20%	28%	-	14%	-	0%	-	-	16%
July	Clipped	0	0	0	0	0	1	0	1	0	0	0	0	0	0	2
	Not Clipped	0	0	0	0	0	0	3	2	0	1	0	0	0	0	6
	Percent Clipped	-	-	-	-	-	100%	0%	33%	-	0%	-	-	-	-	25%
August	Clipped	0	0	0	0	0	0	0	0	0	2	0	0	0	0	2
	Not Clipped	1	1	0	0	0	0	4	1	0	9	1	0	0	0	17
	Percent Clipped	0%	0%	-	-	-	-	0%	0%	-	18%	0%	-	-	-	11%
Total	Clipped	0	1	0	0	0	4	5	10	0	14	0	0	0	0	34
	Not Clipped	2	3	3	0	3	35	33	29	2	51	1	2	0	0	164
	Percent Clipped	0%	25%	0%	-	0%	10%	13%	26%	0%	22%	0%	0%	-	-	17.2%
Coho Salmon																
May	Clipped	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Not Clipped	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1
	Percent Clipped	-	-	-	-	-	-	-	0%	-	-	-	-	-	-	0.0%
June	Clipped	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Not Clipped	0	6	5	0	0	0	0	2	0	2	0	0	0	0	15
	Percent Clipped	-	0%	0%	-	-	-	-	0%	-	0%	-	-	-	-	0.0%
July	Clipped	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Not Clipped	7	4	3	0	0	4	43	21	0	5	0	0	0	0	87
	Percent Clipped	0%	0%	0%	-	-	0%	0%	0%	-	0%	-	-	-	-	0.0%
August	Clipped	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Not Clipped	11	1	9	0	0	7	5	12	0	12	5	0	0	0	62
	Percent Clipped	0%	0%	0%	-	-	0%	0%	0%	-	0%	0%	-	-	-	0.0%
Total	Clipped	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	Not Clipped	18	11	17	0	0	11	48	36	0	19	5	0	0	0	165
	Percent Clipped	0%	0%	0%	-	-	0%	0%	0%	-	0%	0%	-	-	-	0.0%

APPENDIX 7: BIOSAMPLING RESULTS, MAY 2025

	<i>Chinook Salmon</i>		<i>Coho Salmon</i>		<i>Pacific Halibut</i>	
	<i>Number</i>	<i>Percent</i>	<i>Number</i>	<i>Percent</i>	<i>Number</i>	<i>Percent</i>
Number Examined	131		1		53	
Subarea						
3I						
3J	1	1%			3	6%
3K	3	2%				
4A	4	3%				
4B	29	22%			2	4%
4C	20	15%			12	24%
4D	33	25%	1	100%	7	14%
4E	2	2%				
4F	39	30%			22	44%
4G						
4H					4	8%
4L						
Sex						
Male	1	100%				
Female						
Undeterminable						
Unknown	130		1		53	
Trip Type						
Guided	54	41%	1	100%	30	57%
Unguided	77	59%			23	43%
Adipose Clips						
Clipped	28	22%				
Unmarked	101	78%	1	100%		
Unknown						
Head Tagged						
Yes	23	18%				
No	108	82%	1	100%		
Scale Samples			Whatman Samples			
Yes	129	98%	1	100%		
No	2	2%				
Flesh Colour						
Red	17	89%				
White	2	11%				
Marble						
Unknown						