## Subject Index

| abc method, 194, 204 Accelerated gradient descent, 359 Acceleration, 192, 205 Accuracy, 14 after model selection, 402–408 Accurate but not correct, 402 Activation function, 355, 361 leaky rectified linear, 362 rectified linear, 362 ReLU, 362 tanh, 362 Active set, 302, 308 adaboost algorithm, 341–345, 447 Adaboost.M1, 342 Adaptation, 404 Adaptive rate control, 359 Additive model, 324 adaptive, 346 Adjusted compliance, 404 Admixture modeling, 256–260 AIC, see Akaike information criterion Akaike information criterion, 208, 218, 226, 231, 246, 267 Allele frequency, 257 American Statistical Association, 449 Ancillary, 44, 46, 139 Apparent error, 211, 213, 219 arcsin transformation, 95 Arthur Eddington, 447 Asymptotics, xvi, 119, 120 Autoencoder, 362–364 Backfitting, 346 Backpropagation, 356–358 Bagged estimate, 404, 406 Bagging, 226, 327, 406, 408, 419 Balance equations, 256 Barycentric plot, 259 | Basis expansion, 375 Bayes deconvolution, 421–424 factor, 244, 285 false-discovery rate, 279 posterior distribution, 254 posterior probability, 280 shrinkage, 212 t-statistic, 255 theorem, 22 Bayes-frequentist estimation, 412–417 Bayesian inference, 22–37 information criterion, 246 lasso, 420 lasso prior, 415 model selection, 244 trees, 349 Bayesian information criterion, 267 Bayesianism, 3 BCa accuracy and correctness, 206 confidence density, 202, 207, 237, 242, 243 interval, 202 method, 192 Benjamini and Hochberg, 276 Benjamini–Yekutieli, 400 Bernoulli, 338 Best-approximating linear subspace, 363 Best-subset selection, 299 Beta distribution, 54, 239 BH <sub>q</sub> , 276 Bias, 14, 352 Bias-corrected, 330 and accelerated, see BCa method confidence intervals, 190–191 percentile method, 190 |
|--|--|
|--|--|

| Bias-correction value, 191              | Coherent behavior, 261                     |
|---|--|
| Biased estimation, 321                  | Common task framework, 447                 |
| BIC, see Bayesian information criterion | Compliance, 394                            |
| Big-data era, xv, 446                   | Computational bottleneck, 128              |
| Binomial, 109, 117                      | Computer age, xv                           |
|   |  |
| distribution, 54, 117, 239              | Computer-intensive, 127                    |
| log-likelihood, 380                     | inference, 189, 267                        |
| standard deviation, 111                 | statistics, 159                            |
| Bioassay, 109                           | Conditional, 58                            |
| Biometrika, 449                         | Conditional distribution                   |
| Bivariate normal, 182                   | full, 253                                  |
| Bonferroni bound, 273                   | Conditional inference, 45–48, 139, 142     |
| Boole's inequality, 274                 | lasso, 318                                 |
| Boosting, 320, 324, 333–350             | Conditionality, 44                         |
| Bootstrap, 7, 155–180, 266, 327         | Confidence                                 |
| Baron Munchausen, 177                   | density, 200, 201, 235                     |
| Bayesian, 168, 179                      | distribution, 198–203                      |
| cdf, 187                                | interval, 17                               |
| confidence intervals, 181–207           | region, 397                                |
| ideal estimate, 160, 179                | Conjugate, 253, 259                        |
| jackknife after, 179                    | prior, 238                                 |
| moving blocks, 168                      | priors, 237                                |
|   |  |
| multisample, 167                        | Convex optimization, 304, 308, 321, 323,   |
| nonparametric, 159–162, 217             | 377  |
| out of bootstrap, 232                   | Convolution, 422, 445                      |
| packages, 178                           | filters, 368                               |
| parametric, 169–173, 223, 312, 429      | layer, 367                                 |
| probabilities, 164                      | Corrected differences, 411                 |
| replication, 159                        | Correlation effects, 295                   |
| sample, 159                             | Covariance                                 |
| sample size, 179, 205                   | formula, 312                               |
| smoothing, 226, 404, 406                | penalty, 218–226                           |
| t, 196                                  | Coverage, 181                              |
| t intervals, 195–198                    | Coverage level, 274                        |
| Bound form, 305                         | Coverage matching prior, 236–237           |
| Bounding hyperplane, 398                | Cox model, see proportional hazards        |
| Burn-in, 260                            | model                                      |
| $BY_q$ algorithm, 400                   | $C_p$ , 217, 218, 221, 231, 267, 300, 394, |
| Causal inference, xvi                   | 395, 403                                   |
|   | Cramér–Rao lower bound, 44                 |
| Censored                                | Credible interval, 198, 417                |
| data, 134–139                           | Cross-validation, 208–232, 267, 335        |
| not truncated, 150                      | 10-fold, 326                               |
| Centering, 107                          | estimate, 214                              |
| Central limit theorem, 119              | K-fold, 300                                |
| Chain rule for differentiation, 356     | leave one out, 214, 231                    |
| Classic statistical inference, 3–73     |  |
| Classification, 124, 209                | Cumulant generating function, 67           |
| Classification accuracy, 375            | Curse of dimensionality, 387               |
| Classification error, 209               | Dark energy, 210, 231                      |
| Classification tree, 348                | Data analysis, 450                         |
| Cochran-Mantel-Haenszel test, 131       | Data science, xvii, 450, 451               |
|   |  |

| Data sets   | binomial, 54, 117, 239                        |
|---|---|
| ALS, 334  | gamma, 54, 117, 239                           |
| AML, see leukemia                                       | Gaussian, 54                                  |
| baseball, 94  | normal, 54, 117, 239                          |
| butterfly, 78   | Poisson, 54, 117, 239                         |
| cell infusion, 112                                      | Divide-and-conquer algorithm, 325             |
| cholesterol, 395, 402, 403                              | Document retrieval, 298                       |
| CIFAR-100, 365  | Dose–response, 109                            |
| diabetes, 98, 209, 396, 414, 416                        | Dropout learning, 368, 372                    |
| dose-response, 109                                      | DTI, see diffusion tensor imaging             |
| galaxy, 120   | Early computer-age, xvi, 75–268               |
| handwritten digits                                      | Early stopping, 362                           |
| (MNIST), 353  | Effect size, 272, 288, 399, 408               |
| head/neck cancer, 135                                   | Efficiency, 44, 120                           |
| human ancestry, 257                                     | Eigenratio, 162, 173, 194                     |
| insurance, 131  | Elastic net, 316, 356                         |
| kidney function, 157, 222                               | Ellipsoid, 398                                |
| leukemia, 176, 196, 377                                 | EM algorithm, 146–150                         |
| NCOG, 134   | missing data, 266                             |
| nodes, 424, 427, 430, 438, 439, 442                     | Empirical Bayes, 75–90, 93, 264               |
| pediatric cancer, 143                                   | estimation strategies, 421–445                |
| police, 287   | information, 443                              |
| prostate, 249, 272, 289, 408, 410,                      | large-scale testing, 278–282                  |
| 423, 434–436  | Empirical null, 286                           |
| protein classification, 385                             | estimation, 289–290                           |
| shakespear, 81  | maximum-likelihood estimation, 296            |
| <b>spam</b> , 113, 127, 209, 215, 300–302,              | Empirical probability distribution, 160       |
| 325   | Ensemble, 324, 334                            |
| student score, 173, 181, 186,                           | Ephemeral predictors, 227                     |
| 202, 203  | Epoch, 359                                    |
| supernova, 210, 212, 217, 221, 224                      | Equilibrium distribution, 256                 |
| vasoconstriction, 240, 241,                             | Equivariant, 106                              |
| 246, 252  | Exact inferences, 119                         |
| Data snooping, 398                                      | Expectation parameter, 118                    |
| De Finetti, B., 35, 36, 251, 450                        | Experimental design, xvi                      |
| De Finetti–Savage school, 251                           | Exponential family, 53–72, 225                |
| Debias, 318   | <i>p</i> -parameter, 117, 413, 424 curved, 69 |
| Decision rule, 275                                      |   |
| Decision theory, xvi                                    | one-parameter, 116                            |
| Deconvolution, 422                                      | F distribution, 397                           |
| Deep learning, 351–374                                  | F tests, 394                                  |
| Definitional bias, 431                                  | f-modeling, 424, 434, 440–444                 |
| Degrees of freedom, 221, 231, 312–313                   | Fake-data principle, 148, 154, 266            |
| Delta method, 15, 414, 420                              | False coverage                                |
| Deviance, 112, 118, 119, 301                            | control, 399                                  |
| Deviance residual, 123<br>Diffusion tensor imaging, 291 | False discovery, 275                          |
| Direct evidence, 105, 109, 421                          | control, 399                                  |
| Directional derivatives, 158                            | control theorem, 294 proportion, 275          |
| Distribution  | rate, 271–297                                 |
| beta, 54, 239   | False-discovery                               |
| oom, 5 1, 257   | Tuise discovery                               |

## Subject Index

| rate, 9 Family of probability densities, 64 Family-wise error rate, 274 FDR, see false-discovery rate Feed-forward, 351  | Google flu trends, 230, 232<br>Gradient boosting, 338–341<br>Gradient descent, 354, 356<br>Gram matrix, 381<br>Gram-Schmidt orthogonalization, 322  |
|--|---|
| Fiducial, 267 constructions, 199 density, 200 inference, 51  | Graphical lasso, 321<br>Graphical models, xvi<br>Greenwood's formula, 137, 151<br>Group lasso, 321  |
| Fisher, 79 Fisher information, 29, 41, 59 bound, 41 matrix, 236, 427 Fisherian correctness, 205 Fisherian inference, 38–52, 235 Fixed-knot regression splines, 345 Flat prior, 235 Forward pass, 357 | Hadamard product, 358 Handwritten digits, 353 Haplotype estimation, 261 Hazard rate, 131–134 parametric estimate, 138 Hidden layer, 351, 352, 354 High-order interaction, 325 Hinge loss, 380 |
| Forward pass, 337 Forward-stagewise, 346 fitting, 320 Forward-stepwise, 298–303 computations, 322 logistic regression, 322 regression, 300   | Hints learning with, 369 Hoeffding's lemma, 118 Holm's procedure, 274, 294 Homotopy path, 306 Hypergeometric distribution, 141, 152   |
| Fourier method, 440 transform, 440 Frailties, 439  | Imputation, 149<br>Inadmissable, 93<br>Indirect evidence, 102, 109, 266, 290,<br>421, 440, 443  |
| Frequentism, 3, 12–22, 30, 35, 51, 146, 267  | Inductive inference, 120<br>Inference, 3  |
| Frequentist, 413<br>inference, 12–21<br>strongly, 218  | Inference after model selection, 394–420<br>Inferential triangle, 446<br>Infinitesimal forward stagewise, 320   |
| Fully connected layer, 368 Functional gradient descent, 340 FWER, <i>see</i> family-wise error rate  | Infinitesimal jackknife, 167 estimate, 406 standard deviations, 407 Influence function, 174, 177  |
| g-modeling, 423<br>Gamma, 117<br>distribution, 54, 117, 239  | Influence function, 174–177<br>empirical, 175<br>Influenza outbreaks, 230<br>Input distortion, 369, 373   |
| General estimating equations, xvi<br>General information criterion, 248<br>Generalized<br>linear mixed model, 437–440<br>linear model, 108–123, 266<br>ridge problem, 384                            | Input layer, 355 Insample error, 219 Inverse chi-squared, 262 Inverse gamma, 239, 262 IRLS, <i>see</i> iteratively reweighted least squares   |
| Genome, 257 Genome-wide association studies, 451 Gibbs sampling, 251–260, 267, 414 GLM, <i>see</i> generalized linear model GLMM, <i>see</i> generalized linear mixed model                          | Iteratively reweighted least squares, 301, 322  Jackknife, 155–180, 266, 330 estimate of standard error, 156 standard error, 178  |
|  |   |

| James-Stein                             | Local false-discovery rate, 280, 282–286 |
|---|--|
| estimation, 91-107, 282, 305, 410       | Local regression, 387–390, 393           |
| ridge regression, 265                   | Local translation invariance, 368        |
| Jeffreys                                | Log polynomial regression, 410           |
| prior, 237                              | Log-rank statistic, 152                  |
| Jeffreys'                               | Log-rank test, 131, 139–142, 152, 266    |
| prior, 28–30, 36, 198, 203, 236         | Logic of inductive inference, 185, 205   |
| prior, multiparameter, 242              | Logistic regression, 109–115, 139, 214,  |
| scale, 285                              | 299, 375                                 |
| Jumpiness of estimator, 405             | multiclass, 355                          |
|   | Logit, 109                               |
| Kaplan–Meier, 131, 134, 136, 137        |  |
| estimate, 134–139, 266                  | Loss plus penalty, 385                   |
| Karush–Kuhn–Tucker optimality           | Machine learning, 208, 267, 375          |
| conditions, 308                         | Mallows' $C_p$ , see $C_p$               |
| Kernel                                  | Mantel-Haenzel test, 131                 |
| function, 382                           | MAP, 101                                 |
| logistic regression, 386                | MAP estimate, 420                        |
| method, 375–393                         | Margin, 376                              |
| SVM, 386                                | Marginal density, 409, 422               |
| trick, 375, 381–383, 392                | Markov chain Monte Carlo, see MCMC       |
| Kernel smoothing, 375, 387–390          | Markov chain theory, 256                 |
| Knots, 309                              | Martingale theory, 294                   |
| Kullback-Leibler distance, 112          | Matching prior, 198, 200                 |
|   | Matlab, 271                              |
| $\ell_1$ regularization, 321            | Matrix completion, 321                   |
| Lagrange<br>dual, 381                   | Max pool layer, 366                      |
|   | Maximized a-posteriori probability, see  |
| form, 305, 308                          | MAP                                      |
| multiplier, 391                         | Maximum likelihood, 299                  |
| Large-scale                             | Maximum likelihood estimation, 38–52     |
| hypothesis testing, 271–297             |  |
| testing, 272–275                        | MCMC, 234, 251–260, 267, 414             |
| Large-scale prediction algorithms, 446  | McNemar test, 341                        |
| Lasso, 101, 210, 217, 222, 231, 298–323 | Mean absolute deviation, 447             |
| modification, 312                       | Median unbiased, 190                     |
| path, 312                               | Memory-based methods, 390                |
| penalty, 356                            | Meter reader, 30                         |
| Learning from the experience of others, | Meter-reader, 37                         |
| 104, 280, 290, 421, 443                 | Microarrays, 227, 271                    |
| Learning rate, 358                      | Minitab, 271                             |
| Least squares, 98, 112, 299             | Misclassification error, 302             |
| Least-angle regression, 309–313, 321    | Missing data, 146–150, 325               |
| Least-favorable family, 262             | EM algorithm, 266                        |
| Left-truncated, 150                     | Missing-species problem, 78–84           |
| Lehmann alternative, 294                | Mixed features, 325                      |
| Life table, 131–134                     | Mixture density, 279                     |
| Likelihood function, 38                 | Model averaging, 408                     |
| concavity, 118                          | Model selection, 243–250, 398            |
| Limited-translation rule, 293           | criteria, 250                            |
| Lindsey's method, 68                    | Monotone lasso, 320                      |
| Linearly separable, 375                 | Monotonic increasing function, 184       |
| Link function, 237, 340                 | Multinomial                              |
|   |  |

| distribution, 61-64, 425               | predictor, 221                          |
|--|---|
| from Poisson, 63                       | One-sample nonparametric bootstrap,     |
| Multiple testing, 272                  | 161                                     |
| Multivariate                           | One-sample problems, 156                |
| analysis, 119                          | OOB, see out-of-bag error               |
| normal, 55–59                          | Optical character recognition, 353      |
| <i>n</i> -gram, 385                    | Optimal separating hyperplane, 375–377  |
| N-P complete, 299                      | Optimal-margin classifier, 376          |
| Nadaraya–Watson estimator, 388         | Optimality, 18                          |
| Natural parameter, 116                 | Oracle, 275                             |
| Natural spline model, 430              | Orthogonal parameters, 262              |
| NCOG, see Northern California          | Out-of-bag error, 232, 327, 329–330     |
| Oncology Group                         | Out-the-box learning algorithm, 324     |
| Nested models, 299                     | Output layer, 352                       |
| Neural Information Processing Systems, | Outsample error, 219                    |
| 372                                    | Over parametrized, 298                  |
| Neural network, 351–374                | Overfitting, 304                        |
| adaptive tuning, 360                   | Overshrinks, 97                         |
| number of hidden layers, 361           | <i>p</i> -value, 9, 282                 |
| Neurons, 351                           | Package/program                         |
| Neyman's construction, 181, 183, 193,  | gbm, 335, 348                           |
| 204                                    | glmnet, 214, 315, 322, 348              |
| Neyman-Pearson, 18, 19, 293            | h2o, 372                                |
| Non-null, 272                          | lars, 312, 320                          |
| Noncentral chi-square variable, 207    | liblineaR, 381                          |
| Nonlinear transformations, 375         | locfdr, 289-291, 296, 437               |
| Nonlinearity, 361                      | lowess, 6, 222, 388                     |
| Nonparameteric                         | nlm, 428                                |
| regression, 375                        | randomForest, 327, 348                  |
| Nonparametric, 53, 127                 | selectiveInference, 323                 |
| MLE, 150, 160                          | Pairwise inner products, 381            |
| percentile interval, 187               | Parameter space, 22, 29, 54, 62, 66     |
| Normal                                 | Parametric bootstrap, 242               |
| correlation coefficient, 182           | Parametric family, 169                  |
| distribution, 54, 117, 239             | Parametric models, 53–72                |
| multivariate, 55–59                    | Partial likelihood, 142, 145, 151, 153, |
| regression model, 414                  | 266, 341                                |
| theory, 119                            | Partial logistic regression, 152        |
| Northern California Oncology Group,    | Partial residual, 346                   |
| 134                                    | Path-wise coordinate descent, 314       |
| Nuclear norm, 321                      | Penalized                               |
| Nuisance parameters, 142, 199          | least squares, 101                      |
| Objective Bayes, 36, 267               | likelihood, 101, 428                    |
| inference, 233–263                     | logistic regression, 356                |
| intervals, 198–203                     | maximum likelihood, 226, 307            |
| prior distribution, 234–237            | Percentile method, 185–190              |
| OCR, see optical character recognition | central interval, 187                   |
| Offset, 349                            | Permutation null, 289, 296              |
| OLS                                    | Permutation test, 49–51                 |
| algorithm, 403                         | Phylogenetic tree, 261                  |
| estimation, 395                        | Piecewise                               |
|  |   |

| linear, 313                            | sampling variance, 330   |
|--|--|
| nonlinear, 314                         | standard error, 330–331  |
| Pivotal                                | Randomization, 49–51   |
| argument, 183                          | Rao-Blackwell, 227, 231  |
| quantity, 196, 198                     | Rate annealing, 360  |
| statistic, 16                          | Rectified linear, 359  |
| .632 rule, 232                         | Regression, 109  |
| Poisson, 117, 193                      | Regression rule, 219   |
| distribution, 54, 117, 239             | Regression to the mean, 33                                       |
| regression, 120–123, 249, 284, 295,    | Regression tree, 124–128, 266, 348                               |
| 435                                    | Regularization, 101, 173, 298, 379, 428                          |
| Poisson regression, 171                | path, 306  |
| Polynomial kernel, 382, 392            | Relevance, 290–293   |
| Positive-definite function, 382        | Relevance function, 293  |
| Post-selection inference, 317, 394–420 | Relevance theory, 297  |
| Posterior density, 235, 238            | Reproducing kernel Hilbert space, 375,                           |
| Posterior distribution, 416            | 384, 392   |
| Postwar era, 264                       | Resampling, 162  |
| Prediction                             | plans, 162–169   |
| errors, 216                            | simplex, 164, 169  |
| rule, 208–213                          | vector, 163  |
| Predictors, 124, 208                   | Residual deviance, 283   |
| Principal components, 362              | Response, 124, 208   |
| Prior distribution, 234–243            | Ridge regression, 97–102, 209, 304, 327,                         |
| beta, 239                              | 332, 372, 381  |
| conjugate, 237–243                     | James–Stein, 265   |
| coverage matching, 236–237 gamma, 239  | Ridge regularization, 368  |
| normal, 239                            | logistic regression, 392   |
| objective Bayes, 234                   | Right-censored, 150  |
| proper, 239                            | Risk set, 144  |
| Probit analysis, 112, 120, 128         | RKHS, see reproducing-kernel Hilbert                             |
| Propagation of errors, 420             | Space Pobbins' formula 75, 77, 422, 440                          |
| Proper prior, 239                      | Robbins' formula, 75, 77, 422, 440<br>Robust estimation, 174–177 |
| Proportional hazards model, 131,       | Royal Statistical Society, 449                                   |
| 142–146, 266                           |  |
| Proximal-Newton, 315                   | S language, 271  |
| <i>q</i> -value, 280                   | Sample correlation coefficient, 182                              |
| QQ plot, 287                           | Sample size coherency, 248                                       |
| QR decomposition, 311, 322             | Sampling distribution, 312                                       |
| Quadratic program, 377                 | SAS, 271   |
| Quasilikelihood, 266                   | Savage, L. J., 35, 36, 51, 199, 233, 251, 450                    |
| Quetelet, Adolphe, 449                 | Scale of evidence  |
| R, 178, 271                            | Fisher, 245  |
| Random forest, 209, 229, 324–332,      | Jeffreys, 245  |
| 347–350                                | Scheffé  |
| adaptive nearest-neighbor estimator,   | interval, 396, 397, 417  |
| 328                                    | theorem, 398   |
| leave-one-out cross-validated error,   | Score function, 42   |
| 329                                    | Score tests, 301   |
| Monte Carlo variance, 330              | Second-order accuracy, 192–195                                   |
| ,                                      | ,  |

## Subject Index

| Selection bias, 33, 408–411            | Studentized range, 418                   |
|--|--|
| Self-consistent, 149                   | Subgradient                              |
| Separating hyperplane, 375             | condition, 308                           |
| geometry, 390                          | equation, 312, 315                       |
| Seven-league boots, 448                | Subjective prior distribution, 233       |
| Shrinkage, 115, 316, 338               | Subjective probability, 233              |
| estimator, 59, 91, 94, 96, 410         | Subjectivism, 35, 233, 243, 261          |
| Sigmoid function, 352                  | Sufficiency, 44                          |
| Significance level, 274                | Sufficient                               |
| Simulation, 155–207                    | statistic, 66, 112, 116                  |
| Simultaneous confidence intervals,     | vector, 66                               |
| 395–399                                | Supervised learning, 352                 |
| Simultaneous inference, 294, 418       | Support                                  |
| Sinc kernel, 440, 445                  | set, 377, 378                            |
| Single-nucleotide polymorphism, see    | vector, 377                              |
| SNP                                    | vector classifiers, 381                  |
| Smoothing operator, 346                | vector machine, 319, 375-393             |
| SNP, 257                               | SURE, see Stein's unbiased risk estimate |
| Soft margin classifier, 378–379        | Survival analysis, 131–154, 266          |
| Soft-threshold, 315                    | Survival curve, 137, 279                 |
| Softmax, 355                           | SVM                                      |
| Spam filter, 115                       | Lagrange dual, 391                       |
| Sparse                                 | Lagrange primal, 391                     |
| models, 298–323                        | loss function, 391                       |
| principal components, 321              | Taylor series, 157, 420                  |
| Sparse matrix, 316                     | Theoretical null, 286                    |
| Sparsity, 321                          | Tied weights, 368                        |
| Split-variable randomization, 327, 332 | Time series, xvi                         |
| SPSS, 271                              | Training set, 208                        |
| Squared error, 209                     | Transformation invariance, 183–185, 236  |
| Standard candles, 210, 231             | Transient episodes, 228                  |
| Standard error, 155                    | Trees                                    |
| external, 408                          | averaging, 348                           |
| internal, 408                          | best-first, 333                          |
| Standard interval, 181                 | depth, 335                               |
| Stein's                                | terminal node, 126                       |
| paradox, 105                           | Tricube kernel, 388, 389                 |
| unbiased risk estimate, 218, 231       | Trimmed mean, 175                        |
| Stepwise selection, 299                | Triple-point, xv                         |
| Stochastic gradient descent, 358       | True error rate, 210                     |
| Stopping rule, 32, 413                 | True-discovery rates, 286                |
| Stopping rules, 243                    | Tukey, J. W., 418, 450                   |
| String kernel, 385, 386                | Tukey, J. W., 418                        |
| Strong rules, 316, 322                 | Tweedie's formula, 409, 419, 440         |
| Structure, 261                         | Twenty-first-century methods, xvi,       |
| Structure matrix, 97, 424              | 271–446                                  |
| Student t                              | Two-groups model, 278                    |
| confidence interval, 396               | Uncorrected differences, 411             |
| distribution, 196, 272                 | Uninformative prior, 28, 169, 233, 261   |
| statistic, 449                         | Universal approximator, 351              |
| two-sample, 8, 272                     | Unlabeled images, 365                    |
|  |  |

Unobserved covariates, 288

Validation set, 213

Vapnik, V., 390

Variable-importance plot, 331–332, 336

Variance, 14

Variance reduction, 324

Velocity vector, 360

Voting, 333

Warm starts, 314, 363

Weak learner, 333, 342

Weight

decay, 356 regularization, 361, 362 sharing, 352, 367 Weighted exponential loss, 345 Weighted least squares, 315 Weighted majority vote, 341 Weights, 352 Wide data, 298, 321 Wilks' likelihood ratio statistic, 246 Winner's curse, 33, 408 Winsorized mean, 175 Working response, 315, 322  $z^{(\alpha)}$ , 188

 $z^{(\alpha)}$ , 188 Zero set, 296