

MRI REPORT –LEFT ANKLE JOINT

TECHNIQUE:

PD FS Axial, coronal

STIR sagittal, T1W Axial, T2 coronal

OBSERVATION:

High degree partial tear in ATFL with diffuse interstitial edema.

Suspicious partial tear in calcaneofibular ligament.

Mild to moderate sprain PTFL

Subtle synovial thickening along the lateral retinaculum of the ankle joint

Mild ankle joint effusion with periarticular subcutaneous mild edema.

Mild tenosynovitis in tibialis posterior tendon.

Small bony spur in anterosuperior aspect of the neck of talus and anterior aspect of the tibia—of concern for anterior ankle joint impingement. few tiny loose calcific bony fragments in the anterior tibiotalar joint space.

Small loose calcific body in the posterolateral aspect of the subtalar joint, measures approximately 5.8 mm.

The lower end of the tibia and fibula appear normal. There is no evidence of bone marrow edema / focal lesion.

The plantar fascia appears normal.

There is no evidence of tendonitis or tendon tear in any of the tendons. No fluid collection is seen in rest of the tendon sheaths.

The Achilles appears normal.

The retro calcaneal bursa and the fat in Kager's triangle appear normal.

The calcaneum, talus, navicular, cuboid and the cuneiform appear normal.