



DEPARTMENT OF RADIOLOGY & IMAGING

ID. No.		Age	24Y	Sex	M
Patient Name	MR				
Order Time	07/12/22 01:11	Report Time	07/12/22 06:24		
Referred by	DR. RIKTA KUNDU, MBBS, FCPS.				

MRI OF RIGHT FOOT

TECHNIQUE:

SE T1WI axial, coronal, sagittal; FST2WI axial, coronal & sagittal; FLASH sagittal; STIR sagittal.

FINDINGS:

1. T2WI & STIR images show, hyperintense and minimal irregular thickening of the anterior talo-fibular, inferior tibiofibular, the posterior inferior tibiofibular and the interosseous ligaments without complete frank discontinuity. Hyperintense thickening is also noticed at dorsal talonavicular & posterior talofibular ligaments too.
2. No evidence of osteochondral compression (stress) fractures or fractures at epiphysial end of the tibia or tarsal-metatarsal bones. Tibio-talar joint effusion distends the anterior & posterior capsules.
3. No evidence of focal articular surface flattening or discontinuity, loss of articular cartilage or irregular contour of the subchondral bones.
4. No evidence of obliteration of tarsal sinus or compression to the artery in tarsal canal. Subluxation & distortion (loss of subtalar arch) is not noticed at talocalcaneo-navicular joint due to laxity or tear of interosious ligaments.

IMPRESSION:

Features are suggestive of traumatic soft tissue sprains without evidence of osteochondral compression (stress) fracture. Adv. Other adjuvant examinations for further evaluation, please.

Reported by Electronic Signature

ASST. PROF. DR. RAFAEL MURSALIN

MBBS (DU), Ph. D (Japan)

CT Scan & MRI Specialist.

Head of the department (Ex), CMCH.

Fellow-Resident, Nagasaki Univ. Hosp.

Member RSNA (USA), ESR (Austria)

Consultant, Radiology & Imaging

Ibn Sina Diagnostic & Imaging Center