



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Ref. Interval
-----	---------------	----------------	------	--------------------------

Complete Haemogram Test

Erythrocytes

1	Total RBC	3.36	10 ⁶ /μL	4.1-6
2	Hemoglobin	11.5	g/dL	13 -17.5
3	Hematocrit (PCV)	33.4	%	33-57
4	Mean Corpuscular Volume (MCV)	99.4	fL	80-96
5	Mean Corpuscular Hemoglobin (MCH)	34.2	pg	27.5-33.2
6	Mean Corpuscular Hemoglobin Concentration (MCHC)	34.4	g/dL	30.4-34.5
7	Red Cell Distribution Width (RDW-CV)	13.0	%	12-15
8	Red Cell Distribution Width-SD(RDW-SD)	45.4	fL	30-64.5
9	Nucleated Red Blood Cells	0.0	cells/μL	0 - 1.36
10	Nucleated Red Blood Cells Percentage	0.0	%	0-4

Platelets

11	Platelet Count	154.0	10 ³ /μL	150-450
12	Mean Platelet Volume (MPV)	10.5	fL	6 - 12
13	Platelet Distribution Width (PDW)	15.0	%	15.5-18.3
14	Plateletcrit (PCT)	0.14	%	0.12-0.37

Leucocytes

15	Total Leucocytes Count	5.8	10 ³ /μL	4.4-11
16	Neutrophils	58.0	%	40-77
17	Lymphocyte Percentage	34.0	%	16-44
18	Monocytes Percentage	5.0	%	2.0-10.0
19	Eosinophils Percentage	3.0	%	0-7
20	Basophils Percentage	0.0	%	0 - 1
21	Neutrophils-Absolute Count	3.36	10 ³ /μL	1.8-7.8
22	Lymphocytes-Absolute Count	1.97	10 ³ /μL	1-4.8
23	Monocytes-Absolute Count	0.29	10 ³ /μL	0.1-1.0
24	Eosinophils-Absolute Count	0.17	10 ³ /μL	0 - 0.45



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)





Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Ref. Interval
-----	---------------	----------------	------	--------------------------

Leucocytes

25	Basophils-Absolute Count	0.00	10 ³ /μL	0-0.2
----	--------------------------	------	---------------------	-------

Peripheral Blood Smear

26	RBC Morphology	Normocytic Normochromic
27	WBC Morphology	Within Normal Range
28	Platelets	Adequate On Smear

Interpretation

Sample type: EDTA whole blood.

Test Methods:

RBC/WBC/Platelets: Impedance method,

Hemoglobin: Photometric measurement,

Differential count: VCSn Technology,

MCV, MPV: Measured parameter Indices,

Absolute counts: Calculated.

(Processed on Fully Automated 5 parts differential Hematology analyzer).



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify
Page 2 of 8



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Reference Interval
-----	---------------	----------------	------	-------------------------------

ESR (Erythrocyte Sedimentation Rate)

1	ESR (Erythrocyte Sedimentation Rate) EDTA Whole Blood, Method: Automated ESR, infrared emitting and receiving technology	45	mm/hr	0-15
---	---	----	-------	------

Interpretation

ESR is used to screen for inflammation and/or tissue destruction in the body. ESR is not diagnostic for any disease but indicates need for further studies. ESR is also used to follow-up the course of chronic inflammatory conditions.



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify
Page 3 of 8



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Reference Interval
-----	---------------	----------------	------	-------------------------------

Liver Function Test

1	Bilirubin Total Serum, Method: Jendrassik Grof	0.68	mg/ dL	0.2-1.2
2	Bilirubin Direct Serum, Method: Diazotization	0.28	mg/ dL	0.01 - 0.4
3	Bilirubin Indirect Serum, Method: Calculated	0.4	mg/dL	0.01-1.0
4	Aspartate Transaminase (AST/SGOT) Serum, Method: IFCC without P5P	99.0	U/ L	<50
5	Alanine Transaminase (ALT/SGPT) Serum, Method: IFCC without P5P	46.0	U/ L	<50
6	Alkaline Phosphatase Serum, Method: AMP – pNPP Kinetic	153.5	U/L	30 - 130
7	Total Protein Serum, Method: Biuret end point	6.95	g/dL	6.4 - 8.2
8	Albumin Serum, Method: BCG	3.50	g/dL	3.4 - 5
9	Globulin Serum, Method: Calculated	3.45	g/dL	1.9-3.9
10	A/G ratio Serum, Method: Calculated	1.01	Ratio	1.0 - 2.0
11	Gamma GT Serum, Method: Glupa-C;standardized against IFCC	48.22	U/L	5 - 85



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify
Page 4 of 8



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Reference Interval
-----	---------------	----------------	------	-------------------------------

Kidney Profile

1	BUN (Blood Urea Nitrogen) Serum, Method: Calculated	9.0	mg/dL	3.3 - 18.7
2	Creatinine Serum, Method: Alkaline picrate kinetic	0.65	mg/dL	0.5 - 1.3
3	Urea Serum, Method: Urease-GLDH	20.2	mg/dL	7 - 40
4	BUN/Creatinine ratio Serum, Method: Calculated	13.85		4.0 - 21.5
5	Uric Acid Serum, Method: Uricase, UV	4.65	mg/ dL	2.1 - 7.5
6	Calcium Serum, Method: O cresolphthalein complexone	8.94	mg/dL	8.5 - 10.5
7	eGFR (estimated Glomerular Filtration Rate) Serum, Method: Calculated (MDRD formula)	113.38	mL/min/1.73 m ²	Normal: > 90 Mild decrease in GFR: 60- 89 Moderate decrease in GFR: 30-59 Severe decrease in GFR: 15-29 Kidney failure: < 15

Interpretation

A renal function panel could be ordered when a patient has risk factors for kidney dysfunction such as high blood pressure (hypertension), diabetes, cardiovascular disease, obesity, elevated cholesterol, or a family history of kidney disease. A renal function panel may also be ordered when someone has signs and symptoms of kidney disease, though early kidney disease often does not cause any noticeable symptoms. It may be initially detected through routine blood or urine testing. Renal function panel results are not diagnostic but rather indicate that there may be a problem with the kidneys and that further testing is required to make a diagnosis and determine the cause. Results of the panel are usually considered together, rather than separately. Individual test result can be abnormal due to causes other than kidney disease, but taken together with risks and signs and symptoms, they may give an indication of whether kidney disease is present.



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Reference Interval
-----	---------------	----------------	------	-------------------------------

Lipid Profile

1	Total Cholesterol Serum, Method: Cholesterol oxidase,esterase,peroxidase	183.0	mg/dL	Desirable: <200; Borderline high = 200-239; High: > 240
2	Triglycerides Serum, Method: Enzymatic, end point GPO-POD	263.0	mg/dL	Desirable: <150 Borderline High: 150 - 199 High: > 200 - 499
3	HDL-Cholesterol Serum, Method: Enzymatic Immunoinhibition	70.0	mg/dL	30 - 60
4	LDL- Cholesterol Serum, Method: Calculated	60.0	mg/dL	Optimal: <100; Near Optimal: 100-129; Borderline High: 130-159; High: 160-189; Very high: >190
5	Cholesterol/HDL ratio Serum, Method: Calculated	2.61		Optimal: <3.5 Near Optimal: 3.5 - 5.0 High >5.0
6	VLDL Cholesterol Serum, Method: Calculated	53.0	mg/dL	6 - 40
7	Non HDL Cholesterol Serum, Method: Calculated	113.00	mg/dl	Desirable: <130 Borderline high: 130-159 High : 160-189 Very High :>190
8	LDL /HDL ratio Serum, Method: Calculated	0.86		Optimal: <2.5 Near Optimal: 2.5-3.5 High >3.5

Interpretation

- When serum Triglyceride concentration exceeds 500 mg/dl (4.52 mmol/L) or when chylomicrons are present (Post fed stage), Friedel Equation for estimation of VLDL -C =TG/5 (Friedal et al 1972) is not used. VLDL -C are not calculated. When triglycerides are very high greater than 1000mg/dl , there is a risk of developing pancreatitis in children and adults. Triglycerides change dramatically in response to meals, increasing as much as 5 to 10 times higher than fasting levels just a few hours after eating. Therefore modest changes in fasting triglycerides measured on different days are not considered to be abnormal."
- HDL -Cholesterol : HDL-C is also called Good cholesterol because it removes excess cholesterol from tissues and carries it to the liver for disposal. If HDL-C is less than 40mg/dl for men and less than 50mg/dl for women , there is increased risk of heart disease that is independent of other risk factors including the LDL-C level. NCEP guidelines suggest that an HDL cholesterol value greater than 60 mg/dl is protective.
- LDL Cholesterol : Desired goal for LDL-C levels change based on individual risk factors, For young adults less than 120 mg/dl is acceptable. Values between 120 to 159 mg/dl are considered as borderline high. Values greater than 160 mg/dl are considered high. Low levels of LDL cholesterol may be seen in people with an inherited lipoprotein deficiency and in people with hyperthyroidism, infection, inflammation or cirrhosis.



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

No.	Investigation	Observed Value	Unit	Biological Reference Interval
-----	---------------	----------------	------	-------------------------------

HbA1c (Whole Blood)

1 HBA1c-Glycated Haemoglobin EDTA Whole Blood, Method: HPLC	10.4	%	Non-diabetic: 4-6 Excellent Control: 6-7 Fair to good control: 7-8 Unsatisfactory control: 8-10 Poor Control: >10	
2 Estimated Average Glucose (eAG) EDTA Whole Blood, Method: Calculated	251.78	mg/dL	90-120 mg/dL : Good control 121-150 mg/dL : Fair control 151-180 mg/dL : Unsatisfactory control >180 mg/dL : Poor control	

Interpretation

1.The term HbA1c refers to Glycated Haemoglobin. Measuring HbA1c gives an overall picture of what the average blood sugar levels have been over a period of weeks/month. Higher the HbA1c, the greater the risk of developing diabetes-related complications.

2.HbA1c has been endorsed by clinical groups and ADA (American Diabetes Association) guidelines 2012, for the diagnosis of diabetes using a cut-off point of 6.5%. ADA defined biological reference range for HbA1c is between 4-6%. Patients with HBA1c value between 6.0-6.5% are considered at risk for developing diabetes in the future. Trends in HbA1c area a better indicator of glucose control than standalone test.

3.To estimate the eAG from the HbA1c value, the following equation is used: $eAG(mg/dl) = 28.7 * A1c - 46.7$.

4.Diabetic must aspire to keep values under 7% to avoid the various complications resulting from diabetes.

5.Certain conditions can give rise to a spuriously low HbA1C values. Such conditions include Hemolytic anemias, certain hemoglobinopathies (Hb SS, HbSC, Hb CC, unknown variant), recent blood transfusion,acute blood loss, hypertriglyceridemia, drugs (eg dapsone, ribavirin, trimethoprim-sulfamethoxazole, hydroxyurea, vitamin C/E), chronic liver disease.

6.Certain conditions can give rise to a spuriously high HbA1C values. Such conditions include Iron deficiency, vitamin B12 deficiency, alcoholism, uremia, hyperbilirubinemia, drugs (chronic ingestion of salicylates in high doses and opiate addiction).

Note:The reportable range for HbA1C HPLC analyser is 3.8 % to 18.5 %, eAG calculation not possible above or below this range. In such scenario, observed HbA1c results may not be truly representative of the glycemic control and need to be cross checked by other methods of testing like fructosamine test.

Glucose (Random)

3 Glucose (Random) Fluoride Plasma(R), Method: Hexokinase	247.0	mg/dL	Normal : 79 - 140 Pre - Diabetes: 141 - 200 Diabetes: > 200
---	--------------	--------------	---



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



Scan To Verify
Page 7 of 8



Name:	gyanaranjan Behera	Age/Gender:	52 Years/Male
Referred By:	N.A	Client Name:	N.A
Collection Date:	10-06-2026 11:25:00	Report Release Date:	10-06-2026 22:01:24

Curebay at LAB 2.0

Interpretation

Glucose is the primary energy source for the body's cells and the only energy source for the brain and nervous system. High levels of glucose most frequently indicate diabetes, but many other diseases and conditions can also cause elevated blood glucose. Hypoglycemia is characterized by a drop in blood glucose to a level where first it causes nervous system symptoms (sweating, palpitations, hunger, trembling, and anxiety), then begins to affect the brain (causing confusion, hallucinations, blurred vision, and sometimes even coma and death).

End Of Report



Scan For Report

CRM No :1781069464161
Sample Recd. Time: 10-06-2026 20:49
Report Time: 10-06-2026 22:01
Patient Name: gyanaranjan Behera
Patient ID: 16021300

Authorized Signatory
Dr. Deepika Sahu
MBBS MD (Path)



QUALITY POLICY

GENERAL DIAGNOSTICS INTERNATIONAL (P) Ltd. maintains the highest standards of quality control in all aspects of laboratory work. The purpose of our laboratory's Quality Management System is to ensure that:

- Principles of all accreditations, including that of NABL – ISO1518:2012 (National Accreditation Board of Laboratories) are adhered for each test in the scope of the accreditation, and beyond.
- Test methods, processes and control mechanisms are timely updated and fully validated to ensure the accuracy and reliability of our test results.

The objectives of our Quality Control system are:

- Use Bar-Coded operations to enable full traceability throughout the sample flow process and to ensure sample handling procedures and environmental conditions are managed well and there is no or minimal affect on the results.
- Continually improve the practices of our clients, franchise partners, associate doctors, clinics and hospitals and monitor their training needs. Be proactive in identifying gaps in the processes being followed. Guide them to ensure that the patients are served in the best possible way.
- Report the results with accuracy and clarity in a timely manner. Do a root cause analysis whenever there is a deviation against protocols and find solutions to the identified causes.
- Ensure a continual enhancement, implementation and maintenance of the quality system and seek improvement in the effectiveness of the quality system from experts at regular intervals.
- Meet and exceed expectations with respect to turn-around time, sample collection hygiene & reliability of service.
- Ensure that each test is performed by qualified and trained staff. Provide opportunities to the staff so that they can increase their knowledge and use the same for self and organizational betterment.
- Ensure that the equipment used are best in class, properly maintained and calibrated and where possible, measurements are traceable to recognized standards. Also explore methods which may lead to improvement in equipment performance and methodologies used for conducting tests.
- Enable technology upgrades to achieve higher accuracy and reduced complexities.
- Use internal audits and other checks to ensure the quality system complies with requirements; ensure problems are investigated promptly, root cause(s) established and effective action taken to prevent a recurrence.
- Have a smooth communication mechanism to ensure information is made available as rapidly as possible to those who need it, both internal and external to the organization.
- Monitor, help and support our franchise and service partners to be sensitive on all aspects of service delivery and to ensure quality standards are followed with no exceptions.

CONDITIONS of REPORTING

01. It is presumed that the specimen accompanying the TRF (Test Requisition Form where the details of patient are recorded) is of the same patient whose details are there in the TRF.
02. A test requested might not be performed due to the following reasons(s):
 - 2.1 Insufficient quantity of specimen required to conduct the test.
 - 2.2 Poor quality of the Specimen not meeting the quality criteria (hemolysis of sample/clotted.)
 - 2.3 Incorrect specimen type as required to conduct a test.
03. Test(s) may be partly or fully cancelled due to incorrect test code, incorrect name of the test or incorrect type of specimen. A communication shall be made and it is expected that a fresh specimen will be sent to laboratory for analysis of same parameter(s).
04. The results of laboratory investigation are dependent on the quality of the specimen as well as the assay procedures/technologies used. All samples collected for tests are required to be prepared, stored, labeled and brought to processing laboratory as per the prescribed guidelines of GENERAL DIAGNOSTICS.
05. GENERAL DIAGNOSTICS laboratory cannot be held liable for incorrect results of a sample which deviated from the guidelines issued.
06. There can be several factors like sample's unintended exposure to heat or travel through rough terrain which affect the quality of test results. Therefore a 2% chance of error/deviation in results is a possibility.
07. For certain category of tests, the report may carry a "PRELIMINARY" status implying that the results are yet to be reported for one (or more) tests. For example, in the case with certain microbiology tests, a "FINAL" culture, identification or drug susceptibility result might be pending. In such case, the status "RESULT PENDING" will be mentioned on report. The same shall be replaced by the test results whenever it is ready.
08. If the collection date or any other details was not stated in the Test Requisition Form, the same will not be printed on the report. In cases where the missing information is mandatory for report generation or meeting accreditation guidelines, the sample shall not be processed at all.
09. Tests parameters excluded from the "scope" of NABL accreditation shall be marked by asterisks.
10. In case you are not the intended recipient of the report, please immediately inform the same to the issuing entity. Any use, disclosure, copy or distribution of any contents of such report, is unlawful and is strictly prohibited.
11. Some test may be referred to other laboratories to provide a wider test menu to the patients. The details of the laboratory where the sample was referred to, can be obtained from Customer Care department.
12. Claims of comparing results against that from a different laboratory shall be looked into only if it was the same sample which was split and sent in same conditions to all laboratories and processed on the same technology.



The greatness of a NATION is judged by the way it treats its ANIMALS.