

# Sepsis Monitor AI - Clinical Report

Telemetry Timestamp: 2026-08-14 19:26:19 UTC  
Authorized Practitioner Role: Chief of Department / Sef de Sectie


## 1. Initial Baseline Configuration

| Parameter Matrix                  | Active Interface Entry            |
|-----------------------------------|-----------------------------------|
| Gestational Profile               | Preterm 28w / Prematur 28 sapt    |
| Configured Patient Weight         | 2.50 kg                           |
| Renal Function Tracker (AKI Menu) | Normal Baseline / Functie Normala |
| Kangaroo Care Interaction         | Active / In Bratele Mamei         |
| Music Therapy State               | Active / Meloterapie Pornita      |

## 2. Active Sensor Grid Real-Time Frame

|                                           |                                        |
|-------------------------------------------|----------------------------------------|
| Heart Rate: <b>140 bpm</b>                | Temperature: <b>36.7 °C</b>            |
| Oxygen Saturation (SpO2): <b>98%</b>      | Blood Pressure (BP): <b>65/40 mmHg</b> |
| C-Reactive Protein (CRP): <b>4.2 mg/L</b> | Procalcitonin (PCT): <b>0.4 ng/mL</b>  |

## 3. Historical Biomarker Levels (Blue: CRP | Green: PCT)



## 4. AI Decision Support & Active Guardrail Outputs

[Clinical Analysis & Report Summary]  
[Clinical analysis based on augmented context] Patient Vitals Data: - Heart Rate (hr): 140 beats per minute (tachycardia) - Temperature (temp): 36.7°C (normal) - Oxygen Saturation (spo2): 98% (normal) - Blood Pressure (bp): 65/40 mmHg (hypotension) - C-Reactive Protein (pcr): 4.2 mg/dL (elevated) - Procalcitonin (pct): 0.4 ng/mL (elevated) - Weight (weight\_kg): 2.5 kg (consistent with gestational profile) - Gestational Profile: Preterm 28 weeks / Premature 28 weeks - Renal Status: Normal Baseline / Normal Function - Kangaroo Care Active: Yes - Music Therapy Active: Yes Clinical Severity Stratification: Moderate Inflammation Potential Sepsis Indicators: Elevated C-Reactive Protein (pcr) and Procalcitonin (pct) levels, hypotension (bp), and tachycardia (hr) Intrusion Alert: Potential SQL injection attempt detected. Standard emergency

*antibiotic protocol enforced.*

**[Individualized Medication & Dose Architecture]**

*[Mandated antibiotic or monitoring layout based on clinical guidelines] Ampicillin 50 mg/kg IV q 12 hours (Gram-positive coverage: *Listeria*, Group B *Streptococcus*) Gentamicin 2.5 mg/kg IV q 8 hours (Gram-negative synergy coverage: *Escherichia coli*, *Pseudomonas*) Monitor for signs of sepsis, including respiratory distress, apnea, bradycardia, and decreased urine output Monitor renal function and adjust antibiotic dosing as necessary*

**[Family-Centered Care (FCC) Co-Validation]**

*[Evaluation of non-pharmacological family-centered care interventions] Kangaroo Care: Continue to promote skin-to-skin contact between mother and infant to enhance bonding and promote infant stability Music Therapy: Continue to provide soothing music to promote relaxation and reduce stress for both mother and infant Family Education: Educate parents on signs of sepsis and the importance of monitoring for potential complications [STABLE CLINICAL FCC CORRELATION]: The neonatal subject exhibits physiological stability. Clinical staff encourages extending the Kangaroo Care regimen beyond the baseline 2-hour daily threshold. Prolonged skin-to-skin contact actively reinforces neurodevelopmental maturation and enhances systemic metabolic clearance.*