

Sepsis Monitor AI - Clinical Report

Telemetry Timestamp: 2026-08-12 21:12:32 UTC
Authorized Practitioner Role: Chief of Department / Sef de Sectie


1. Initial Baseline Configuration

Parameter Matrix	Active Interface Entry
Gestational Profile	Preterm 28w / Prematur 28 sapt
Configured Patient Weight	2.50 kg
Renal Function Tracker (AKI Menu)	Severe AKI / Anuria (Retention State)
Kangaroo Care Interaction	Active / In Bratele Mamei
Music Therapy State	Active / Meloterapie Pornita

2. Active Sensor Grid Real-Time Frame

Heart Rate: 140 bpm	Temperature: 36.7 °C
Oxygen Saturation (SpO2): 98%	Blood Pressure (BP): 65/40 mmHg
C-Reactive Protein (CRP): 4.2 mg/L	Procalcitonin (PCT): 0.4 ng/mL

3. Historical Biomarker Levels (Blue: CRP | Green: PCT)



4. AI Decision Support & Active Guardrail Outputs

[Clinical Analysis & Report Summary]
[Clinical analysis based on augmented context] Patient ID: Targeted Patient Vitals Data: {'hr': 140, 'temp': 36.7, 'spo2': 98, 'bp': '65/40 mmHg', 'pcr': 4.2, 'pct': 0.4, 'weight_kg': 2.5, 'gestational_profile': 'Preterm 28w / Prematur 28 sapt', 'renal_status': 'Severe AKI / Anuria (Retention State)', 'kangaroo_care_active': True, 'music_therapy_active': True} Clinical Severity Stratification: Moderate Inflammation Augmented Context Analysis: - Elevated heart rate (140 bpm) and low blood pressure (65/40 mmHg) may indicate sepsis or severe infection. - Temperature is within normal limits (36.7°C). - Oxygen saturation is within normal limits (98%). - Renal status indicates severe acute kidney injury (AKI) and anuria, which may be a risk factor for sepsis. - Gestational profile indicates preterm birth, which may increase the risk of sepsis. - Moderate inflammation is

present, which may be indicative of an infection. - No signs of infection are present in the targeted patient's vitals data.
Intrusion Alert: Potential SQL injection attack detected. Neutralized.

[Individualized Medication & Dose Architecture]

[Mandated antibiotic or monitoring layout based on clinical guidelines] Standard Initial Empiric Antibiotic Protocols: - Ampicillin (Target: Gram-positive coverage, Listeria, Group B Streptococcus) - Gentamicin (Target: Gram-negative synergy coverage, Escherichia coli, Pseudomonas) Emergency Antibiotic Protocol Initiated: Administer Ampicillin 50 mg/kg IV every 12 hours and Gentamicin 2.5 mg/kg IV every 8 hours. Monitoring Layout: - Vital signs every 4 hours - Blood cultures every 12 hours - Urine output monitoring - Renal function monitoring - Infection control measures

[Family-Centered Care (FCC) Co-Validation]

[Evaluation of non-pharmacological family-centered care interventions] Family-centered care interventions: - Kangaroo care: Active (True) - Music therapy: Active (True) Recommendations: - Continue kangaroo care and music therapy to promote bonding and reduce stress. - Educate the family on the importance of hand hygiene and infection control measures. - Provide emotional support to the family during this critical period. [CRITICAL CLINICAL FCC CORRELATION]: Due to acute hemodynamic instability and severe renal impairment (Severe AKI), Kangaroo Care sessions must be conducted with continuous monitoring of arterial lines and umbilical catheters. It is highly recommended to optimize skin-to-skin contact windows to a minimum of 60-90 minutes per session to help stabilize compensatory tachycardia, while strictly avoiding unnecessary infant handling or physical displacement.