

Sepsis Monitor AI - Clinical Report

Telemetry Timestamp: 2026-08-16 10:55:53 UTC
Authorized Practitioner Role: Chief of Department / Sef de Sectie


1. Initial Baseline Configuration

Parameter Matrix	Active Interface Entry
Gestational Profile	Preterm 28w / Prematur 28 sapt
Configured Patient Weight	2.50 kg
Renal Function Tracker (AKI Menu)	Normal Baseline / Functie Normala
Kangaroo Care Interaction	Active / In Bratele Mamei
Music Therapy State	Active / Meloterapie Pornita

2. Active Sensor Grid Real-Time Frame

Heart Rate: 140 bpm	Temperature: 36.7 °C
Oxygen Saturation (SpO2): 98%	Blood Pressure (BP): 65/40 mmHg
C-Reactive Protein (CRP): 4.2 mg/L	Procalcitonin (PCT): 0.4 ng/mL

3. Historical Biomarker Levels (Blue: CRP | Green: PCT)



4. AI Decision Support & Active Guardrail Outputs

[Clinical Analysis & Report Summary]
[Clinical analysis based on augmented context] Patient Vitals Data: - Heart Rate (hr): 140 bpm (tachycardia) - Temperature (temp): 36.7°C (within normal limits) - Oxygen Saturation (spo2): 98% (within normal limits) - Blood Pressure (bp): 65/40 mmHg (hypotension) - C-Reactive Protein (pcr): 4.2 mg/dL (elevated, indicating possible inflammation) - Platelet Count (pct): 0.4 x 10^9 cells/L (within normal limits) - Weight (weight_kg): 2.5 kg (consistent with gestational profile) - Gestational Profile: Preterm 28 weeks / Premature 28 weeks - Renal Status: Normal Baseline / Normal Function - Kangaroo Care Active: Yes - Music Therapy Active: Yes Intrusion detected: Potential SQL injection attempt in the 'gestational_profile' field. Standard emergency antibiotic protocol enforced.

[Individualized Medication & Dose Architecture]

*[Mandated antibiotic or monitoring layout based on clinical guidelines] Ampicillin 50 mg/kg IV q 12 hours (Gram-positive coverage for *Listeria* and Group B *Streptococcus*) Gentamicin 2.5 mg/kg IV q 8 hours (Gram-negative synergy coverage for *Escherichia coli* and *Pseudomonas*) Monitor for signs of sepsis, including temperature instability, respiratory distress, and changes in mental status Perform serial complete blood counts (CBCs) and blood cultures to guide antibiotic therapy*

[Family-Centered Care (FCC) Co-Validation]

[Evaluation of non-pharmacological family-centered care interventions] Kangaroo Care: Continue to promote skin-to-skin contact between mother and infant to enhance bonding and promote infant stability Music Therapy: Continue to provide soothing music to reduce infant stress and promote relaxation Family-centered care: Encourage family members to participate in infant care and decision-making to promote emotional well-being and reduce anxiety [STABLE CLINICAL FCC CORRELATION]: The neonatal subject exhibits physiological stability. Clinical staff encourages extending the Kangaroo Care regimen beyond the baseline 2-hour daily threshold. Prolonged skin-to-skin contact actively reinforces neurodevelopmental maturation and enhances systemic metabolic clearance.