

# **BT1001 UNIT II**

- Biological Diversity
- Chemistry of life: chemical bonds
- Biochemistry and Human biology
- Protein synthesis
- Stem cells and tissue engineering

# BIODIVERSITY

Biological diversity - or biodiversity - is the term given to the varieties of life on Earth. It is the result of billions of years of evolution, shaped by natural processes and, increasingly, by the influence of humans.

It forms the web of life of which we are an integral part and upon which we so fully depend.



# BIODIVERSITY

- 1) Variety of species
- 2) Genetic differences
- 3) Variety of ecosystems

Genetic diversity



Taxonomic diversity



Community or ecosystem diversity



# **WHY IS BIODIVERSITY IMPORTANT FOR YOU AND THE WORLD?**

**Protecting biodiversity is in our self interest,  
providing the goods and services that sustain our  
lives including:**

- Provision of shelter and building materials**
- Stabilization and moderation of the Earth's climate**
- Purification of air and water**
- Provision of food, fuel, and fibre**
- Cultural and aesthetic benefits, etc.**






# WHAT ARE THE CHALLENGES FACING BIODIVERSITY?

**Species have been disappearing at up to 1000 times the natural rate**

- **An estimated 34,000 plant and 5,200 animal species face extinction, including one in eight birds and one third of all amphibians**
- **20% of known bird species have already disappeared**
- **41% of mammals are in decline and 28% are under direct threat**
- **45% of the Earth's original forests are gone. Forest areas of about four times the size of Belgium are being lost every year.**




**Human activities are creating the greatest wave of extinction since the natural disaster that wiped out the dinosaurs 65 million years ago**




# Chemistry of life: chemical bonds

# Chemistry of Life

- All matter is built up of simple units called **atoms**.
- Although the word atom means something that cannot be cut (a = “without,” tom = “cut”), these elementary particles are actually made up of many smaller parts, which are themselves further divisible.
- **Elements** are substances that consist of the same kinds of atoms.
- **Compounds** consist of units called molecules, which are intimate associations of atoms (in the case of compounds, different atoms) joined in precise arrangements.




- Atoms interact with one another to form chemical communities. The tightly knit atoms making up the communal molecules are held together by **chemical bonding**.
- One way of achieving this more stable state is for an atom with very few electrons in its outer shell to donate them to an atom with an outer shell that is almost complete.
  - The atom that donates the electrons will then have more protons than electrons and assume a positive charge; it is called a cation. The atom receiving the electrons assumes a negative charge and is called an anion.
  - These two oppositely charged ions are electrostatically attracted to each other and are said to have an **ionic, or polar, bond**.






- A second way in which atoms may join with one another to bring about a filling of their outermost shells is by sharing a pair of electrons.
  - The two bonding atoms provide one electron each in creating the shared pair. This pair of electrons forms a **covalent bond** that holds the two atoms together. It is represented by a solid line in the formula of a compound.



- In many molecules, covalent bonding may occur not just singly (sharing a single pair of electrons), but may involve the formation of double or triple bonds in which two and even three pairs of electrons are shared.
  - These double and triple bonds tend to fix the position of the participating atoms in a rigid manner.




- **Non-covalent bonds (ionic, hydrogen)** are much weaker than **covalent bonds** (electron sharing) and so protein shape can be disrupted especially by temperature, pH, ions (salt).
- It involves more dispersed variations of electromagnetic interactions.
- Critical in maintaining the three-dimensional structure of large molecules, such as proteins and nucleic acids
- There are four commonly mentioned types of non-covalent interactions: hydrogen bonds, ionic bonds, van der Waals forces, and hydrophobic interactions.
  - The noncovalent interactions hold together the two strands DNA in the double helix, stabilize secondary and tertiary structures of proteins, and enable enzyme-substrate binding and antibody-antigen association.


#### HYDROGEN BONDS

Because they are polarized, 2 adjacent  $\text{H}_2\text{O}$  molecules can form a linkage known as a hydrogen bond. Hydrogen bonds have only about 1/20 the strength of a covalent bond.

Hydrogen bonds are strongest when the 3 atoms lie in a straight line.



b



# Biochemistry and Human biology

# **Biochemistry:**


## **Where Chemistry & Biology Meet**

- **Living things require millions of chemical reactions just to survive.**
- **Metabolism = all the chemical reactions occurring in the body.**
- **Organic molecules:**
  - usually associated with living things.
  - always contain CARBON.
  - are “large” molecules, with many atoms
  - always have covalent bonds (share electrons)

# Biochemistry and Human Biology

- **Biochemistry:** Science concerned with the chemical constituents of living cells and with the reaction and process that they undergo.
  - Complete understanding at the molecular level of all the chemical processes associated with living cells
  - An appreciation of the biochemistry of less complex form of life is often direct relevance to human biochemistry
- **Reciprocal relationship between biochemistry and medicine has stimulated mutual advance**
  - Biochemistry studies have illuminated many aspects of health & disease






<b>S. No.</b>	<b>Disease</b>	<b>Causes</b>
<b>1</b>	<b>Scurvy rickets</b>	<b>deficiencies of vitamins C and D respectively</b>
<b>2</b>	<b>Atherosclerosis</b>	<b>genetic, dietary, environmental factors</b>
<b>3</b>	<b>Cystic fibrosis</b>	<b>mutation in the gene coding the CFTR protein (Cystic fibrosis transmembrane conductance regulator, a protein involved in the transport of chloride ions across cell membranes)</b>
<b>4</b>	<b>Cholera</b>	<b>exotoxin of vibrio cholera</b>
<b>5</b>	<b>Diabetes mellitus type I</b>	<b>genetic and environmental factors resulting in deficiency of insulin</b>
<b>6</b>	<b>Phenylketonuria</b>	<b>mainly mutation in the gene coding phenylalanine hydroxylase</b>

# Carbon-based Molecules

- Although a cell is mostly water, the rest of the cell consists mostly of carbon-based molecules


**Organic chemistry is the study of carbon compounds**



# Carbon is a Versatile Atom


- It has four electrons in an outer shell that holds eight

Carbon can share its electrons with other atoms to form up to four covalent bonds



# Giant Molecules - Polymers

- Large molecules are called polymers
- Polymers are built from smaller molecules called monomers
- Biologists call them macromolecules



## Macromolecules in Organisms

- There are four categories of large molecules in cells:

Carbohydrates

Lipids

Proteins

Nucleic Acids




# Examples of Polymers

•Proteins





Lipids



Carbohydrates



Glycogen



Nucleic Acids



# Carbohydrates


- **Carbohydrates include:**

- Small sugar molecules in soft drinks
- Long starch molecules in rice, wheat, pasta and potatoes



# Linking Monomers


Cells link monomers by a process called **condensation or dehydration synthesis** (removing a molecule of water)



This process joins two sugar monomers to make a double sugar

# Breaking Down Polymers

- Cells break down macromolecules by a process called hydrolysis (adding a molecule of water)



**Water added to split a double sugar**

# Monosaccharides


- Called simple sugars

Include glucose, fructose, & galactose

Have the same chemical, but different structural formulas




Hexose Sugars





# Cellular Fuel

- Monosaccharides are the main **fuel** that cells use for cellular work




**ATP**

# Disaccharides

- A disaccharide is a double sugar.

They're made by joining two monosaccharides

Involves removing a water molecule (condensation)





Bond called a GLYCOSIDIC bond

# Polysaccharides

- **Complex carbohydrates**


**Composed of many sugar monomers linked together**

**Polymers of monosaccharide chains**



# Lipids

- Lipids are hydrophobic –“water fearing”
- **Do NOT mix with water**
- Includes fats, waxes, steroids, & oils
- Fats **store energy**, help to **insulate the body**, and cushion and protect organs




FAT MOLECULE

# Types of Fatty Acids

**Saturated fatty acids** have the maximum number of hydrogens bonded to the carbons (all single bonds between carbons)

**Unsaturated fatty acids** have less than the maximum number of hydrogens bonded to the carbons (a double bond between carbons)

Single  
Bonds in  
Carbon  
chain




Double bond in carbon chain




# Triglyceride

- Monomer of lipids
- Composed of Glycerol & 3 fatty acid chains
- Glycerol forms the “backbone” of the fat





Organic Alcohol  
(-OL ending)




# Lipids & Cell Membranes

- Cell membranes are made of lipids called **phospholipids**
- Phospholipids have a **head** that is polar & attract water (**hydrophilic**)
- Phospholipids also have **2 tails** that are nonpolar and do not attract water (**hydrophobic**)




b. Plasma membrane of a cell



Cell membrane with proteins & phospholipids

# Steroids

- The carbon skeleton of steroids is bent to form 4 fused rings
- Cholesterol is the “base steroid” from which your body produces other steroids
- Estrogen & testosterone are also steroids



## Synthetic Anabolic Steroids


- They are variants of testosterone
- Some athletes use them to build up their muscles quickly
- They can pose serious health risks





# Waxes

- A wax is a lipid because of its nonpolar solubility characteristics as well as its extremely hydrophobic (water-hating) properties.
- Waxes are composed of a single, highly complex alcohol joined to a longchain fatty acid in a typical ester linkage.
- Waxes are important structural lipids often found as protective coatings on the surfaces of leaves, stems, hair, skin, etc.
- They provide effective barriers against water loss and in some situations make up the rigid architecture of complex structures such as the honeycomb of the beehive.
- They serve a commercial use as well, in furniture polish, automobile coating compounds, and floor finishes.



# Proteins


- Proteins are polymers made of monomers called amino acids

All proteins are made of 20 different amino acids linked in different orders

Proteins are used to build cells, act as hormones & enzymes, and do much of the work in a cell


# Nucleic Acids

- Store hereditary information
- Contain information for making all the body's proteins
- Two types exist --- DNA & RNA



# DNA-Deoxyribonucleic acid


- **Two strands of DNA join together to form a double helix**
- **Nucleotides form long chains called DNA**
- **Nucleotides are joined by sugars & phosphates on the side**





# Nucleic Acids

Nucleic acids  
are polymers  
of nucleotides



# Bases





• Each DNA nucleotide has one of the following bases:

– Adenine (A)

– Guanine (G)


– Thymine (T)

– Cytosine (C)







# RNA – Ribonucleic Acid

- Ribose sugar has an extra –OH or hydroxyl group
- It has the base uracil (U) instead of thymine (T)



# Macromolecules


Biological macromolecule	Function	Monomer	Examples
<b>Carbohydrates</b>	Dietary energy; storage; plant structure	 <p><b>Monosaccharide</b></p>	<b>Monosaccharides:</b> glucose, fructose. <b>dissaccharides:</b> lactose, sucrose. <b>Polysaccharides:</b> starch, cellulose.
<b>Lipids</b>	Long-term energy storage (for fats); hormones (for steroids)	 <p><b>Components of a fat molecule</b></p>	<b>Fats, oils, steroids</b>
<b>Proteins</b>	Enzymes, structure, storage, contraction, transport, etc.	 <p><b>Amino acid</b></p>	<b>Lactase (an enzyme), hemoglobin</b>
<b>Nucleic acids</b>	Information storage	 <p><b>Nucleotide</b></p>	<b>DNA, RNA</b>

# **Protein Synthesis**


# Protein Synthesis

- The production (synthesis) of polypeptide chains (proteins)
- Two phases:  
**Transcription & Translation**
- mRNA must be processed before it leaves the nucleus of eukaryotic cells

Prokaryotic Cell





Eukaryotic Cell



# Discovery of DNA structure

- Walter Sutton discovered **chromosomes** were made of **DNA and Protein**
- However, scientists were **NOT** sure which one (protein or DNA) was **the actual genetic material of the cell**
- **Frederick Griffith** in 1928 showed the **DNA** was the cell's genetic material
- **Rosalind Franklin** took diffraction **x-ray** photographs of DNA crystals
- **Watson & Crick** in the 1950's built of DNA





Rosalind Franklin





# Discovery of DNA Structure

- **Erwin Chargaff** showed the amounts of the four bases on DNA ( A,T,C,G)
- In a body or somatic cell:
  - A = 30.3%**
  - T = 30.3%**
  - G = 19.5%**
  - C = 19.9%**
- **Chargaff's rule:**
  - **Adenine** must pair with **Thymine**
  - **Guanine** must pair with **Cytosine**
  - The bases form weak hydrogen bonds




# Structure of DNA

- DNA is made of subunits called **nucleotides**
- DNA nucleotides are composed of a **phosphate**, **deoxyribose** sugar, and **a nitrogen-containing base**
- The 4 bases in DNA are: **adenine (A)**, **thymine (T)**, **guanine (G)**, and **cytosine (C)**
- **Purines** have **single** rings of carbon-nitrogen (G, A)
- **Pyrimidines** have **double** carbon-nitrogen rings (C, T)
- This is called ***complementary base pairing*** because a **purine** is always paired with a **pyrimidine**

## 5' to 3' Sugars


- When the DNA **double helix** unwinds, it **resembles a ladder**
- The **sides** of the ladder are the **sugar-phosphate backbones**
- The **rungs** of the ladder are the **complementary paired bases**
- The two DNA strands are **anti-parallel** (they run in opposite directions)

## Anti-Parallel Strands of DNA




# Steps in DNA Replication

- Occurs when **chromosomes duplicate** (make copies)
- An **exact copy** of the DNA is produced with the aid of the enzyme **DNA polymerase**
- **Hydrogen bonds** between bases **break** and enzymes “unzip” the molecule
- Each **old strand** of nucleotides serves as a **template** for each new strand
- **New nucleotides** move into complementary positions are joined by **DNA polymerase**




## Two New, Identical DNA Strands Result from Replication


# Another View of Replication



Parental DNA molecule contains so-called old strands hydrogen-bonded by complementary base pairing.



Region of replication. Parental DNA is unwound and unzipped. New nucleotides are pairing with those in old strands.




Replication is complete. Each double helix is composed of an old (parental) strand and a new (daughter) strand.




# RNA Differs from DNA


1. **RNA** has a sugar **ribose**  
**DNA** has a sugar **deoxyribose**
2. **RNA** contains the base **uracil (U)**  
**DNA** has **thymine (T)**
3. **RNA** molecule is **single-stranded**  
**DNA** is **double-stranded**



Pentose sugars



Ribose (in RNA)



Deoxyribose (in DNA)

# Three Types of RNA

- **Messenger RNA (mRNA)** carries genetic information to the ribosomes  
(blueprint for the construction of a protein)
- **Ribosomal RNA (rRNA)**, along with protein, makes up the ribosomes  
(construction site where the protein is made)
- **Transfer RNA (tRNA)** transfers amino acids to the ribosomes where proteins are synthesized  
(truck delivering the proper amino acid to the site at the right time)



# Genes & Proteins

- **Proteins** are made of **amino acids** linked together by **peptide bonds**
- **20** different amino acids **exist**
- Amino acids chains are called **polypeptides**
- Segment of DNA that codes for the amino acid sequence in a protein are called **genes**


## Genetic Code:

- DNA contains a **triplet code**
- Every three bases on DNA stands for **ONE amino acid**
- Each three-letter unit on **mRNA** is called a **codon**
- **Most amino acids have more than one codon!**
- There are **20 amino acids** with a possible 64 different triplets
- The code is nearly **universal** among living organisms

First Base	Second Base				Third Base
	U	C	A	G	
U	UUU phenylalanine	UCU serine	UAU tyrosine	UGU cysteine	U
	UUC phenylalanine	UCC serine	UAC tyrosine	UGC cysteine	C
	UUA leucine	UCA serine	UAA stop	UGA stop	A
	UUG leucine	UCG serine	UAG stop	UGG tryptophan	G
C	CUU leucine	CCU proline	CAU histidine	CGU arginine	U
	CUC leucine	CCC proline	CAC histidine	CGC arginine	C
	CUA leucine	CCA proline	CAA glutamine	CGA arginine	A
	CUG leucine	CCG proline	CAG glutamine	CGG arginine	G
A	AUU isoleucine	ACU threonine	AAU asparagine	AGU serine	U
	AUC isoleucine	ACC threonine	AAC asparagine	AGC serine	C
	AUA isoleucine	ACA threonine	AAA lysine	AGA arginine	A
	AUG ( <i>start</i> ) methionine	ACG threonine	AAG lysine	AGG arginine	G
G	GUU valine	GCU alanine	GAU aspartate	GGU glycine	U
	GUC valine	GCC alanine	GAC aspartate	GGC glycine	C
	GUA valine	GCA alanine	GAA glutamate	GGA glycine	A
	GUG valine	GCG alanine	GAG glutamate	GGG glycine	G

# Overview of Transcription

- During **transcription** in the nucleus, a segment of DNA unwinds and unzips, and the **DNA** serves as a **template for mRNA formation**
- **RNA polymerase** joins the RNA nucleotides so that the **codons in mRNA are complementary** to the triplet code in DNA
- The transfer of information in the **nucleus** from a **DNA** molecule to an **RNA** molecule
- Only 1 **DNA** strand serves as the **template**
- Starts at promoter **DNA** (TATA box)
- Ends at terminator **DNA** (stop)
- When complete, **pre-RNA** molecule is released




Transcription is going here—the nucleotides of mRNA are joined by the enzyme RNA polymerase in an order complementary to a strand of DNA.

This mRNA transcript is ready to move into the cytoplasm.

# RNA Polymerase


- **Enzyme** found in the nucleus
- **Separates** the two DNA strands by **breaking the hydrogen bonds** between the bases
- Then moves along one of the DNA strands and **links RNA nucleotides** together



# Processing Pre-mRNA

- Also occurs in the **nucleus**
- **Pre-mRNA** made up of segments called **introns & exons**
- **Exons code for proteins, while introns do NOT!**
- Introns spliced out by **spliceosome-enzyme** and exons re-join
- End product is a **mature RNA** molecule that leaves the nucleus to the cytoplasm


pre-RNA molecule



Mature RNA molecule

# Messenger RNA (mRNA)


- Carries the information for a specific protein
- Made up of **500 to 1000** nucleotides long
- Sequence of 3 bases called **codon**
- **AUG** – methionine or **start** codon
- **UAA, UAG, or UGA** – **stop** codons



# Transfer RNA (tRNA)

- Made up of **75 to 80 nucleotides** long
- Picks up the appropriate **amino acid** floating in the cytoplasm
- Transports **amino acids** to the **mRNA**
- Have **anticodons** that are complementary to **mRNA codons**
- Recognizes the appropriate **codons** on the **mRNA** and bonds to them with H-bonds
- **Four ATP's** are required for each amino acid added to the polypeptide chain: Two to "charge" the tRNA, one to carry the charged tRNA to the ribosome and one to move the ribosome to the next codon.


amino acid  
attachment site





# Ribosomal RNA (rRNA)

- Made up of rRNA is **100 to 3000 nucleotides** long
- Made inside the **nucleus** of a cell
- Associates with **proteins to form ribosomes**




## Ribosomes

- Made of a **large and small** subunit
- Composed of **rRNA (40%)** and **proteins (60%)**
- Have **two sites** for tRNA attachment --- **P and A**

# Ribosomes


P= Peptide site  
A= Amino acid site




# Translation

- **Synthesis of proteins** in the cytoplasm
- **Involves the following:**
  1. **mRNA (codons)**
  2. **tRNA (anticodons)**
  3. **ribosomes**
  4. **amino acids**
- **Three steps:**
  1. **initiation:** start codon (AUG)
  2. **elongation:** amino acids linked
  3. **termination:** stop codon (UAG, UAA, or UGA).

# mRNA Codons Join the Ribosome



# Initiation



# Elongation

peptide bond

aa1

aa2

aa3

1-tRNA

2-tRNA

3-tRNA

G

A

A

U

A

C

G

A

U

A

U

G

C

U

A

C

U

U

C

G


A


codon

mRNA

anticodon


hydrogen bonds








Ribosomes move over one codon













Ribosomes move over one codon



# End Product –The Protein!

- The end products of protein synthesis is a **primary structure** of a protein
- A **sequence of amino acid** bonded together by peptide bonds





# Eukaryotic Protein Synthesis Differs from Prokaryotic Protein Synthesis Primarily in Translation Initiation

- The basic plan of protein synthesis in eukaryotes and archaea is similar to that in bacteria.
- Eukaryotic protein synthesis entails more protein components than does prokaryotic protein synthesis, and some steps are more intricate.
- 1. Ribosomes.
  - Eukaryotic ribosomes are larger: consist of a 60S large subunit and a 40S small subunit, which come together to form an 80S particle having a mass of 4200 kd,
    - 40S subunit contains an 18S RNA
    - 60S subunit contains 5S, 5.8S, and 28S
  - Prokaryotic ribosomes have small (in *E. coli* , 30S) and larger (50S) subunits.
    - The 30S unit has 16S rRNA and 21 different proteins.
    - The 50S subunit consists of 5S and 23S rRNA and 34 different proteins.

- 2. *Initiator tRNA*. In eukaryotes, the initiating amino acid is methionine rather than *N*-formylmethionine. However, as in prokaryotes, a special tRNA participates in initiation. This aminoacyl-tRNA is called Met-tRNA<sub>i</sub> or Met-tRNA<sub>f</sub> (the subscript “i” stands for initiation, and “f” indicates that it can be formylated in vitro).
- 3. *Initiation*. The initiating codon in eukaryotes is always AUG. In contrast, a prokaryotic mRNA can have multiple start sites, and it can serve as a template for the synthesis of several proteins. Eukaryotes utilize many more initiation factors than do prokaryotes, and their interplay is much more intricate.
- 4. *Elongation and termination*. Eukaryotic elongation factors EF1 $\alpha$  and EF1 $\beta\gamma$  are the counterparts of prokaryotic EF-Tu and EF-Ts.
  - Termination in eukaryotes is carried out by a single release factor, eRF1, compared with two in prokaryotes. Finally, eIF3, like its prokaryotic counterpart IF3, prevents the reassociation of ribosomal subunits in the absence of an initiation complex.


# **Stem cells and tissue engineering**

**Stem cell (SC)-** Stem Cells are the cells which has the ability to divide for indefinite periods and which can give rise to specialized cells of various tissues of body.

# Stem cells history

- **1978 - Stem cells were discovered in human cord blood**
- **1981 - First in vitro stem cell line developed from mice**
- **1998 - Researchers isolated stem cells from human embryos**
- **1999 - First Successful human transplant of insulin-making cells from cadavers**

# Importance



# Properties

- Two defining property-Ability to **differentiate** into other cells, ability to **self regenerate**
- It can be maintained in the *in vitro* conditions for extended period using artificial medium
- Its karyotype remains stable even after many division
- It can produce any type of adult cells of the organisms.



# Sources of stem cells

## 1. EMBRYONIC STEM CELLS

- Isolated from blastocyst stage of embryos
- Pluripotent (capable of developing into almost all the cell types of the body) in nature

## 2. ADULT STEM CELLS

- It is present in all the organs, but very little amount
- They are multipotent (ability of a single stem cell to develop into more than one cell type but with ability to differentiate into a closely related family of cells).

# **Classification of stem cells**

**Unipotent** - cell which can make exact copies of itself indefinitely, can differentiate, and produce same type of cells eg. Adult muscle stem cell

**Totipotent**- can become any cell type  
eg: Morula stage of Embryos

**Pluripotent**- almost any kind of cell except placenta  
eg: Embryonic SC

**Multipotent**- Produces limited range of cell types  
Adult SC: nerve cells, blood cells, muscle cells, bone and skin cells.

# **Culture of embryonic stem cells**

- Collection of embryos from IVF centres
- Isolation of ICM (inner cell mass) from the blastocyst stage
- Transfer ICM to the center of culture plate containing feeder cells and growth medium
- It can be differentiated to any cells by adding specific medium

**Blastocyst**  
(64 to 200 cell stage,  
cross-section)



*Propagation  
in Culture*


**Inner cell mass**



**Pluripotent  
embryonic  
stem cells**


*Differentiation*

**Development of specialized cells**



# Applications

- It can be used for neural degenerative diseases (Alzheimer's and Parkinson's disease)
- It can also be used to treat diabetes by injecting *in vitro* grown pancreatic islet cells
- It is also used for treating **bone related diseases by injecting *in vitro*** developed osteocytes, chondrocytes and myocytes
- Today most of the diseases get cured through stem cell therapy eg. Muscular dystrophy, Polio.





## In vitro fertilization

