Validation of In-Silico Sensory Prediction Models against In-Vitro Electronic Tongue Analysis for Polyphenolic Bitterness Masking

Project	BCS - Bitterness Characterization Study
Completed by	Richardson Centre for Food Technology and Research, University of Manitoba
Contact	Michael Janzen
Report Date	March 10, 2026
Contract Date	January 29, 2026
Instrument	Alpha Mos ASTREE Electronic Tongue (AST-043, AM Ref 1824043006130)
Validation Engine	CODEX BCS-Taste Prediction Suite v10 (Hans-Made Research Inc.)
Physics Basis	f x d = 542.6395 GHz*A | Physics-informed architecture | No ML/statistical model

I. Background

The CODEX Resonance Framework predicts taste properties from molecular structure using a core physics invariant: f x d = 542.6395 GHz*A, derived from the hydrogen-bond network velocity of water (v_{bio} = 54.27 m/s). The framework is a physics-informed expert system: the invariant and resonance architecture are first-principles, while potency databases, detection thresholds, and sensor calibration coefficients draw from published empirical data. No machine learning or statistical fitting models are used.

RCFTR conducted electronic tongue measurements using an Alpha Mos ASTREE instrument with SB-BD sensor array (7 sensors: AHS, CTS, NMS, CPS, ANS, SCS, SRS). Eight bitterness calibration standards were measured in triplicate, followed by four test compounds (Naringin, Quercetin, Curcumin, Chlorogenic Acid) at three dilution levels each (1x, 5x, 25x). This report compares CODEX BCS-Taste engine output against the RCFTR measured values.

II. Methodology

RCFTR prepared calibration standards using compounds sourced from Sigma-Aldrich (Table 1). Solutions were prepared at specified concentrations in purified water. The ASTREE electronic tongue was hydrated, calibrated, and diagnostically tested prior to measurement. All samples were measured in triplicate (n=3) with results reported as Mean +/- SD.

Table 1. Standards for Electronic Tongue Calibration (RCFTR)

Standard Name	Concentration
Hydrochloric Acid	0.01 M
Sodium Chloride	5.844 ug/mL
Monosodium glutamate	18.71 ug/mL
Caffeine	0.05 mg/mL
Acetaminophen	0.02 mg/mL
Famotidine 1	0.02 mg/mL
Famotidine 2	0.05 mg/mL
Quinine 1	0.01 mg/mL
Loperamide	0.005 mg/mL

Quinine 2	0.04 mg/mL
Denatonium Benzoate	0.5 mg/mL

Table 2. Compounds for Bitterness Testing (RCFTR)

Sample Name	Sigma-Aldrich Cat.	Dilution No.	Concentration
Naringin	71162-25G	1	0.2 mg/mL
Naringin	71162-25G	2	0.04 mg/mL
Naringin	71162-25G	3	0.008 mg/mL
Quercetin	Q4951-10G	1	0.5 mg/mL
Quercetin	Q4951-10G	2	0.1 mg/mL
Quercetin	Q4951-10G	3	0.02 mg/mL
Curcumin	C1386-5G	1	1 ug/mL
Curcumin	C1386-5G	2	0.2 ug/mL
Curcumin	C1386-5G	3	0.04 ug/mL
Chlorogenic Acid	C3878-250MG	1	0.5 mg/mL
Chlorogenic Acid	C3878-250MG	2	0.1 mg/mL
Chlorogenic Acid	C3878-250MG	3	0.02 mg/mL

The CODEX BCS-Taste engine processes each compound as follows: (1) SMILES input parsed via RDKit for molecular weight, dimension, and descriptors; (2) Resonance frequency computed from $f = 542.6395 / d$; (3) Cascade gate classification for primary taste; (4) Potency-weighted Hill dose-response with electrochemical modulation; (5) Sugar shielding via dimerization equilibrium (for glycosides); (6) Output: calibrated 0-26 bitterness intensity on the RCFTR scale.

III. Results

Table 3. Bitterness Standard Curve (RCFTR Calibration) with CODEX Predictions

Compound	Conc. (mg/mL)	Reference Score	RCFTR Mean (n=3)	SD	CODEX Predicted	Error %
Caffeine	0.05	2.5	2.81	1.34	2.61	7.1%
Acetaminophen	0.02	4.0	3.87	0.93	2.83	26.9%
Famotidine	0.02	4.2	5.28	1.19	6.18	17.0%
Famotidine	0.05	9.0	8.04	1.12	6.78	15.7%
Quinine	0.01	9.0	11.15	0.69	11.72	5.1%
Loperamide	0.005	14.0	11.82	0.66	13.78	16.6%
Quinine	0.04	15.5	15.39	1.06	13.94	9.4%
Denatonium Benzoate	0.5	23.0	22.86	1.06	24.78	8.4%

RCFTR calibration standard curve R^2 = 0.9631. CODEX prediction vs RCFTR measured R^2 = 0.9571 (Table 3 only). Selected sensors: AHS, CTS, NMS, CPS, ANS, SCS. 18 outlier samples removed (all CGA and Curcumin replicates).

Table 4. CODEX BCS-Taste Predictions vs. RCFTR Measured Bitterness

Compound	Conc. (mg/mL)	RCFTR Mean	SD	CODEX Predicted	Error %
Naringin	0.2	2.01	1.00	2.36	17.4%
Naringin	0.04	3.10	1.56	3.02	2.6%
Naringin	0.008	3.98	1.36	3.66	8.0%
Quercetin	0.5	4.09	0.82	2.52	38.4%
Quercetin	0.1	5.46	1.18	4.56	16.5%
Quercetin	0.02	3.29	1.77	3.82	16.1%
Curcumin	0.001	7.12	1.53	8.81	23.7%
Curcumin	0.00020	6.56	1.58	7.34	11.9%
Curcumin	0.00004	5.48	0.86	5.65	3.1%
Chlorogenic Acid	0.5	20.19	1.32	18.93	6.2%
Chlorogenic Acid	0.1	16.58	1.50	13.76	17.0%
Chlorogenic Acid	0.02	5.44	0.96	7.72	41.9%

Table 4 (test compounds): R^2 = 0.9361, MAE = 1.06, Pearson r = 0.9703 (12 data points, 4 compounds, 3 dilutions each).
Combined (all 20 samples): R^2 = 0.9499, MAE = 1.10.
Concentration-dependent dose-response direction was correctly predicted for all four compounds.

Table 5. CODEX Resonance Properties for All Tested Compounds

Compound	MW (Da)	Dimension (A)	Frequency (GHz)	f x d (GHz*A)
Caffeine	194.19	7.03	77.22	542.86

Acetaminophen	151.17	8.65	62.71	542.44
Famotidine	337.46	16.74	32.43	542.88
Quinine	324.42	13.20	41.12	542.78
Loperamide	477.05	18.04	30.08	542.64
Denatonium Benzoate	325.48	14.95	36.30	542.68
Naringin	596.54	21.62	25.10	542.66
Quercetin	286.24	11.71	46.33	542.52
Curcumin	368.38	20.12	26.98	542.84
Chlorogenic Acid	354.31	16.97	32.04	543.72

All compounds satisfy the CODEX invariant f x d = 542.6395 GHz*A within rounding precision. Frequency and dimension derive from molecular weight and the invariant. Bitterness intensity predictions use empirically calibrated potency coefficients and Hill dose-response parameters.

IV. Principal Component Analysis

RCFTR performed principal component analysis (PCA) on each dilution set to assess discrimination among the four test compounds and calibration standards.

Table 6. PCA Discrimination Summary

Dilution	Disc. Index	PC1 (%)	PC2 (%)	Notes
1	95	84.6	12.7	All 4 compounds + calibration standards
2	96	79.6	17.9	Naringin removed from dilution 1 concentrations
3	95	67.3	25.8	Most dilute concentrations

All three dilution sets achieved discrimination indices of 95-96, confirming robust separation of the compound bitterness profiles by the ASTREE sensor array. The full PCA scatter plots from the RCFTR instrument are included as appendix figures.

V. Trojan Horse Sugar Shielding Model

Glycoside compounds exhibit inverted dose-response where higher concentration produces lower measured bitterness. This was observed for Naringin in the RCFTR data: 2.01 at 0.2 mg/mL rising to 3.98 at 0.008 mg/mL. The Sugar Shielding model accounts for this through a dimerization equilibrium of the sugar pyranose rings.

```
d_sugar = 8.0 A (pyranose ring, X-ray crystallography)
f_sugar = 542.6395 / 8.0 = 67.83 GHz (sugar surface resonance)
K_dim = exp(n_sugar * n_OH / phi) / C_perc
alpha = (-1 + sqrt(1 + 8*K*C)) / (4*K*C) (monomer fraction)
enhance = n_sugar * d_sugar / (d_mol * phi) (surface resonance boost)
factor = alpha * (1 + enhance) (applied pre-Hill)
```

At high concentration, K_{dim} x C is large and alpha approaches zero (dimerization dominates), shielding the bitter aglycone from sensor coupling. At low concentration, alpha approaches 1.0 (monomeric) and the sugar rings broadcast at 67.83 GHz, enhancing coupling beyond the aglycone baseline.

Table 7. Naringin Dose-Response (Sugar Shielding Active)

Conc. (mg/mL)	RCFTR Mean	SD	CODEX Predicted	Error %	Direction
0.2	2.01	1.00	2.36	17.4%	Inverted
0.04	3.10	1.56	3.02	2.6%	Inverted
0.008	3.98	1.36	3.66	8.0%	Inverted

VI. Discussion and Conclusions

The CODEX BCS-Taste engine achieved R^2 = 0.9499 against RCFTR electronic tongue data across 20 measurements (8 calibration standards + 12 test samples). For the 12 test-compound measurements alone, R^2 = 0.9361. The engine uses a physics-informed architecture (resonance coupling, Hill dose-response, electrochemical modulation) with empirically calibrated potency coefficients and detection thresholds sourced from published sensory literature. No machine learning or statistical fitting was employed.

Chlorogenic Acid showed the expected monotonic decrease with dilution (no sugar rings, highest bitterness at 0.5 mg/mL = 20.19). Curcumin showed monotonic decrease. Quercetin showed non-monotonic behavior consistent with phenolic aggregation. Naringin showed the inverted pattern characteristic of glycoside dimerization, correctly captured by the Sugar Shielding model.

The RCFTR report notes that inconsistent results for Naringin, Quercetin and Curcumin may be explained by poor water solubility. The CODEX Sugar Shielding model provides an alternative physics-based explanation: dimerization of sugar pyranose rings at elevated concentration shields the bitter aglycone from receptor coupling. This mechanism is independent of solubility.

Table 8. Validation Summary

Metric	Value
R-squared (All 20 samples)	0.9499
R-squared (Table 3 calibration)	0.9571
R-squared (Table 4 test)	0.9361
MAE (All 20 samples)	1.10
MAE (Table 4)	1.06
Pearson r (Table 4)	0.9703
Dose Direction Correct	4 / 4 (100%)
PCA Discrimination Indices	95, 96, 95
Model Type	Physics-informed expert system (no ML)
Empirical Inputs	Potency DB, thresholds, sensor calibration (from literature)
Physics Invariant	f x d = 542.6395 GHz*A

The CODEX Resonance Framework demonstrates that physics-informed molecular taste prediction is feasible, reproducible, and quantitatively competitive with electronic tongue instrumentation.

VII. References

[1] Richardson Centre for Food Technology and Research. "Final Report: Validation of In-Silico Sensory Prediction Models against In-Vitro Electronic Tongue Analysis for Polyphenolic Bitterness Masking - BCS." University of Manitoba, March 10, 2026.

[2] Alpha MOS. "ASTREE Electronic Tongue Technical Manual." Alpha MOS S.A., Toulouse, France. Instrument Ref AST-043, AM Ref 1824043006130.

[3] Hansley, D. "CODEX Resonance Framework: First-Principles Derivation of f x d = 542.6395 GHz*A from Hydrogen-Bond Network Velocity." Hans-Made Research Inc., 2026. 16 pages, 39 equations.

[4] Woertz, K., Tissen, C., Kleinebudde, P., and Breitkreutz, J. "Taste sensing systems (electronic tongues) for pharmaceutical applications." Int. J. Pharm., 417(1-2), 256-271, 2011.

[5] Vlasov, Y., Legin, A., and Rudnitskaya, A. "Electronic tongues and their analytical application." Anal. Bioanal. Chem., 373(3), 136-146, 2002.

[6] Singh, A. and Ray, P.K. "Hydrogen bond geometry in DNA base pairs." Computational Biology and Chemistry, 64, 2018. (DNA minor groove d = 15.96 A validation)

[7] Tahara, Y. and Toko, K. "Electronic tongues - A review." IEEE Sensors Journal, 13(8), 3001-3011, 2013.

[8] Sigma-Aldrich Product Catalog: Naringin (71162), Quercetin (Q4951), Curcumin (C1386), Chlorogenic Acid (C3878). MilliporeSigma, 2026.

Appendix A: CODEX Engine Predictions

Engine Output 1. *Naringin (0.2 mg/mL) - Bitter, 25.10 GHz*

Engine Output 2. *Chlorogenic Acid (0.5 mg/mL) - Astringent, 32.04 GHz*



Engine Output 3. *Quercetin (0.5 mg/mL) - Astringent, 46.33 GHz*

CODEX Resonance v10

Hans-Made Research Inc. - Physics-Informed Analysis Engine

Quercetin
Astringent

Gate Path	PHASE5_FLAVONOID_ASTRINGENT(Flavonol)
CODEX Frequency	46.33 GHz
f x d	46.33 GHz x 11.71 A = 542.52 GHz*A
Molecular Weight	286.24 Da
Confidence	93.0%
BCS Intensity @ 0.5 mg/mL	2.52

Physics Invariant: f x d = 542.6395 GHz*A | Derived from hbar / (2 * m_H2O * r_OO)

Engine Output 4. *Curcumin (0.001 mg/mL) - Bitter, 26.98 GHz*

CODEX Resonance v10

Hans-Made Research Inc. - Physics-Informed Analysis Engine

Curcumin
Bitter

Gate Path	PHASE4_CURCUMINOID_BITTER(Curcuminoid)
CODEX Frequency	26.98 GHz
f x d	26.98 GHz x 20.12 A = 542.84 GHz*A
Molecular Weight	368.38 Da
Confidence	91.0%
BCS Intensity @ 0.001 mg/mL	8.81

Physics Invariant: f x d = 542.6395 GHz*A | Derived from hbar / (2 * m_H2O * r_OO)

Appendix A (cont.): Calibration Standard Engine Output

Engine Output 5. *Caffeine (0.05 mg/mL) - Bitter, 77.22 GHz*

Engine Output 6. *Quinine (0.04 mg/mL) - Bitter, 41.12 GHz*



Engine Output 7. *Denatonium Benzoate (0.5 mg/mL) - Bitter, 36.30 GHz*

CODEX Resonance v10

Hans-Made Research Inc. - Physics-Informed Analysis Engine

Denatonium Benzoate

Bitter

Gate Path	PHASE4_QUATERNARY_BITTER(Benzamide)
CODEX Frequency	36.30 GHz
f x d	36.30 GHz x 14.95 A = 542.68 GHz*A
Molecular Weight	325.48 Da
Confidence	92.0%
BCS Intensity @ 0.5 mg/mL	24.78

Physics Invariant: f x d = 542.6395 GHz*A | Derived from hbar / (2 * m_H2O * r_OO)

Figure 1. *CODEX Predicted vs RCFTR Measured Bitterness (Table 4)*



Figure 2. *Table 3 Calibration: CODEX vs RCFTR*



Appendix B: Prediction Comparison Charts

Figure 3. *Dose-Response Curves: CODEX vs RCFTR for All Four Test Compounds*



Figure 4. *Naringin Inverted Dose-Response (Sugar Shielding)*


Naringin Inverted Dose-Response
(Sugar Shielding: higher conc = lower bitterness)
RCFTR Measured
CODEX Predicted
Bitterness Score
4.00
3.75
3.50
3.25
3.00
2.75
2.50
2.25
2.00
10^{-2}
10^{-1}
Concentration (mg/mL)

Appendix C: RCFTR Instrument Output

Figure 5. *RCFTR Bitterness Standard Curve (R^2 = 0.9413)*

Statistics report

Model supervisor ACCESS LEVEL — *Model creation date:* 3/4/2026 3:43:41 P M

Instrument :
Astree *Ref* AST-043
AM *Ref* 1824043006130

Library:
Codex Resonance pks excluded.lbx
Method:

Preprocessing method
AverageValue Time1 = 100 Time2 = 120
NoPretrait

Data processing method Sensory score correlation



Sample : B0_5_137 Sample type : Calibration Date : 2/27/2026 7:49:24 PM Reference : 23.00 Measured : 21.46, Indice

Selected sensor : AHS CTS NMS CPS ANS SCS

Removed Samples / outliers : Chlorogenic_Acid_1_148 Chlorogenic_Acid_1_204 Chlorog enic_Acid_1_260 Chlorogenic_Acid_2_157 Chlorogenic_Acid_2_213 Chlo rogenic_Acid_2_269 Chlorogenic_Acid_3_165 Chlorogenic_Acid_3_221 Chlorogenic_Acid_3_277 Curcumin_1_1 51 Curcumin_1_207 Curcumin_1_263 Curcumin_2_159 Curcumin_2_215 Curcumin_2_271 Curcumin_ 3_167 Curcumin_3_223 Curcumin_3_279

Signature reason	Signature name	Signature date

Note:

Visa:

User: ACCESS LEVEL — *Date:* 3/4/2026 3:44:02 P M — *Page* 1

UNIVERSITY OF MANITOBA ☎ XXX XXXXXX XXXXX

Figure 6. *PCA Dilution No. 1 (Discrimination Index = 95, PC1 = 84.6%, PC2 = 12.7%)*

Statistics report

Model supervisor ACCESS LEVEL *Model creation date:* 3/3/2026 9:01:04 A M

Instrument :
Astree *Ref* AST-043
AM *Ref* 1824043006130

Library: **Codex Resonance calibration.lbx**
Method:

Preprocessing method
AverageValue Time1 = 100 Time2 = 120
NoPretrait

Data processing method PCA

Data reduction : No



Sample : B0_5_249 Sample type : Calibration Date : 2/27/2026 11:18:01 PM PC1 - 84.646% : -167.90 PC2 - 12.667% : 1,550.24, Indice

Selected sensor : AHS CTS NMS CPS ANS SCS

Removed Samples / outliers : Chlorogenic_Acid_2_157 Chlorogenic_Acid_2_213 Chlorog enic_Acid_2_269 Chlorogenic_Acid_3_165 Chlorogenic_Acid_3_221 Chlo rogenic_Acid_3_277 Curcumin_2_159 Curcumin_2_215 Curcumin_2_271 Curcumin_3_167 Curcumin_3_223 C urcumin_3_279 Naringin_2_153 Naringin_2_209 Naringin_2_265 Naringin_3_161 Naringin_3_217 Naringin_ 3_273 Quercetin_2_155 Quercetin_2_211 Quercetin_

Signature reason	Signature name	Signature date

Note:

Visa:

User: ACCESS LEVEL *Date:* 3/3/2026 9:01:26 A M *Page* 1

UNIVERSITY OF MANITOBA ☎ XXX XXXXXX XXXXX

Figure 7. *PCA Dilution No. 2 (Discrimination Index = 96, PC1 = 79.6%, PC2 = 17.9%)*

Statistics report

Model supervisor ACCESS LEVEL *Model creation date:* 3/3/2026 9:01:48 A M

Instrument :
Astree *Ref* AST-043
AM *Ref* 1824043006130

Library:
Codex Resonance calibration.lbx
Method:

Preprocessing method
AverageValue Time1 = 100 Time2 = 120
NoPretrait

Data processing method PCA

Data reduction : No



Selected sensor : AHS CTS NMS CPS ANS SCS

Removed Samples / outliers : Chlorogenic_Acid_1_148 Chlorogenic_Acid_1_204 Chlorog enic_Acid_1_ 260 Chlorogenic_Acid_3_165 Chlorogenic_Acid_3_221 Chlo rogenic_Acid_3_277 Curcumin_1_151 Curcumin_1_ 207 Curcumin_1_263 Curcumin_3_167 Curcumin_3_223 C urcumin_3_279 Naringin_1_142 Naringin_1_198 Naringin_1_254 Naringin_2_153 Naringin_2_209 Naringin_ 2_265 Naringin_3_161 Naringin_3_217 Naringin_3_

Signature reason	Signature name	Signature date

Note:

Visa:

User: ACCESS LEVEL *Date:* 3/3/2026 9:02:04 A M *Page* 1

UNIVERSITY OF MANITOBA ☎ XXX XXXXXX XXXXX

Figure 8. *PCA Dilution No. 3 (Discrimination Index = 95, PC1 = 67.3%, PC2 = 25.8%)*

Statistics report

Model supervisor ACCESS LEVEL *Model creation date:* 3/3/2026 9:03:26 A M

Instrument :
Astree *Ref* AST-043
AM *Ref* 1824043006130

Library:
Codex Resonance calibration.lbx
Method:

Preprocessing method
AverageValue Time1 = 100 Time2 = 120
NoPretrait

Data processing method PCA

Data reduction : No



Sample : CAFFEINE_252 Sample type : Calibration Date : 2/27/2026 11:23:25 PM PC1 - 67.277% : -1,072.97 PC2 - 25.817% : 612.44, Indice

Selected sensor : AHS CTS NMS CPS ANS SCS

Removed Samples / outliers : Chlorogenic_Acid_1_148 Chlorogenic_Acid_1_204 Chlorog enic_Acid_1_260 Chlorogenic_Acid_2_157 Chlorogenic_Acid_2_213 Chlo rogenic_Acid_2_269 Curcumin_1_151 Curcumin_1_207 Curcumin_1_263 Curcumin_2_159 Curcumin_2_215 C urcumin_2_271 Naringin_1_142 Naringin_1_198 Naringin_1_254 Naringin_2_153 Naringin_2_209 Naringin_ 2_265 Quercetin_1_145 Quercetin_1_201 Quercetin_

Signature reason	Signature name	Signature date

Note:

Visa:

User: ACCESS LEVEL *Date:* 3/3/2026 9:03:52 A M *Page* 1

UNIVERSITY OF MANITOBA ☎ XXX XXXXXX XXXXX