

Projective Geometry

Tejaswi

Advisor: Dr. Steven Spallone

Summer 2025

Contents

1	Some Algebra	1
1.1	Groups, Fields, and Rings	1
1.2	Field Extensions	1
2	Conics	2
2.1	Dandelin Spheres	2
2.2	Group Laws on Conics	4
2.3	Generating solutions for algebraic equations	8
3	Affine Geometry	9
3.1	Affine Space	9
3.2	Fundamental Theorem of Affine Geometry	9
3.3	Affine Congruence of Conics	9
4	Projective Geometry	10
4.1	The Projective Space	10
4.2	Fundamental Theorem of Projective Geometry	10
4.3	Some Theorems	10
4.4	Group Laws on Elliptic Curves	10

Chapter 1

Some Algebra

1.1 Groups, Fields, and Rings

1.2 Field Extensions

Chapter 2

Conics

Definition. *A conic section, a conic, or a quadratic curve is a curve obtained from a cone's surface intersecting a plane.*

2.1 Dandelin Spheres

Germinal Pierre Dandelin, a 19th century French-Belgian Professor, discovered this beautiful proof to demonstrate that any plane that cuts through a right circular cone produces a quadratic curve.

Theorem. *When a plane intersects a right circular cone, the curve produced will either be an ellipse, a parabola or a hyperbola.*

Proof. Place a sphere tangent to the intersecting plane π and the cone such that it touches the plane at F , and the cone in a circle C with centre O that lies on a horizontal plane ϵ .¹

Take an arbitrary point P on the curve Q , and extend the line VP from the vertex V of the cone to meet C at point L . Let D be the point on the intersection on the planes π and ϵ such that PD is perpendicular to the line of intersection. (If the planes do not intersect, Q will be a circle)

Drop a perpendicular PM on OL such that $\triangle PML$ and $\triangle PMD$ are both right angled. Denote $\angle PLM$ as α , and $\angle PDM$ as β .

¹Assuming such sphere exists.

From the triangles $\triangle PML$ and $\triangle PMD$

$$\begin{aligned}\sin \alpha &= \frac{PM}{PD} \\ \text{and } \sin \beta &= \frac{PM}{PL} \\ \text{i.e. } \frac{PL}{PD} &= \frac{\sin \alpha}{\sin \beta}\end{aligned}$$




Figure 2.1: When $0 < \alpha < \beta < \frac{\pi}{2}$

Since PL and PF are both tangents from P to the sphere, $PF = PL$. Therefore,

$$\frac{PF}{PD} = \frac{\sin \alpha}{\sin \beta}$$

i.e. $PF = e \cdot PD$, where $e = \sin \alpha / \sin \beta$

It follows from the focus - directrix definition that Q will be an ellipse if $\alpha < \beta$, a parabola if $\alpha = \beta$, or a hyperbola if $\alpha > \beta$. \square

Proof adapted from [BEG12] with modifications to generalize it for all conics.

2.2 Group Laws on Conics

Consider a conic section C and a point $O \in C$. For any $P, Q \in C$, define a binary operation $\oplus : C \times C \rightarrow C$ by $P \oplus Q = R$, where R is such that $l_{PQ} \parallel l_{OR}$.

Theorem. *Set of points of C forms a group $G(C)$ under the binary operation \oplus , with O as the identity element.*

Proof. Closure: The line through O parallel to l_{PQ} necessarily meets C again, (counting algebraic multiplicities) since for any quadratic equation with real coefficients, if one of the roots is real, the other one must be real too.

Existence of Identity Element: The point O serves as the identity element.

Existence of Inverse: Constructively, when Q is such that the line parallel to l_{PQ} that passes through O is tangent to the conic, i.e. when $R = O$, we get $P \oplus Q = O$. So, Q serves as the inverse of P .

Associativity: To prove associativity, we'll find algebraic formula for $P \oplus Q$ for standard conics, i.e. for the circle $x^2 + y^2 = 1$, for the parabola $y = x^2$, and for the hyperbola $xy = 1$. In the next chapter, we'll prove that any ellipse, hyperbola or parabola is affine congruent to its standard form. This result will generalize the result to all conics. The following formulae will be valid for any fields with non-two characteristic.

Let the point P be (p_1, p_2) , Q be (q_1, q_2) , O be (o_1, o_2) , and R be (r_1, r_2) , and let the slope of the line l_{PQ} be $\lambda = (q_2 - p_2) / (q_1 - p_1)$, assuming $P \neq Q$, since associativity would be trivial then. Let ℓ be the line through O with slope λ . The coordinates of R will satisfy $\lambda = \frac{r_2 - o_2}{r_1 - o_1} = \frac{q_2 - p_2}{q_1 - p_1}$, $\Rightarrow r_2 = o_2 + \mu(q_2 - p_2)$ and $r_1 = o_1 + \mu(q_1 - p_1)$ for some $\mu \in \mathbb{F}$.

(i) Circle

Without loss of generality, let $O = (1, 0)$. Since R also lies on C , $r_1^2 + r_2^2 = 1$. i.e.

$$\begin{aligned}
& (1 + \mu(q_1 - p_1))^2 + (0 + \mu(q_2 - p_2))^2 = 1 \\
\Rightarrow & \mu(\mu(q_1 - p_1)^2 + \mu(q_2 - p_2)^2 + 2(q_1 - p_1)) = 0 \\
\Rightarrow & \mu = 0 \text{ or } \mu = -\frac{2(q_1 - p_1)}{(q_1 - p_1)^2 + (q_2 - p_2)^2}
\end{aligned}$$

We assume that $(q_1 - p_1)^2 + (q_2 - p_2)^2 \neq 0$. Because if it was so,

$$\begin{aligned}
& q_1^2 + p_1^2 - 2q_1p_1 + q_2^2 + p_2^2 - 2p_2q_2 = 0 \\
\Rightarrow & 1 - p_1q_1 - p_2q_2 = 0 \\
\Rightarrow & p_1^2q_1^2 = 1 + p_2^2q_2^2 - 2p_2q_2 \\
\Rightarrow & p_1^2q_1^2 = 1 + (1 - p_1^2)(1 - q_1^2) - 2p_2q_2 \\
\Rightarrow & 0 = 2 - p_1^2 - q_1^2 - 2p_2q_2 \\
\Rightarrow & (p_2 - q_2)^2 = 0 \\
\Rightarrow & p_2 = q_2 \text{ and similarly, } p_1 = q_1
\end{aligned}$$

Which is when $P = Q$, which we have assumed not to be true.




Figure 2.2: $R = P \oplus Q$ when C is a circle.

The $\mu = 0$ solution corresponds to O . Considering the other solution,

$$\begin{aligned}
r_1 &= 1 - \frac{2(q_1 - p_1)^2}{(q_1 - p_1)^2 + (q_2 - p_2)^2} \\
&= \frac{(q_2 - p_2)^2 - (q_1 - p_1)^2}{(q_1 - p_1)^2 + (q_2 - p_2)^2} \\
&= \frac{q_2^2 + p_2^2 - 2p_2q_2 - q_1^2 - p_1^2 + 2p_1q_1}{2(1 - p_1q_1 - p_2q_2)} \\
&= \frac{1 - p_1^2 - q_1^2 + p_1q_1 - p_2q_2}{1 - p_1q_1 - p_2q_2} \\
&= \frac{(p_1q_1 - p_2q_2)(1 - p_1q_1 - p_2q_2)}{1 - p_1q_1 - p_2q_2} \\
&= p_1q_1 - p_2q_2 \\
\text{and, } r_2 &= -\frac{2(q_1 - p_1)(q_2 - p_2)}{(q_1 - p_1)^2 + (q_2 - p_2)^2} \\
&= \frac{p_2q_2 + p_2q_1 - p_1p_2 - q_1q_2}{1 - p_1q_1 - p_2q_2} \\
&= \frac{(p_1q_2 + p_2q_1)(1 - p_1q_1 - p_2q_2)}{1 - p_1q_1 - p_2q_2} \\
&= p_1q_2 + p_2q_1
\end{aligned}$$

$$\Rightarrow R = P \oplus Q = (r_1, r_2) = (p_1q_1 - p_2q_2, p_1q_2 + p_2q_1)$$

Using this formula, it can be proved that $(P \oplus Q) \oplus R = P \oplus (Q \oplus R)$.

(ii) **Parabola**

Without loss of generality, let $O = (0, 0)$. The points of the standard parabola can be parameterized as (t, t^2) . Let $P = (p, p^2)$, $Q = (q, q^2)$, and $R = (r, r^2)$. Substituting these in λ ,

$$\begin{aligned}
\lambda &= \frac{r^2}{r} = \frac{q^2 - p^2}{q - p} \Rightarrow r = p + q \\
&\Rightarrow P \oplus Q = (p + q, (p + q)^2)
\end{aligned}$$

Since the parameters just get added, it can be easily proved that $P \oplus (Q \oplus R) = (P \oplus Q) \oplus R$




Figure 2.3: $R = P \oplus Q$ when C is a parabola.

(iii) **Hyperbola**

Without loss of generality, let $O = (1, 1)$. The points of the standard hyperbola can be parameterized as $(t, \frac{1}{t})$. Let $P = (p, \frac{w}{p})$, $Q = (q, \frac{1}{q})$, and $R = (r, \frac{1}{r})$. Substituting these in λ ,

$$\begin{aligned} \lambda &= \frac{\frac{1}{r} - 1}{r - 1} = \frac{\frac{1}{q} - \frac{1}{p}}{p - q} \Rightarrow r = pq \\ \Rightarrow P \oplus Q &= (pq, \frac{1}{pq}) \end{aligned}$$




Figure 2.4: $R = P \oplus Q$ when C is a hyperbola.

Since parameters just get multiplied, it can be easily proved that
 $P \oplus (Q \oplus R) = (P \oplus Q) \oplus R$

□

Proof adapted from [Shi09] with a formula based field independent proof for associativity.

2.3 Generating solutions for algebraic equations

Chapter 3

Affine Geometry

3.1 Affine Space

A set ε is endowed with the structure of an affine space by a vector space E and a mapping Θ that associates a vector of E with any ordered pair of points in ε ,

$$\begin{array}{ccc} \varepsilon \times \varepsilon & \longrightarrow & E \\ (A, B) & \longmapsto & \overrightarrow{AB} \end{array}$$

such that:

- for any point A of ε , the partial map $\Theta_A : B \mapsto \overrightarrow{AB}$ is a bijection from ε to E .
- for any points A , B , and C in ε , we have $\overrightarrow{AB} = \overrightarrow{AC} + \overrightarrow{CB}$.

The vector space E is the direction of ε , or its underlying vector space. The elements of ε are called points, and the dimension of the vector space E is called the dimension of ε . [Aud02]

3.2 Fundamental Theorem of Affine Geometry

3.3 Affine Congruence of Conics

Chapter 4

Projective Geometry

4.1 The Projective Space

4.2 Fundamental Theorem of Projective Geometry

4.3 Some Theorems

Theorem (Desargues's Theorem).

Theorem (Pascal's Theorem).

4.4 Group Laws on Elliptic Curves

An elliptic curve is a non-empty, non-singular, degree 3 projective curve. [Spa]

Bibliography

- [Aud02] Michele Audin. *Geometry*. Universitext. Springer-Verlag, 2002.
- [BEG12] David A. Brannan, Matthew F. Esplen, and Jeremy J. Gray. *Geometry*. Cambridge University Press, second edition, 2012.
- [Shi09] Shailesh Shirali. Groups associated with conics. *The Mathematical Gazette*, March 2009.
- [Spa] Steven Spallone. Introduction to curves.