

Inverse Trigonometric Function Derivatives

Logan Grosz

October 21, 2017

Abstract

The derivatives of inverse trigonometric derivatives are very different than their basic trigonometric counterparts. The following document will describe the process of getting such a visually unorthodox solution to $\frac{d}{dx} f$, where f is an inverse trigonometric function.

1 Declarations

x ; parameter passed into inverse trigonometric function, measured in radians;

For $\sin^{-1} x, x \in [-1, 1]$

For $\cos^{-1} x, x \in [-1, 1]$

For $\tan^{-1} x, x \in [-\infty, \infty]$

For $\cot^{-1} x, x \in [-\infty, \infty]$

For $\sec^{-1} x, x \in (-\infty, -1] \cup [1, \infty)$

For $\csc^{-1} x, x \in (-\infty, -1] \cup [1, \infty)$

$\theta \triangleq x$; used in place of x at times to not confused reader;

$\arcsin f \triangleq f^{-1}$; arc and -1 convention equivalence; $\arcsin x \equiv \sin^{-1} x$

hypotenuse; Latin for "stretching under", the line connecting both catheti; Short: *hyp*

adjacent; Cathetus closest or next to the observed angle; Short: *adj*

opposite Cathetus furthest from or opposite of the observed angle; Short: *opp*

2 Rule

$$\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}} \quad (1)$$

$$\frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1-x^2}} \quad (2)$$

$$\frac{d}{dx} \tan^{-1} x = \frac{1}{x^2+1} \quad (3)$$

$$\frac{d}{dx} \cot^{-1} x = \frac{-1}{x^2+1} \quad (4)$$

$$\frac{d}{dx} \sec^{-1} x = \frac{1}{|x|\sqrt{x^2-1}} \quad (5)$$

$$\frac{d}{dx} \csc^{-1} x = \frac{-1}{|x|\sqrt{x^2-1}} \quad (6)$$

3 Pre-Derivation


The following notation is the inverse of its subscribed function:

$$\sin^{-1}(\sin \theta) = \theta$$







Pythagorean Trigonometric Identity:

$$\sin^2 \theta + \cos^2 \theta = 1$$

The following defines sides of triangle given angle measure θ .



The following are the graphs of all six inverse trigonometric functions for reference:



4 Derivation

4.1 Inverse Sine

$$y = \sin^{-1} x$$

$$\sin y = x$$

$$\frac{d}{dx} \sin y = \frac{d}{dx} x$$

$$\frac{dy}{dx} \cos y = 1$$

$$\frac{dy}{dx} = \frac{1}{\cos y}$$

Using $\sin^2 y + \cos^2 y = 1 \dots$


$$\cos y = \sqrt{1 - \sin^2 y}$$

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - \sin^2 y}}$$

Recall $\sin y = x \dots$

$$\therefore \frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1 - x^2}}$$

Id Est



$$\sin y = x \Rightarrow \frac{\text{opp}}{\text{hyp}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{1}{\cos x}$$

See figure at left...

$$\cos y = \sqrt{1 - x^2}$$

$$\therefore \frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1 - x^2}}$$

4.2 Inverse Cosine

$$y = \cos^{-1} x$$

$$\cos y = x$$

$$\frac{d}{dx} \cos y = \frac{d}{dx} x$$

$$\frac{dy}{dx} - \sin y = 1$$

$$\frac{dy}{dx} = \frac{-1}{\sin y}$$

Using $\sin^2 y + \cos^2 y = 1 \dots$


$$\sin y = \sqrt{1 - \cos^2 y}$$

$$\frac{dy}{dx} = \frac{-1}{\sqrt{1 - \cos^2 y}}$$

Recall $\cos y = x \dots$

$$\therefore \frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1 - x^2}}$$

Id Est



$$\cos y = x \Rightarrow \frac{\text{adj}}{\text{hyp}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{-1}{\sin y}$$

See figure at left...

$$\sin y = \sqrt{1 - x^2}$$

$$\therefore \frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1 - x^2}}$$

4.3 Inverse Tangent

$$y = \tan^{-1} x$$

$$\tan y = x$$

$$\frac{d}{dx} \tan y = \frac{d}{dx} x$$

$$\frac{dy}{dx} \sec^2 y = 1$$

$$\frac{dy}{dx} = \frac{1}{\sec^2 y}$$

Using $\sin^2 y + \cos^2 y = 1 \dots$


$$\sec^2 y = \tan^2 y + 1$$

$$\frac{dy}{dx} = \frac{1}{\tan^2 y + 1}$$

Recall $\tan y = x \dots$

$$\therefore \frac{d}{dx} \tan^{-1} x = \frac{1}{x^2 + 1}$$

Id Est



$$\tan y = x \Rightarrow \frac{\text{opp}}{\text{adj}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{1}{\sec^2 y}$$

See figure at left...

$$\sec^2 y = x^2 + 1$$

$$\therefore \frac{d}{dx} \tan^{-1} x = \frac{1}{x^2 + 1}$$

4.4 Inverse Cotangent

$$y = \cot^{-1} x$$

$$\cot y = x$$

$$\frac{d}{dx} \cot y = \frac{d}{dx} x$$

$$\frac{dy}{dx} - \csc^2 y = 1$$

$$\frac{dy}{dx} = \frac{-1}{\csc^2 y}$$

Using $\sin^2 y + \cos^2 y = 1$...


$$\csc^2 y = \cot^2 y + 1$$

$$\frac{dy}{dx} = \frac{-1}{\cot^2 y + 1}$$

Recall $\cot y = x$...

$$\therefore \frac{d}{dx} \cot^{-1} x = \frac{-1}{x^2 + 1}$$

Id Est



$$\cot y = x \Rightarrow \frac{\text{adj}}{\text{opp}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{-1}{\csc^2 y}$$

See figure at left...

$$\csc^2 y = x^2 + 1$$

$$\therefore \frac{d}{dx} \cot^{-1} x = \frac{-1}{x^2 + 1}$$

4.5 Inverse Secant

$$y = \sec^{-1} x$$


$$\sec y = x$$

$$\frac{d}{dx} \sec y = \frac{d}{dx} x$$

$$\frac{dy}{dx} \sec y \tan y = 1$$

$$\frac{dy}{dx} = \frac{1}{\sec y \tan y}$$

Id Est



$$\sec y = x \Rightarrow \frac{\text{hyp}}{\text{adj}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{1}{\sec y \tan y}$$

See figure at left...

$$\tan y = \sqrt{x^2 - 1}$$

Recall $\sec y = x$


And note $\frac{d}{dx} \sec^{-1} \theta > 0$, so, $|x| = \{x | x > 0\}$

$$\therefore \frac{d}{dx} \sec^{-1} x = \frac{1}{|x| \sqrt{x^2 - 1}}$$

4.6 Inverse Cosecant

$$\begin{aligned}
 y &= \csc^{-1} x \\
 \csc y &= x \\
 \frac{d}{dx} \csc y &= \frac{d}{dx} x \\
 \frac{dy}{dx} - \csc y \cot y &= 1 \\
 \frac{dy}{dx} &= \frac{1}{\csc y \cot y}
 \end{aligned}$$

Id Est



$$\csc y = x \Rightarrow \frac{\text{hyp}}{\text{opp}} = \frac{x}{1}$$

See above that...

$$\frac{dy}{dx} = \frac{-1}{\csc y \cot y}$$

See figure at left...

$$\cot y = \sqrt{x^2 + 1}$$

Recall $\csc y = x$

And note, $\frac{d}{dx} \csc^{-1} \theta < 0$, so, $|x| = \{x | x > 0\}$

$$\therefore \frac{d}{dx} \csc^{-1} x = \frac{-1}{|x| \sqrt{x^2 - 1}}$$

5 Exempli Gratia

Examples of important instances