

# Calibration Report: High N Sedimentary Site Base Case

Kaveh Gholamhossein Siah

21 March 2021

# Contents

Hydrology . . . . .	4
Soil Solution Results . . . . .	4
Lysimeter Comparisons . . . . .	10
Weathering Results . . . . .	13
Litter Pool Results . . . . .	16
Soil Organic Matter Results . . . . .	18
Tree Nutrient Content . . . . .	21
Analysis 1: Stack Flux Data . . . . .	22
Cation Exchange Capacity . . . . .	25
Anion Exchange Capacity . . . . .	26
Other . . . . .	28


# List of Figures

1 Monthly Calcium Concentrations by Soil Layer . . . . .	6
2 Monthly Magnesium Concentrations by Soil Layer . . . . .	6
3 Monthly Potassium Concentrations by Soil Layer . . . . .	6
4 Monthly Sodium Concentrations by Soil Layer . . . . .	7
5 Monthly Sulfate Concentrations by Soil Layer . . . . .	7
6 Monthly Chloride Concentrations by Soil Layer . . . . .	7
7 Monthly Aluminum Concentrations by Soil Layer . . . . .	8
8 Monthly SiO <sub>2</sub> Concentrations by Soil Layer . . . . .	8
9 Monthly Organic Acid Base (R-) Concentrations by Soil Layer . . . . .	8
10 Monthly pH by Soil Layer . . . . .	9
11 Yearly Ammonium concentration by Soil Layer . . . . .	9
12 Yearly Nitrate concentration by Soil Layer . . . . .	9
13 Calcium Weathering (All Layer) . . . . .	13
14 Magnesium Weathering (All Layer) . . . . .	13
15 Potassium Weathering (All Layer) . . . . .	14
16 Aluminum Weathering (All Layer) . . . . .	14
17 Phosphate Weathering (All Layer) . . . . .	15
18 Silica Weathering (All Layer) . . . . .	15
19 Sodium Weathering (All Layer) . . . . .	16
20 Forest Floor (O-Layer) Carbon Content Over Simulation Period . . . . .	16
21 Forest Floor (O-Layer) Carbon Content Over Simulation Period . . . . .	17
22 Forest Floor/O-horizon Ca content over time (a). and net annual Ca return in litterfall (b). . . . .	17
23 Forest Floor/O-horizon Mg content over time (a). and net annual Mg return in litterfall (b). . . . .	18
24 Forest Floor/O-horizon K content over time (a). and net annual K return in litterfall (b). . . . .	19
25 Forest Floor/O-horizon S content over time (a). and net annual S return in litterfall (b). . . . .	19
26 Forest Floor/O-horizon P content over time (a). and net annual P return in litterfall (b). . . . .	20
27 Forest Floor/O-horizon N content over time (a). and net annual N return in litterfall (b). . . . .	20
28 Base Cation Nutrient Content in Simulated Forest . . . . .	21
29 N, S, and P Nutrient Contents and biomass per compartment . . . . .	22
30 Calcium input and output comparison graphs . . . . .	22
31 Magnesium input and output comparison graphs . . . . .	23
32 Potassium input and output comparison graphs . . . . .	23
33 Sulfur input and output comparison graphs . . . . .	24
34 Nitrogen input and output comparison graphs . . . . .	24
35 Calcium and Magnesium on exchangerover time . . . . .	25
36 Potassium and Sodium on exchangerover time . . . . .	25
37 Ammonium and Aluminum on exchangerover time . . . . .	25
38 N and P Potential Uptake to Actual Uptake Difference . . . . .	28
39 Ca and Mg Potential Uptake to Actual Uptake Difference . . . . .	29
40 K and S Potential Uptake to Actual Uptake Difference . . . . .	29

## List of Tables

1	Lysimeter Measured Soil Solution Concentrations of Reliable Months (2005) . . . . .	5
2	Simulated Lysimeter Fluxes by Depth (2005-2006) . . . . .	11
3	Actual Average Lysimeter Fluxes (2005) . . . . .	12

## Hydrology



## Soil Solution Results

Table 1: Lysimeter Measured Soil Solution Concentrations of Reliable Months (2005)

Layer	\$\\mu\$mol/L																								
	Ca	Ca SD	Mg	Mg SD	K	K SD	Na	Na SD	NO3	NO3 SD	NH4	NH4 SD	SO4	SO4 SD	Cl	Cl SD	P <sup>a</sup>	P SD	DOC	DOC SD	Al <sup>b</sup>	Al SD	Si <sup>c</sup>	Si SD	pH <sup>d</sup>
1	24	4.97	45	12.1	2.3	0.76	237	30	46	15	1.1	0.32	20	4.0	271	77	0.037	0.93	106	13	2.76	0.856	20	4.6	4.5
2	17	0.78	35	2.3	2.4	1.16	226	24	29	12	1.0	0.20	22	1.7	218	22	0.058	1.98	38	13	0.63	0.091	25	7.6	6.3

<sup>a</sup> Average based on TP annual average

<sup>b</sup> Does not distinguish between organic-Al and free Al

<sup>c</sup> Model does not simulate Si uptake

<sup>d</sup> From Hynicka et al., 2017 (10-50cm) extrapolated to 1m




Figure 1: Monthly Calcium Concentrations by Soil Layer




Figure 2: Monthly Magnesium Concentrations by Soil Layer




Figure 3: Monthly Potassium Concentrations by Soil Layer




Figure 4: Monthly Sodium Concentrations by Soil Layer




Figure 5: Monthly Sulfate Concentrations by Soil Layer




Figure 6: Monthly Chloride Concentrations by Soil Layer




Figure 7: Monthly Aluminum Concentrations by Soil Layer




Figure 8: Monthly  $\text{SiO}_2$  Concentrations by Soil Layer




Figure 9: Monthly Organic Acid Base (R-) Concentrations by Soil Layer




Figure 10: Monthly pH by Soil Layer




Figure 11: Yearly Ammonium concentration by Soil Layer




Figure 12: Yearly Nitrate concentration by Soil Layer

## Lysimeter Comparisons





Table 2: Simulated Lysimeter Fluxes by Depth (2005-2006)

Depth	YEAR	kg/ha											
		Ca	Mg	K	Na	NO3	NH4	SO4	Cl	P	DOC	Al	Si
2	2005	14.9	16	1.37	35	14	0.99	10.5	109	0.131	17	0.77	8.7
2	2006	9.6	11	0.98	18	11	0.69	8.9	31	0.089	17	0.65	6.5
8	2005	10.8	13	1.2	60	6.5	0.90	9.5	93	0.0182	12	0.059	15.7
8	2006	6.9	10	1.0	34	4.3	0.34	9.3	42	0.0018	12	0.074	7.2

Table 3: Actual Average Lysimeter Fluxes (2005)

Depth	kg/ha												
	NH4	NO3	TN	DOC	TP	Cl	SO4	Ca	Mg	K	Na	Al	Si
20	0.247777178	9.763424509	16.05235558	16.52820633	0.015397959	147.7048599	7.135287478	14.6687802	16.66489783	1.089048305	84.64727009	1.026315671	15.44797242
100	0.207849949	5.757736987	11.92369571	10.69157732	0.024684756	105.034089	6.944387278	9.669335876	11.42146102	1.013541339	70.18746399	0.216742935	20.16864975

## Weathering Results




Figure 13: Calcium Weathering (All Layer)




Figure 14: Magnesium Weathering (All Layer)




Figure 15: Potassium Weathering (All Layer)




Figure 16: Aluminum Weathering (All Layer)




Figure 17: Phosphate Weathering (All Layer)




Figure 18: Silica Weathering (All Layer)




Figure 19: Sodium Weathering (All Layer)

### Litter Pool Results






Figure 20: Forest Floor (O-Layer) Carbon Content Over Simulation Period

Looking at a range of soil carbon studies in Douglas-fir forests of the Pacific Northwest, forest floor (defined as non-mineral OM) C content goes from a lower bound of 3,700 kg C/ha in a 9-yr old stand (Cromack et al., 1999) to 8200 kg C/ha in an average 38 year old stand (Edmonds and Chappell, 1993). These stands were notably N rich compared to the site simulated for the low N site, the soil C should be lower in the simulations as there is about half as much soil N in the low N simulated site as in the sites described in Edmonds and Chappel, 1993. The high N site has about 21,000 kg N/ha at 1m depth, so it should be modeled to be at the higher end of organic and litter C buildup.




(a) Carbon content of fine litter (kg/ha)




(b) Carbon content of humic litter (kg/ha)

Figure 21: Forest Floor (O-Layer) Carbon Content Over Simulation Period

Note that the fine litter pool (the stage between humus and fresh/coarse litter) is growing in this model. This might deviate from observed behavior.




(a) Ca Content in each litter decomposition stage (kg/ha)



(b) Ca net annual return in litterfall (kg/ha)

Figure 22: Forest Floor/O-horizon Ca content over time (a). and net annual Ca return in litterfall (b).

## Soil Organic Matter Results



Mineral soil SOM C content is very high compared to other pools of carbon in the ecosystem, soil carbon should buildup over time assuming available surfaces exist for soil carbon “stabilization”. In NutsFor, the SOM pool is represented by an active microbial pool, so there are issues with building up SOM in the soil as one might expect from a real stand. Microbial growth is limited by soil moisture and nutrient availability like the tree pool, so it is not a wholly adequate representation of C stabilization. Instead of calibrating this output to show buildup, I calibrated it such that it was “level”, thus, soil carbon additions to the mineral soil are dictated by DOC percolation with water flow.






Figure 23: Forest Floor/O-horizon Mg content over time (a). and net annual Mg return in litterfall (b).



(a) K Content in each litter decomposition stage (kg/ha)


Litter Type

- Coarse Litter
- Fine Litter
- Humic Litter
- Total Litter



(b) K net annual return in litterfall (kg/ha)


Figure 24: Forest Floor/O-horizon K content over time (a). and net annual K return in litterfall (b).



(a) S Content in each litter decomposition stage (kg/ha)


Litter Type

- Coarse Litter
- Fine Litter
- Humic Litter
- Total Litter



(b) S net annual return in litterfall (kg/ha)


Figure 25: Forest Floor/O-horizon S content over time (a). and net annual S return in litterfall (b).



(a) P Content in each litter decomposition stage (kg/ha)


Litter Type

- Coarse Litter
- Fine Litter
- Humic Litter
- Total Litter



(b) P net annual return in litterfall (kg/ha)


Figure 26: Forest Floor/O-horizon P content over time (a). and net annual P return in litterfall (b).



(a) N Content in each litter decomposition stage (kg/ha)

Litter Type

- Coarse Litter
- Fine Litter
- Humic Litter
- Total Litter



(b) N net annual return in litterfall (kg/ha)

Figure 27: Forest Floor/O-horizon N content over time (a). and net annual N return in litterfall (b).

## Tree Nutrient Content







Figure 28: Base Cation Nutrient Content in Simulated Forest




(a) Nitrogen content in each biomass compartment



(b) Sulfur content in each biomass compartment




(c) Phosphorous content in each biomass compartment




(d) Biomass of each compartment

Figure 29: N, S, and P Nutrient Contents and biomass per compartment

### Analysis 1: Stack Flux Data




(a) Ca input and output fluxes over time




(b) Total Average Ca input and output fluxes

Figure 30: Calcium input and output comparison graphs




(a) Mg input and output fluxes over time




(b) Total Average Mg input and output fluxes

Figure 31: Magnesium input and output comparison graphs



(a) K input and output fluxes over time




(b) Total Average K input and output fluxes

Figure 32: Potassium input and output comparison graphs




(a) S input and output fluxes over time




(b) Total Average S input and output fluxes

Figure 33: Sulfur input and output comparison graphs

The sulfate adsorbed pool depletes itself, the organic sulfur pool becomes increasingly dominant. This behavior is not unreasonable, however I would expect higher sulfate adsorption.



(a) N input and output fluxes over time



(b) Total Average N input and output fluxes

Figure 34: Nitrogen input and output comparison graphs

Notice how SOM mineralization starts off highly negative (-500+ kg/ha/yr N). This is strange as the SOM pool is set such that it starts with the same concentration of N (roughly a C:N of 18) as is entered for the microbial target concentration. It does not make sense that so much N would be initially uptaken into the SOM pool, this could be a coding error in NutsFor or an issue with initial parameter entry.

## Cation Exchange Capacity




Figure 35: Calcium and Magnesium on exchangerover time




Figure 36: Potassium and Sodium on exchangerover time







Figure 37: Ammonium and Aluminum on exchangerover time

### Anion Exchange Capacity





The phosphate adsorption is set from the original parameterization I received from Gregory. It tends to build up, which implies a high soil solution concentration (adsorption is determined by concentration).

Sulfate adsorption is weak and drains easily, I set a low adsorbed sulfate pool following IFS data from the Thompson site (glacial outwash, inceptisol). According to the book *Atmospheric Sulfur Deposition: Environmental and Health Impacts*, sulfur is mostly locked in organic compartments rather than on the adsorption surfaces. We might expect that sulfate, like phosphate, would increase on the AEC, however the input of sulfate relative to the adsorption and uptake of sulfate is likely too low to facilitate adsorption. This is well supported by IFS data that show low sulfate adsorption on potentially high capacity adsorbing soils. The higher sulfate concentrations observed at the high N site could well be due to a higher inherent sulfur pool, possibly a condition of higher sulfate-mineral weathering, or due to a competitive response with phosphate.

## Other






Figure 38: N and P Potential Uptake to Actual Uptake Difference




Figure 39: Ca and Mg Potential Uptake to Actual Uptake Difference




Figure 40: K and S Potential Uptake to Actual Uptake Difference

For the high N sedimentary site, the current calibration clearly shows Ca and P limitation. P limitation is effectively the same as in the low N sedimentary site as I used the exact same adsorbed P pool initially, although the SOM contains more organically bound P than in the low N scenario. The additional organic P is the reason why a strong phosphate adsorption response is seen in this site, as the decomposition rate and P release rates are enough to stimulate high enough phosphate concentrations to stimulate adsorption.