




# Supporting Information

Toettcher et al. 10.1073/pnas.1005615107




**Fig. S1.** Population and single cells data of p53 dynamics after zinc stimulation. (A) Time-course of endogenous p53 and exogenous p53–CFP levels in MCF7 cells after addition of 25 or 50  $\mu\text{M}$  zinc. Cells were collected every hour and analyzed by Western blot (see *SI Appendix, “Estimating pulse statistics from trajectories”* for details regarding accounting for the endogenous protein level in our calculations of p53 pulse amplitude and damping). (B) Time-course of total p53 induction, quantified from the Western blot in A. All samples are normalized to tubulin and are shown relative to the value at  $t = 0$ . (C) Fraction of cells that divide within 24 h after zinc treatment. (D) Timing between successive p53 pulses, computed for the same cells shown in Fig. 1 *H–I*. Cells were treated with varying zinc doses; at least 50 cells were analyzed per condition. The fraction of cells observed undergoing each p53 pulse (E) or Mdm2 pulse (F) is plotted against pulse number from the same time-lapse microscopy experiments of Fig. 1 *F* and *G*. Colors representing each zinc dose are the same as shown in D.





**Fig. S3.** Predictions for addition of synthetic feedback loops on p53. Prediction of p53 damping (A, D), p53 amplitude (B, E), and p53 frequency coefficient (C, F) for the model incorporating a second negative feedback in addition to the core p53–Mdm2 loop (A–C; 2NF model) and a synthetic positive feedback loop in addition to the core p53–Mdm2 loop (D–F; NPF model). The x and y axes represent 51 logarithmically spaced values from 0.1 to 10 times the nominal parameter value for the MTF1 protein production delay  $\gamma_{t0}$  and synthetic feedback strength  $\beta_{p0}$ , respectively.



**Fig. S4.** Fraction of pulsing cells in the presence of additional synthetic feedback loops on p53. The number of cells observed undergoing each pulse are plotted against the number of pulses from the same single cell time-lapse microscopy experiments as Fig. 3 E and F.



**Table S1. Parameters for the core p53–Mdm2 negative feedback (1NF) model**

Parameter name	Parameter value	Description	Units
$\rho_z$	0.527	Zinc-mediated p53 production rate	1/(Zn h)
$\alpha_p$	1	p53 protein production rate	p53/h
$\delta_p$	16,388	Mdm2-mediated p53 degradation rate	1/h
$K_p$	3.568	Saturation of Mdm2-mediated p53 degradation	p53
$\alpha_{m0}$	0.069	Mdm2 mRNA basal transcription rate	<i>mdm2</i> /h
$\beta_{m0}$	0.058	p53-mediated Mdm2 mRNA transcription rate	<i>mdm2</i> /h
$K_{m0}$	0.117	Saturation of p53-mediated Mdm2 production	p53
$\gamma_{m0}$	1.854	Mdm2 production delay	1/h
$\gamma_m$	0	Mdm2 degradation rate	1/h
$\delta_m$	0.128	Mdm2-mediated Mdm2 degradation rate	1/h
$K_m$	$1.25 \times 10^{-4}$	Saturation of Mdm2-mediated Mdm2 degradation	Mdm2

The table indicates each parameter's name, its nominal value, a description of its effect, and the units in which it is measured. Italicized units are mRNA; otherwise units represent protein levels.

**Table S2. Parameters for the models with additional synthetic feedback loops on p53**

Parameter name	Parameter value	Description	Units
$\alpha_{p0}$	0.040	P53 mRNA basal transcription rate	1/(Zn h)
$\beta_{p0,NPF}$	14.82	NPF-mediated <i>p53</i> mRNA transcription rate	<i>p53</i> /h
$\beta_{p0,2NF}$	0.317	2NF-mediated <i>p53</i> mRNA transcription rate	<i>p53</i> /h
$K_f$	2	Saturation of feedback-mediated p53 production	FB
$\gamma_{p0}$	3.293	p53 production delay	1/h
$\alpha_p$	1	p53 production rate	p53/h
$\delta_p$	16,388	Mdm2-mediated p53 degradation rate	p53
$\alpha_{m0}$	0.069	<i>mdm2</i> mRNA basal transcription rate	1/h
$K_p$	3.568	Saturation of Mdm2-mediated p53 degradation	p53
$\beta_{m0}$	0.058	p53-mediated <i>mdm2</i> mRNA transcription rate	Mdm2/h
$K_{m0}$	0.117	Saturation of p53-mediated Mdm2 production	p53
$\gamma_{m0}$	1.854	Mdm2 production delay	1/h
$\gamma_m$	0	Mdm2 degradation rate	1/h
$\delta_m$	0.128	Mdm2-mediated Mdm2 degradation rate	Mdm2/h
$K_m$	$1.25 \times 10^{-4}$	Saturation of Mdm2-mediated Mdm2 degradation	Mdm2
$\alpha_{f0}$	0.069	Feedback species mRNA transcription rate	<i>FB</i> /h
$\beta_{f0,NPF}$	0.396	NPF p53-mediated feedback species production	<i>FB</i> /h
$\beta_{f0,2NF}$	31.05	2NF p53-mediated feedback species production	<i>FB</i> /h
$K_{f0}$	0.117	Saturation of p53-mediated feedback production	p53
$\gamma_{f0}$	1	Feedback protein production delay	1/h
$\gamma_f$	1	Feedback protein degradation rate	1/h

The table indicates each parameter's name, its nominal value, a description of its effect, and the units in which it is measured. Dynamics of the core 1NF loop in the absence of feedback (obtained by setting  $\beta_{f0}$  to zero) are identical to those obtained from the original 1NF model. Italicized units are mRNA; otherwise units represent protein levels. For simplicity, units of the feedback species (either MTF1 or MTF1-KRAB) are shown as "FB."

## Other Supporting Information Files

[SI Appendix \(PDF\)](#)