	ı ne propabili ıday is:	ty of a leap yea	r selected at	random contain 53	
		(b) 1/7	(c) 2/7	(d) 53/365	
2.	A bag contains	s 3 red and 2 b	lue marbles. A	A marble is drawn at	
		bability of draw			
	(a) 3/5	(b) 2/5		(d) 1/5	
3 . ⁻	The probabilit	y that it will rai	n tomorrow is	(d) 1/5 o.85. What is the	
	bability that it	will not rain to	morrow		
	(a) 0.25	(b) 0.145		(d) none of these	
4.				cted from the numbers	
(1,	2, 3,,15) is a multiple o	of 4?		
		(b) 4/5	(c) 2/15	(d) 1/3	
5 .	Wnat are tne			ow three coins?	
	(a) 4	(b) 5	· · · ·	(d) 7	
6.	The probability	ty that a prime	numper selec	ted at random from the	
nur	nbers (1,2,3, .	35) is :			
	(a) 12/35	(1	(c) 13/3	(d) none of these	
7.	The sum of th	e probability of	an event and	l non event is :	
	(a) 2	(c)	0 (d) no	one of these.	
8.	The following	probabilities a	re given; cho	ose the correct answer	
for	that which is i	not possible.			
				(d) none of these.	
			ultaneously, t	han the probability of	
get	ting at least tv	vo heads, is: 🧫			
	(a) 1/4	(b) 3/8		(d) 1/8	
10.	A letter is ch	losen at randol	m from the let	tters of the word	
*	SSASSINATIO	$DN oldsymbol{\diamond}$. The prob	ability that th	e letter chosen has:	
	(a) 6/13	(b) 7/13		(d) none of these.	
11 .	A dice is thro			etting an even number.	
(A)	2/3	(B) 1	(C) 5/6	(D) 1/2	
12. Two coins are thrown at the same time. Find the probability of					
getting both heads.					
_	3/4 (P)	(C) 1/2	(D) 0		
13	Two dice are	thrown simults	neously The	probability of getting a	

sum of 9 is:

(A) 1/10	(B) 3/10	(0) 1/ 2	(D) 4/9				
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4		(2)	(D) 2	9/100			
15. A bag contains 5 red balls and some blue balls . If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:							
(A) 5		(C) 15 ((D) 20				
taken out at non-defectiv	16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is:						
(A) 143/150		(C) 1/	/25 ((D) 1/50			
17. Cards marked with numbers 2 to 101 are placed in a box and mixed thoroughly. One card is drawn from this box randomly, then the probability that the number on card is a perfect square. (B) 1/10 (C) 3/10 (D) 19/100							
18. What is (A) 1/7		y of getting 53 M (C) 2/7					
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit.							
		C) 7/52 (D) 1					
equally likel 1,2,312 ;	y to come to re then the proba	nsists of spinning est pointing to o ability that it will (C) 7/12	ne of the nu point to an	mber			
	,	, ,	. ,				
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (B) $1/2$ (C) 1 (D) $1/4$							

22. Riya and Kajal are friends same birthday is the same bir	•	both will have the				
(A) 364/365 (B) 31/365		(D) 1/133225				
23. A number x is chosen at a 2. Then the probability that x (A) 1/5 (B) 2/5	² < 2 is?	numbers -2, -1, 0 , 1,				
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (D) 7						
25. A number is selected at ra Then the probability that it is (A) 7/50 (B) 4/25 (C) 1/2	a multiple of 3 an					
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$ b) $\frac{5}{42}$	c) $\frac{1}{21}$	21				
27. Runs scored by batsman in 5 one day matches are 50, 70, 82, 93, and 20. The standard deviation is						
b) 25.49	c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $\frac{1}{2}$ b) $\frac{1}{3}$		d) $^{1}/_{6}$				
30. X is a variate between 0 a a) 8 b) 7	nd 3. The value o	f E(X²) is				
21 The random variables V or	nd V hove veriens	00 0 2 and 0 E				

31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?

32.Out of the following values, which one is not possible in probability? a) $P(x) = 1$ b) $\sum x P(x) = 3$						
c) $P(x) = 0.5$		- O E				
	= 2 and E(z) = 4,	then E(z – x) =?			
3) 3	b) 6	c) 0	d) Insu	fficient data		
34.The cov	ariance of two ii	ndependent	random variable	is		
a) 1		c) - 1	d) Und	efined		
35.If Σ P(x)) = k ² – 8 then, t	he value of k	c is?			
a) 0	b) 1	` -		ufficient data		
	0.5 and x = 4, th b) 0.5	h en E(x) = ? c) 4				
37.In a disciss always?	crete probability	distribution	, the sum of all p	orobabilities		
a) 0	b) Infinite	~\1	d) Unde	efined		
38.If the pr	robability of hitt	ing the targe	et is 0.4, find me	an and		
	b) 0.6, 0.	.24	c) 0.4, 0.16	d) 0.6, 0.16		
•	robability that a % and if 10 bom	nbs are drop	•			
40. Find the mean of tossing 8 coins. a) 2 b) 40 c) 8 c) 8 d) 1 41. What is the mean and variance for standard normal distribution?						

c) 5

a) 3

b) 4

) Mean is 1 and variance is 0 c) Mean is 0 and variance is ∞ d) Mean is ∞ and variance is 0									
	42. Variance of a random variable X is given by a) $E(X)$ b) $E(X2)$ c $E(X)$ d) $E(X)$ 2								
43.	43.Mean of a random variable X is given by b) E(X2) c) E(X2) - (E(X))2 d) (E(X))2								
44.N a) 0	44.Mean of a constant 'a' is a) 0								
45.\	45.Variance of a constant 'a' is b) a c) a/2 d) 1								
46.Find the mean and variance of X?									
x 0 1 2 3 4									
	Х	U			J	·			
ļ	f(x)	1/9	2/9	3/9	2/9	1/9			

47. Find the expectation of a random variable X?

	X	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		,		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**
- a) \sqrt{np}
- b) \sqrt{pq} c) (np)2