# Lab 1 – MATH 240 – Computational Statistics

First Author Affiliation Department email@domain

#### Abstract

This document provides a basic template for the 2-page labs we will complete each week. Here, you should provide a succinct summary about what you did and why it might be helpful.

**Keywords:** What topics does the lab cover with respect to class?

## 1 Instructions

For this lab, you will

- 1. Install R and RStudio
- Install tinytex (if necessary): install.packages("tinytex")
- Create a GitHub account here, and email me you username.
- 4. Install GitHub desktop.
- 5. Accept the LAB 1 assignment here.
- 6. Recreate this document (except put your name/info at the top) to get used to writing in LATEX and to see the types of things we can do when creating a document to convey statistical information. Make sure to commit and push your work using GitHub desktop as you finish each section.

**Remark:** You will find the class Sweave cheatsheet to be *incredibly* \emph{incredibly} helpful.

# 2 Word Processing Tasks

#### 2.1 Centering Text

We can center text in Sweave

#### 2.2 Bold, Italics, and Underlining

We can **bold**, *italicize*, <u>underline</u>, and *emhpasize* text in Sweave

Note, I did a column break here so that the list wasn't broken across columns.

#### 2.3 Lists, and Numbered Lists

We can write an unordered list in Sweave.

- first item
- second item
- third item

We can write a numbered list in Sweave.

- 1. first item
- 2. second item
- 3. third item

We can write a lettered list in Sweave.

- a. first item
- b. second item
- c. third item

#### 2.4 Submissions

This part of the midterm is due Sunday November 14 by 5p. I will not accept late submissions. Note that you may use this template to help build your introduction and methods sections, and you can use the work you did as a group during the datathon. Still, I expect this submission to be your own summary and extension of that work without collaboration.

#### 2.5 Typing Mathematical Equations

We can write a one line equation that is centered like this

$$\hat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{1i} x_{2i} + \epsilon_i.$$

This can be written in the text, as  $\widehat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{1i} x_{2i} + \epsilon_i$  using as well.

When we need to show multiple steps, we can create a multi-line equation that is centered like this:

$$8(x-5) + x = 9(x-5) + 5$$

$$8x - 40 + x = 9x - 45 + 5$$
 (Distributing)
$$9x - 40 = 9x - 40$$
 (Combining like terms)
$$9x = 9x$$
 (Adding 40 to both sides)
$$x = x$$
 (Dividing both sides by 9)

The equality holds for any x.

Note, I did a page break here so that the next section started on a clean page.

### 2.6 Running R Code

Code chunks can be entered into Sweave; e.g., here are some comments.

```
# R code goes here
# Output is automatically printed in the pdf
```

Below, you can see that we can algebra with R.

```
8*(9-5) + 9 # 8(x-5) + x for x=9
## [1] 41
```

Below, we show we can produce the code without evaluating it.

```
8*(9-5) + 9 # 8(x-5) + x for x=9
```

Alternatively, we can produce the output without the code.

```
## [1] 41
```

We can also call object values from R directly.

```
result <- 8*(9-5) + 9 # 8(x-5) + x for x = 9 result.with.error <- result + rnorm(1, mean = 0, sd = 0.1) result.with.error ## [1] 40.91967
```

The result is 40.9196733. Note that I did not type the result, but I used the **\Sexpr{}** command.

### 2.7 Plotting

We can also plot with R.

```
#Plot a histogram of random normally distributed data
hist(rexp(100))
```

## Histogram of rexp(100)




Figure 1: A histogram of random normally distributed data, n=100.

#### 2.8 Tables

Below, we load and take a peek at some data about the death rate per 1000 in Virginia in 1940 (Molyneaux et al., 1947).

```
data(VADeaths)
head(VADeaths) # Take a peek of the data
         Rural Male Rural Female Urban Male Urban Female
##
## 50-54
               11.7
                              8.7
                                        15.4
                                                       8.4
## 55-59
                             11.7
                                        24.3
                                                      13.6
               18.1
## 60-64
               26.9
                             20.3
                                        37.0
                                                      19.3
## 65-69
                             30.9
                                        54.6
               41.0
                                                      35.1
## 70-74
                             54.3
                                        71.1
                                                      50.0
               66.0
```

If we want to print this nicely, we can do so using the xtable package (Dahl et al., 2019), which we can reference using the label (Table 1)

Rural Male	e Rural Female	Urban Male	Urban Female
11.70	8.70	15.40	8.40
18.10	11.70	24.30	13.60
26.90	20.30	37.00	19.30
41.00	30.90	54.60	35.10
66.00	54.30	71.10	50.00

Table 1: Death Rates per 1000 in Virginia (1940).

### References

Dahl, D. B., Scott, D., Roosen, C., Magnusson, A., and Swinton, J. (2019). xtable: Export Tables to LaTeX or HTML. R package version 1.8-4. Horst, A. M., Hill, A. P., and Gorman, K. B. (2020). palmerpenguins: Palmer Archipelago (Antarctica) penguin data. R package version 0.1.0. Molyneaux, L., Gilliam, S. K., and Florant, L. (1947). Differences in Virginia death rates by color, sex, age and rural or urban residence. American Sociological Review, 12(5):525–535.

## 3 Appendix

Below is a table from a paper I'm currently working on. Without the analysis object in R, I have to create this table myself.

Term	SS (Type III)	df	F	p-value	$\epsilon_p^2$
(Intercept)	4.95	1.00	5.37	0.0209	
White-Poor $(Z)$	3.17	1.00	3.44	0.0642	0.02
Zero-Sum (Z)	17.96	1.00	19.48	< 0.0001	0.03
Education (Z)	0.39	1.00	0.42	0.5161	0.00
Income (Z)	0.16	1.00	0.17	0.6817	0.00
Democrat	9.60	1.00	10.42	0.0013	0.02
Black-Poor (Z)	1.92	1.00	2.08	0.1496	0.00
White-Poor $(Z) \times Zero-Sum(Z)$	7.96	1.00	8.63	0.0034	0.01
Residuals	506.92	550.00			

Table 2: ANOVA table for Case Study I.

The palmerpenguins package for R (Horst et al., 2020) provides data on adult foraging penguins near Palmer Station, Antarctica. Figure 2 is too big to fit nicely in our column-based-template above, so I've placed it here in the abstract by saving it and presenting it scaled to 0.75.

```
library(palmerpenguins)
pdf("figure/penguins.pdf", width = 8, height = 5)
plot(penguins)
dev.off()
```




Figure 2: Data on adult foraging penguins near Palmer Station, Antarctica.

**Bibliography:** Note that when you add citations to your bib.bib file and you cite them in your document, the bibliography section will automatically populate here.