Lab 1 – MATH 240 – Computational Statistics

Caroline Devine
Affiliation
Department
cdevine@colgate.edu

Abstract

This document provides a basic template for the 2-page labs we will complete each week. Here, you should provide a succinct summary about what you did and why it might be helpful.

Keywords: What topics does the lab cover with respect to class?

1 Instructions

For this, you will

- 1. Install R and RStudio
- 2. Install tinytex (if necessary):
 install.packages("tinytex")
- Create a GitHub account here, and email me your username.
- 4. Install GitHub desktop.
- 5. Accept the LAB 1 assignment here
- 6. Recreate this document (except put your name/info at the top) to get used to writing in LATEX and to see that types of things we can do when creating a document to convey statistical information. Make sure to commit and push your working using GitHub desktop as you finish each section.

Remark: You will find the class Sweave cheatsheet to be *incredibly* (\emph{incredibly}) helpful.

2 Word Processing Tasks

2.1 Centering Text

We can center text in Sweave.

2.2 Bold, Italics, and Underlining

We can **bold**, *italicize*, <u>underline</u>, and *emphasize* text in Sweave.

Note, I did a column break here so that the list wasn't broken across columns.

2.3 Lists, and Numbered Lists

We can write an unordered list in Sweave.

- first item
- second item
- third item

We can write a numbered list in Sweave.

- 1. first item
- 2. second item
- 3. third item

We can write a lettered list in Sweave.

- a. first item
- b. second item
- c. third item

2.4 Submissions

This part of the midterm is due Sunday November 14 by 5p. I will not accept late submissions. Note that you may use this template to help build your introduction and methods sections, and you can use the work you did as a group during the datathon. Still, I except this submission to be your own summary and extension of that work without collaboration.

2.5 Typing Mathematical Equations

We can write a one line equation that is centered like this

$$\widehat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{1i} x_{2i} + \epsilon_i.$$

This can be written in the text, as $\widehat{y}_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \beta_3 x_{1i} x_{2i} + \epsilon_i$ using as well.

When we need to show multiple steps, we can create a multi-line equation that is centered like this:

$$8(x-5) + x = 9(x-5) + 5$$

$$8x - 40 + x = 9x - 45 + 5$$
 (Distributing)
$$9x - 40 = 9x - 40$$
 (Combing like terms)
$$9x = 9x$$
 (Adding 40 to both sides)
$$x = x$$
 (Dividing both sides by 9)

The equality holds for any x.

Note, I did a page break here so that the next section started on a clean page.

Running R Code 2.6

Code chunks can be entered into Sweave; e.g., here are some comments.

```
# R code goes here
# Output is automatically printed in the pdf.
```

Below, you can see that we can do algebra with R.

```
8*(9-5) + 9 #8*(x-5) + x for x=9
## [1] 41
```

Below, we show we can produce the code without evaluating it.

```
8*(9-5) + 9 #8*(x-5) + x for x=9
```

Alternatively, we can produce the output without the code.

```
## [1] 41
```

We can also call object values from R directly.

```
result <- 8*(9-5) + 9 \#8*(x-5) + x for x=9
result.with.error \leftarrow result + rnorm(1, mean = 0, sd = 0.1)
result.with.error
## [1] 41.07462
```

The result is 41.0746179. Not that I did not type the result, but I used the \Sexpr{} command.

2.6.1 Plotting

We can also plot with R.

```
#Plot a histogram of random exponential data
hist(rexp(100))
```

Histogram of rexp(100)




Figure 1: A histogram of random exponentially distributed data, n = 100.

Tables 2.7

Below, we load and take a peek at some data about the death rates per 1000 in Virgina in 1940 (Molyneaux et al., 1947).

```
data(VADeaths)
head(VADeaths) # Take a peek of the data
         Rural Male Rural Female Urban Male Urban Female
## 50-54
               11.7
                             8.7
                                        15.4
## 55-59
               18.1
                             11.7
                                        24.3
                                                      13.6
## 60-64
               26.9
                             20.3
                                        37.0
                                                      19.3
## 65-69
               41.0
                             30.9
                                        54.6
                                                     35.1
                                        71.1
## 70-74
               66.0
                            54.3
                                                     50.0
```

If we want to print this nicely, we can do so using the xtable package (Dahl et al., 2019), which we can reference using the label (Table 1).

```
library(xtable)
sleep.table<-xtable(VADeaths ,</pre>
               label = "VADeaths.tab",
              caption = "Death Rates per 100 in Virgina (1940).")
```

Rural Male	Rural Female	Urban Male	Urban Female
11.70	8.70	15.40	8.40
18.10	11.70	24.30	13.60
26.90	20.30	37.00	19.30
41.00	30.90	54.60	35.10
66.00	54.30	71.10	50.00

Table 1: Death Rates per 100 in Virgina (1940).

References

Dahl, D. B., Scott, D., Roosen, C., Magnusson, A., and Swinton, J. (2019). xtable: Export Tables to LaTeX or HTML. R package version 1.8-4.
Horst, A. M., Hill, A. P., and Gorman, K. B. (2020). palmerpenguins: Palmer Archipelago (Antarctica) penguin data. R package version 0.1.0.
Molyneaux, L., Gilliam, S. K., and Florant, L. (1947). Differences in virginia death rates by color, sex, age and rural or urban residence. American Sociological Review, 12(5):525-535.

3 Appendix

Below is a table from a paper I'm currently working on. Without the analysis object in R, I have to create this table myself.

Term	SS(Type III)	df	F	p-value	ϵ_p^2
(Intercept)	4.95	1.00	5.37	0.0209	
White-Poor(Z)	3.17	1.00	3.44	0.0642	0.02
$\operatorname{Zero-Sum}(Z)$	17.96	1.00	19.48	< 0.0001	0.03
Education (Z)	0.39	1.00	0.42	0.5161	0.00
Income(Z)	0.16	1.00	0.17	0.6817	0.00
Democrat	9.60	1.00	10.42	0.0013	0.02
Black-Poor(Z)	1.92	1.00	2.08	0.1496	0.00
White-Poor(Z)xZero-Sum(Z)	7.96	1.00	8.63	0.0034	0.01
Residuals	506.92	550.00			

Table 2: ANOVA table for Case Study I.

The palmerpenguins package for R (Horst et al., 2020) provides data on adult foraging penguins near Palmer Station, Antartica. Figure 2 is too big to fit nicely in our column-based-template above, so I've placed it here in the abstract by saving it and presenting it scaled to 0.75.

```
library(palmerpenguins)
pdf("figure/penguins.pdf", width = 8, height = 5)
plot(penguins)
dev.off()
```




Figure 2: Data on adult foraging penguins near Palmer Station, Antarctica