

Lab Eleven – Point Estimation and Resampling

Tips

- Complete the tasks below. Make sure to start your solutions in on a new line that starts with “**Solution:**”.
- Make sure to use the Quarto Cheatsheet. This will make completing and writing up the lab *much* easier.
- Make sure to use “enter” to make your code readable, so it doesn’t run off the page. I switched the Quarto document to automatically render to .pdf, so you can see what you’re submitting by default. You can switch back for the highlighting by moving the html block above the pdf block in the YAML header.
- Don’t forget that you can copy-and-paste code when you’re doing something similar as we did in class!
- Make sure to plot all computed probabilities. This is good **ggplot** practice AND will help make sure you don’t make a mistake when computing the probability.

1 Introduction

When performing point estimation, there isn’t just one *right* answer. For example, the Method of Moments (MOM) and Maximum Likelihood Estimates (MLEs) aren’t always the same.

One question that comes up, is why we calculate the sample variance as

$$S_{n-1}^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2,$$

when it is supposed to be the “average squared distance from the mean, i.e.,

$$S_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2.$$

2 Comparing Estimators

2.1 Initial Simulation

In this lab, we will explore why we use S_{n-1}^2 and not S_n^2 . To do this, we will generate data from four known distributions:

- Poisson($\lambda = 2$)
- Binomial($n = 50, p = \frac{5-\sqrt{21}}{10}$)
- Normal($\mu = 0, \sigma = \sqrt{2}$)
- Exponential($\lambda = \frac{1}{\sqrt{2}}$).

Remark: Note that I have selected the parameters so that the true variance for all four models is 2.

For each distribution, write a loop that completes the following 1000 times. Put `set.seed(7272)` at the top of your code chunk.

- Generate a sample of size n from the distribution.
- Calculate and save S_{n-1}^2 and S_n^2 .

Start with $n = 20$. Is there evidence that S_{n-1}^2 is a better estimator than S_n^2 ?

Solution

```

1  set.seed(7272)
2  n = 20
3  n.sims=1000
4  # Poisson
5  lambda = 2
6  results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
7  for(i in 1:n.sims){
8    x = rpois(n, lambda)
9    (results_p$var[i] = var(x) ) # variance
10   (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
11  }
12  means_p = tibble(var_mean = mean(results_p$var), not_var_mean = mean(results_p$not_var))
13
14  # Binomial
15  size = 50
16  p = (5-sqrt(21))/10
17  results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
18  for(i in 1:n.sims){
19    x = rbinom(n, size, p)
20    (results_b$var[i] = var(x) ) # variance
21    (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
22  }
23  means_b = tibble(var_mean = mean(results_b$var), not_var_mean = mean(results_b$not_var))
24
25  # Normal
26  mean = 0
27  sd = sqrt(2)
28  results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
29  for(i in 1:n.sims){
30    x = rnorm(n, mean, sd)
31    (results_n$var[i] = var(x) ) # variance
32    (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
33  }
34  means_n = tibble(var_mean = mean(results_n$var), not_var_mean = mean(results_n$not_var))
35
36  # Exponential
37  lambda = 1/sqrt(2)
38  results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
39  for(i in 1:n.sims){
40    x = rexp(n, lambda)
41    (results_e$var[i] = var(x) ) # variance
42    (results_e$not_var[i] = (n-1)/n * var(x)) # the "not" variance - avg. squared dist from mean
43  }
44  means_e = tibble(var_mean = mean(results_e$var), not_var_mean = mean(results_e$not_var))

```

Since we know the true variance of all the models is 2, we can compare the accuracy of the calculated variance and the calculated “not-variance” by seeing how close they are to 2. For every model, at n=20, the variance is closer to 2 than the “not-variance”, which shows that it is a better estimator.

Create a plot or combination of plots (using patchwork) to show the results across all four distributions. Make sure to create a corresponding table that numerically summarizes the results of the simulation. You should include

$$\text{Bias} = \text{Average of Estimates} - \text{True Value}$$

$$\text{Precision} = \frac{1}{\text{Variance of Estimates}}$$

$$\text{Mean Square Error} = \text{Bias}^2 + \text{Variance of Estimates.}$$

Solution

```

1 # biases
2 bias_b = tibble(varbias = mean(results_b$var) - 2, notvarbias=mean(results_b$not_var) - 2)
3 bias_p = tibble(varbias = mean(results_p$var) - 2, notvarbias=mean(results_p$not_var) - 2)
4 bias_n = tibble(varbias = mean(results_n$var) - 2, notvarbias=mean(results_n$not_var) - 2)
5 bias_e = tibble(varbias = mean(results_e$var) - 2, notvarbias=mean(results_e$not_var) - 2)
6
7 # precisions
8 precision_b = tibble(var_prec = 1/var(results_b$var), notvar_prec = 1/var(results_b$not_var))
9 precision_p = tibble(var_prec = 1/var(results_p$var), notvar_prec = 1/var(results_p$not_var))
10 precision_n = tibble(var_prec = 1/var(results_n$var), notvar_prec = 1/var(results_n$not_var))
11 precision_e = tibble(var_prec = 1/var(results_e$var), notvar_prec = 1/var(results_e$not_var))
12
13 # MSE
14 mse_b = tibble(var_mse = (bias_b$varbias)^2 + var(results_b$var),
15 var_notmse = bias_b$notvarbias)^2 + var(results_b$not_var)
16
17 mse_p = tibble(var_mse = (bias_p$varbias)^2 + var(results_p$var),
18 var_notmse = bias_p$notvarbias)^2 + var(results_p$not_var)
19
20 mse_n = tibble(var_mse = (bias_n$varbias)^2 + var(results_n$var),
21 var_notmse = (bias_n$notvarbias)^2 + var(results_n$not_var))
22
23 mse_e = tibble(var_mse = (bias_e$varbias)^2 + var(results_e$var),
24 var_notmse = (bias_e$notvarbias)^2 + var(results_e$not_var))
25
26
27 # combining all calculated values into a tibble for each distrib
28 combined_p = bind_cols(mse_p, bias_p, precision_p, means_p)
29 combined_b = bind_cols(mse_b, bias_b, precision_b, means_b)
30 combined_e = bind_cols(mse_e, bias_e, precision_e, means_e)
31 combined_n = bind_cols(mse_n, bias_n, precision_n, means_n)
32
33 all_dists = bind_rows(combined_p, combined_b, combined_n, combined_e) |>
34   add_column(Distribution = c("Poisson", "Binomial", "Normal", "Exponential"), .before = 1)
35
36 all_dists |>
37   knitr::kable(digits = 2)

```

Table 1: Different properties of the distributions

Distribution	var_mse	var_notmse	varbias	notvarbias	var_prec	notvar_prec	var_mean	not_var_mean
Poisson	0.66	0.44	-0.02	-0.12	2.10	2.33	1.98	1.88
Binomial	0.64	0.44	-0.04	-0.13	2.15	2.38	1.96	1.87
Normal	0.49	0.45	0.00	-0.10	2.03	2.25	2.00	1.90
Exponential	1.43	1.30	-0.02	-0.12	0.70	0.77	1.98	1.88


```

1 poisson_var = ggplot(results_p) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 poisson_notvar = ggplot(results_p) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")

```


```
1 poisson_var + poisson_notvar
```

Figure 1: Boxplots comparing the variances of the two estimators for variance, with data from the Poisson distribution



```
1 binom_var = ggplot(results_b) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 binom_notvar = ggplot(results_b) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
12
13 binom_var + binom_notvar
```

Figure 2: Boxplots comparing the variances of the two estimators for variance, with data from the Binomial distribution




```
1 norm_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
```

```

6
7 norm_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1
norm_var + norm_notvar

```

Figure 3: Boxplots comparing the variances of the two estimators for variance, with data from the Normal distribution




```

1 exp_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 exp_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1
exp_var + exp_notvar

```

Figure 4: Boxplots comparing the variances of the two estimators for variance, with data from the Exponential distribution



2.2 Simulation over various sample sizes

Consider the effect of the sample size have on these results. Provide a graphical and numerical summary of what happens as n increases – say $n = 20$, $n = 50$, $n = 100$, and $n = 500$. Does the result in your initial simulation hold across samples? Or, do things change as the sample size changes? Ensure to set your randomization seed.

Solution

Here, I reuse my original code, and change n to 50.

```
1 set.seed(7272)
2 n = 50
3 n.sims=1000
4 # Poisson
5 lambda = 2
6 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
7 for(i in 1:n.sims){
8   x = rpois(n, lambda)
9   (results_p$var[i] = var(x) ) # variance
10  (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
11 }
12 means_p = tibble(var_mean = mean(results_p$var), not_var_mean = mean(results_p$not_var))
13
14 # Binomial
15 size = 50
16 p = (5-sqrt(21))/10
17 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
18 for(i in 1:n.sims){
19   x = rbinom(n, size, p)
20   (results_b$var[i] = var(x) ) # variance
21   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
22 }
23 means_b = tibble(var_mean = mean(results_b$var), not_var_mean = mean(results_b$not_var))
24
25 # Normal
26 mean = 0
27 sd = sqrt(2)
28 results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
29 for(i in 1:n.sims){
30   x = rnorm(n, mean, sd)
31   (results_n$var[i] = var(x) ) # variance
32   (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
33 }
34 means_n = tibble(var_mean = mean(results_n$var), not_var_mean = mean(results_n$not_var))
35
36 # Exponential
37 lambda = 1/sqrt(2)
38 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
39 for(i in 1:n.sims){
40   x = rexp(n, lambda)
41   (results_e$var[i] = var(x) ) # variance
42   (results_e$not_var[i] = (n-1)/n * var(x)) # the "not" variance - avg. squared dist from mean
43 }
44 means_e = tibble(var_mean = mean(results_e$var), not_var_mean = mean(results_e$not_var))
45 # biases
46 bias_b = tibble(varbias = mean(results_b$var) - 2, notvarbias=mean(results_b$not_var) - 2)
47 bias_p = tibble(varbias = mean(results_p$var) - 2, notvarbias=mean(results_p$not_var) - 2)
48 bias_n = tibble(varbias = mean(results_n$var) - 2, notvarbias=mean(results_n$not_var) - 2)
49 bias_e = tibble(varbias = mean(results_e$var) - 2, notvarbias=mean(results_e$not_var) - 2)
```

```

50 # precisions
51 precision_b = tibble(var_prec = 1/var(results_b$var), notvar_prec = 1/var(results_b$not_var))
52 precision_p = tibble(var_prec = 1/var(results_p$var), notvar_prec = 1/var(results_p$not_var))
53 precision_n = tibble(var_prec = 1/var(results_n$var), notvar_prec = 1/var(results_n$not_var))
54 precision_e = tibble(var_prec = 1/var(results_e$var), notvar_prec = 1/var(results_e$not_var))
55
56 # MSE
57 mse_b = tibble(var_mse = (bias_b$varbias)^2 + var(results_b$var),
58 var_notmse = bias_b$notvarbias^2 + var(results_b$not_var))
59
60 mse_p = tibble(var_mse = (bias_p$varbias)^2 + var(results_p$var),
61 var_notmse = bias_p$notvarbias^2 + var(results_p$not_var))
62
63 mse_n = tibble(var_mse = (bias_n$varbias)^2 + var(results_n$var),
64 var_notmse = (bias_n$notvarbias)^2 + var(results_n$not_var))
65
66 mse_e = tibble(var_mse = (bias_e$varbias)^2 + var(results_e$var),
67 var_notmse = (bias_e$notvarbias)^2 + var(results_e$not_var))
68
69
70 # combining all calculated values into a tibble for each distrib
71 combined_p = bind_cols(mse_p, bias_p, precision_p, means_p)
72 combined_b = bind_cols(mse_b, bias_b, precision_b, means_b)
73 combined_e = bind_cols(mse_e, bias_e, precision_e, means_e)
74 combined_n = bind_cols(mse_n, bias_n, precision_n, means_n)
75
76
77 all_dists = bind_rows(combined_p, combined_b, combined_n, combined_e) |>
78   add_column(Distribution = c("Poisson", "Binomial", "Normal", "Exponential"), .before = 1)
79
80 all_dists |>
81   knitr::kable(digits = 2)

```

Table 2: Different properties of the distributions, when n = 100


Distribution	var_mse	var_notmse	varbias	notvarbias	var_prec	notvar_prec	var_mean	not_var_mean
Poisson	0.22	0.19	-0.03	-0.07	5.17	5.39	1.97	1.93
Binomial	0.22	0.18	-0.01	-0.05	5.31	5.53	1.99	1.95
Normal	0.17	0.16	-0.01	-0.05	5.98	6.23	1.99	1.95
Exponential	0.70	0.67	-0.02	-0.06	1.43	1.49	1.98	1.94

```

1 poisson_var = ggplot(results_p) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 poisson_notvar = ggplot(results_p) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
12
13 poisson_var + poisson_notvar


```

Figure 5: Boxplots comparing the variances of the two estimators for variance, with data from the Poisson distribution, when $n = 50$



```
1 binom_var = ggplot(results_b) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 binom_notvar = ggplot(results_b) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1
1 binom_var + binom_notvar
```

Figure 6: Boxplots comparing the variances of the two estimators for variance, with data from the Binomial distribution, when $n = 50$




```
1 norm_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
```

```

7 norm_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 norm_var + norm_notvar

```

Figure 7: Boxplots comparing the variances of the two estimators for variance, with data from the Normal distribution, where $n = 50$




```

1 exp_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 exp_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 exp_var + exp_notvar

```

Figure 8: Boxplots comparing the variances of the two estimators for variance, with data from the Exponential distribution where $n = 50$



Here, I do the same thing, but change n to 100.

```
1 set.seed(7272)
2 n = 100
3 n.sims=1000
4 # Poisson
5 lambda = 2
6 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
7 for(i in 1:n.sims){
8   x = rpois(n, lambda)
9   (results_p$var[i] = var(x) ) # variance
10  (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
11 }
12 means_p = tibble(var_mean = mean(results_p$var), not_var_mean = mean(results_p$not_var))
13
14 # Binomial
15 size = 50
16 p = (5-sqrt(21))/10
17 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
18 for(i in 1:n.sims){
19   x = rbinom(n, size, p)
20   (results_b$var[i] = var(x) ) # variance
21   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
22 }
23 means_b = tibble(var_mean = mean(results_b$var), not_var_mean = mean(results_b$not_var))
24
25 # Normal
26 mean = 0
27 sd = sqrt(2)
28 results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
29 for(i in 1:n.sims){
30   x = rnorm(n, mean, sd)
31   (results_n$var[i] = var(x) ) # variance
32   (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
33 }
34 means_n = tibble(var_mean = mean(results_n$var), not_var_mean = mean(results_n$not_var))
35
36 # Exponential
37 lambda = 1/sqrt(2)
38 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
39 for(i in 1:n.sims){
40   x = rexp(n, lambda)
41   (results_e$var[i] = var(x) ) # variance
42   (results_e$not_var[i] = (n-1)/n * var(x)) # the "not" variance - avg. squared dist from mean
43 }
44 means_e = tibble(var_mean = mean(results_e$var), not_var_mean = mean(results_e$not_var))
45
46 # biases
47 bias_b = tibble(varbias = mean(results_b$var) - 2, notvarbias=mean(results_b$not_var) - 2)
48 bias_p = tibble(varbias = mean(results_p$var) - 2, notvarbias=mean(results_p$not_var) - 2)
49 bias_n = tibble(varbias = mean(results_n$var) - 2, notvarbias=mean(results_n$not_var) - 2)
50 bias_e = tibble(varbias = mean(results_e$var) - 2, notvarbias=mean(results_e$not_var) - 2)
51
52 # precisions
53 precision_b = tibble(var_prec = 1/var(results_b$var), notvar_prec = 1/var(results_b$not_var))
54 precision_p = tibble(var_prec = 1/var(results_p$var), notvar_prec = 1/var(results_p$not_var))
55 precision_n = tibble(var_prec = 1/var(results_n$var), notvar_prec = 1/var(results_n$not_var))
56 precision_e = tibble(var_prec = 1/var(results_e$var), notvar_prec = 1/var(results_e$not_var))
```

```

56 # MSE
57 mse_b = tibble(var_mse = (bias_b$varbias)^2 + var(results_b$var),
58 var_notmse = bias_b$notvarbias)^2 + var(results_b$not_var)
59
60
61 mse_p = tibble(var_mse = (bias_p$varbias)^2 + var(results_p$var),
62 var_notmse = bias_p$notvarbias)^2 + var(results_p$not_var)
63
64 mse_n = tibble(var_mse = (bias_n$varbias)^2 + var(results_n$var),
65 var_notmse = (bias_n$notvarbias)^2 + var(results_n$not_var))
66
67 mse_e = tibble(var_mse = (bias_e$varbias)^2 + var(results_e$var),
68 var_notmse = (bias_e$notvarbias)^2 + var(results_e$not_var))
69
70
71 # combining all calculated values into a tibble for each distrib
72 combined_p = bind_cols(mse_p, bias_p, precision_p, means_p)
73 combined_b = bind_cols(mse_b, bias_b, precision_b, means_b)
74 combined_e = bind_cols(mse_e, bias_e, precision_e, means_e)
75 combined_n = bind_cols(mse_n, bias_n, precision_n, means_n)
76
77 all_dists = bind_rows(combined_p, combined_b, combined_n, combined_e) |>
78   add_column(Distribution = c("Poisson", "Binomial", "Normal", "Exponential"), .before = 1)
79
80 all_dists |>
81   knitr::kable(digits = 2)

```

Table 3: Different properties of the distributions, when n = 100


Distribution	var_mse	var_notmse	varbias	notvarbias	var_prec	notvar_prec	var_mean	not_var_mean
Poisson	0.11	0.10	-0.02	-0.04	10.07	10.27	1.98	1.96
Binomial	0.12	0.11	0.00	-0.02	9.30	9.49	2.00	1.98
Normal	0.09	0.09	0.01	-0.01	11.37	11.60	2.01	1.99
Exponential	0.30	0.29	0.02	0.00	3.33	3.40	2.02	2.00

```

1 poisson_var = ggplot(results_p) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 poisson_notvar = ggplot(results_p) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
12
13 poisson_var + poisson_notvar

```

Figure 9: Boxplots comparing the variances of the two estimators for variance, with data from the Poisson distribution, when $n = 100$




```

1 binom_var = ggplot(results_b) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 binom_notvar = ggplot(results_b) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1
1 binom_var + binom_notvar

```

Figure 10: Boxplots comparing the variances of the two estimators for variance, with data from the Binomial distribution, when $n = 100$



```

1 norm_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6


```

```

7 norm_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 norm_var + norm_notvar

```

Figure 11: Boxplots comparing the variances of the two estimators for variance, with data from the Normal distribution, where $n = 100$




```

1 exp_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 exp_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 exp_var + exp_notvar

```

Figure 12: Boxplots comparing the variances of the two estimators for variance, with data from the Exponential distribution where $n = 100$



Finally, I do the same thing again but with $n = 500$.

```
1 set.seed(7272)
2 n = 500
3 n.sims=1000
4 # Poisson
5 lambda = 2
6 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
7 for(i in 1:n.sims){
8   x = rpois(n, lambda)
9   (results_p$var[i] = var(x) ) # variance
10  (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
11 }
12 means_p = tibble(var_mean = mean(results_p$var), not_var_mean = mean(results_p$not_var))
13
14 # Binomial
15 size = 50
16 p = (5-sqrt(21))/10
17 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
18 for(i in 1:n.sims){
19   x = rbinom(n, size, p)
20   (results_b$var[i] = var(x) ) # variance
21   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
22 }
23 means_b = tibble(var_mean = mean(results_b$var), not_var_mean = mean(results_b$not_var))
24
25 # Normal
26 mean = 0
27 sd = sqrt(2)
28 results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
29 for(i in 1:n.sims){
30   x = rnorm(n, mean, sd)
31   (results_n$var[i] = var(x) ) # variance
32   (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
33 }
34 means_n = tibble(var_mean = mean(results_n$var), not_var_mean = mean(results_n$not_var))
35
36 # Exponential
37 lambda = 1/sqrt(2)
38 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
39 for(i in 1:n.sims){
40   x = rexp(n, lambda)
41   (results_e$var[i] = var(x) ) # variance
42   (results_e$not_var[i] = (n-1)/n * var(x)) # the "not" variance - avg. squared dist from mean
43 }
44 means_e = tibble(var_mean = mean(results_e$var), not_var_mean = mean(results_e$not_var))
45
46 # biases
47 bias_b = tibble(varbias = mean(results_b$var) - 2, notvarbias=mean(results_b$not_var) - 2)
48 bias_p = tibble(varbias = mean(results_p$var) - 2, notvarbias=mean(results_p$not_var) - 2)
49 bias_n = tibble(varbias = mean(results_n$var) - 2, notvarbias=mean(results_n$not_var) - 2)
50 bias_e = tibble(varbias = mean(results_e$var) - 2, notvarbias=mean(results_e$not_var) - 2)
51
52 # precisions
53 precision_b = tibble(var_prec = 1/var(results_b$var), notvar_prec = 1/var(results_b$not_var))
54 precision_p = tibble(var_prec = 1/var(results_p$var), notvar_prec = 1/var(results_p$not_var))
55 precision_n = tibble(var_prec = 1/var(results_n$var), notvar_prec = 1/var(results_n$not_var))
56 precision_e = tibble(var_prec = 1/var(results_e$var), notvar_prec = 1/var(results_e$not_var))
```

```

56 # MSE
57 mse_b = tibble(var_mse = (bias_b$varbias)^2 + var(results_b$var),
58 var_notmse = bias_b$notvarbias)^2 + var(results_b$not_var)
59
60
61 mse_p = tibble(var_mse = (bias_p$varbias)^2 + var(results_p$var),
62 var_notmse = bias_p$notvarbias)^2 + var(results_p$not_var)
63
64 mse_n = tibble(var_mse = (bias_n$varbias)^2 + var(results_n$var),
65 var_notmse = (bias_n$notvarbias)^2 + var(results_n$not_var))
66
67 mse_e = tibble(var_mse = (bias_e$varbias)^2 + var(results_e$var),
68 var_notmse = (bias_e$notvarbias)^2 + var(results_e$not_var))
69
70
71 # combining all calculated values into a tibble for each distrib
72 combined_p = bind_cols(mse_p, bias_p, precision_p, means_p)
73 combined_b = bind_cols(mse_b, bias_b, precision_b, means_b)
74 combined_e = bind_cols(mse_e, bias_e, precision_e, means_e)
75 combined_n = bind_cols(mse_n, bias_n, precision_n, means_n)
76
77 all_dists = bind_rows(combined_p, combined_b, combined_n, combined_e) |>
78   add_column(Distribution = c("Poisson", "Binomial", "Normal", "Exponential"), .before = 1)
79
80 all_dists |>
81   knitr::kable(digits = 2)

```

Table 4: Different properties of the distributions


Distribution	var_mse	var_notmse	varbias	notvarbias	var_prec	notvar_prec	var_mean	not_var_mean
Poisson	0.02	0.02	0	-0.01	50.32	50.52	2	1.99
Binomial	0.02	0.02	0	-0.01	55.41	55.63	2	1.99
Normal	0.02	0.02	0	0.00	58.58	58.81	2	2.00
Exponential	0.06	0.06	0	0.00	15.71	15.77	2	2.00

```

1 poisson_var = ggplot(results_p) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 poisson_notvar = ggplot(results_p) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
12
13 poisson_var + poisson_notvar


```

Figure 13: Boxplots comparing the variances of the two estimators for variance, with data from the Poisson distribution, when $n = 500$



```
1 binom_var = ggplot(results_b) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 binom_notvar = ggplot(results_b) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1
1 binom_var + binom_notvar
```

Figure 14: Boxplots comparing the variances of the two estimators for variance, with data from the Binomial distribution, when $n = 500$




```
1 norm_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
```

```

7 norm_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 norm_var + norm_notvar

```

Figure 15: Boxplots comparing the variances of the two estimators for variance, with data from the Normal distribution, where $n = 500$




```

1 exp_var = ggplot(results_n) +
2   geom_boxplot(aes(x="", y = var), width=.1) +
3   geom_hline(yintercept = 2, color = "red") +
4   theme_bw() +
5   labs(x = "", y="Variance", title = "Variance estimator")
6
7 exp_notvar = ggplot(results_n) +
8   geom_boxplot(aes(x="", y = not_var), width=.1) +
9   geom_hline(yintercept = 2, color = "red") +
10  theme_bw() +
11  labs(x = "", y="Variance", title = "'Not-variance' estimator")
1 exp_var + exp_notvar

```

Figure 16: Boxplots comparing the variances of the two estimators for variance, with data from the Exponential distribution where $n = 500$



Looking at the different tables produced for varying values of n , we can see that the bias and mse approaches 0 as n increases. We can also see that the mean approaches 2, and that the precision increases as n increases. The boxplots show the same trend, which can be seen by the median approaching the true mean of 2, suggesting that as the size of the data increases, outliers have a smaller effect on the data. However, the variance is still more accurate than the “not-variance”, even at larger sample sizes, although the difference is not as large as it was at small sizes.

2.3 Simulation

You have written code to draw a random sample of various sizes from each distribution, and to calculate and store the calculated estimates. Rework your code to produce a density plot of the s_{n-1}^2 and s_n^2 . Do you have a guess about their distribution? Ensure to set your randomization seed.


Solution

```

1 n = 20
2 # Poisson
3 lambda = 2
4 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_p$var[i] = var(x) ) # variance
8   (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }
10
11 #poisson
12 set.seed(7272)
13 pvar_dplot=ggplot(data=results_p) +
14   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
15   xlab("(S_(n-1))^2")
16 pnotvar_dplot=ggplot(data=results_p) +
17   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
18   xlab("(S_(n))^2")
19
20 pvar_dplot + pnotvar_dplot

```

Figure 17: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Poisson distribution when $n = 20$



```

1 n = 50
2 # Poisson
3 lambda = 2
4 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_p$var[i] = var(x) ) # variance


```

```

8 (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }
1 #poisson
2 set.seed(7272)
3 pvar_dplot=ggplot(data=results_p) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 pnotvar_dplot=ggplot(data=results_p) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 pvar_dplot + pnotvar_dplot

```

Figure 18: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Poisson distribution when $n = 50$




```

1 n = 100
2 # Poisson
3 lambda = 2
4 results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_p$var[i] = var(x) ) # variance
8   (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }
1 #poisson
2 set.seed(7272)
3 pvar_dplot=ggplot(data=results_p) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 pnotvar_dplot=ggplot(data=results_p) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 pvar_dplot + pnotvar_dplot

```

Figure 19: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Poisson distribution when $n = 100$




```

1  n = 500
2  # Poisson
3  lambda = 2
4  results_p = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5  for(i in 1:n.sims){
6    x = rpois(n, lambda)
7    (results_p$var[i] = var(x)) # variance
8    (results_p$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9  }

10 #poisson
11 set.seed(7272)
12 pvar_dplot=ggplot(data=results_p) +
13   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
14   xlab("(S_{n-1})^2")
15 pnotvar_dplot=ggplot(data=results_p) +
16   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
17   xlab("(S_n)^2")
18 pvar_dplot + pnotvar_dplot

```

Figure 20: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Poisson distribution when $n = 500$




```

1 n = 20
2 # binomial
3 lambda = 2
4 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_b$var[i] = var(x) ) # variance
8   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 set.seed(7272)
2 bvar_dplot=ggplot(data=results_b) +
3   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4   xlab("(S_(n-1))^2")
5 bnotvar_dplot=ggplot(data=results_b) +
6   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7   xlab("(S_(n))^2")
8
9 bvar_dplot + bnotvar_dplot

```

Figure 21: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Binomial distribution when $n = 20$




```

1 n = 50
2 # binomial
3 lambda = 2
4 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_b$var[i] = var(x) ) # variance
8   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 set.seed(7272)
2 bvar_dplot=ggplot(data=results_b) +
3   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4   xlab("(S_(n-1))^2")
5 bnotvar_dplot=ggplot(data=results_b) +
6   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7   xlab("(S_(n))^2")
8
9 bvar_dplot + bnotvar_dplot

```

Figure 22: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Binomial distribution when $n = 50$




```

1  n = 100
2  # binomial
3  lambda = 2
4  results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5  for(i in 1:n.sims){
6    x = rpois(n, lambda)
7    (results_b$var[i] = var(x) ) # variance
8    (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9  }

10 #poisson
11 set.seed(7272)
12 bvar_dplot=ggplot(data=results_b) +
13   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
14   xlab("(S_{(n-1)})^2")
15 bnotvar_dplot=ggplot(data=results_b) +
16   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
17   xlab("(S_n)^2")
18 bvar_dplot + bnotvar_dplot

```

Figure 23: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Binomial distribution when $n = 100$




```

1 n = 500
2 # binomial
3 lambda = 2
4 results_b = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_b$var[i] = var(x) ) # variance
8   (results_b$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 #poisson
2 set.seed(7272)
3 bvar_dplot=ggplot(data=results_b) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 bnotvar_dplot=ggplot(data=results_b) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 bvar_dplot + bnotvar_dplot

```

Figure 24: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Binomial distribution when $n = 500$




```

1 n = 20
2 # exponential
3 lambda = 2
4 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_e$var[i] = var(x) ) # variance
8   (results_e$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 #poisson
2 set.seed(7272)
3 evar_dplot=ggplot(data=results_e) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 enotvar_dplot=ggplot(data=results_e) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 evar_dplot + enotvar_dplot

```

Figure 25: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Exponential distribution when $n = 20$




```

1  n = 50
2  # exponential
3  lambda = 2
4  results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5  for(i in 1:n.sims){
6    x = rpois(n, lambda)
7    (results_e$var[i] = var(x) ) # variance
8    (results_e$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9  }

10 #poisson
11 set.seed(7272)
12 evar_dplot=ggplot(data=results_e) +
13   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
14   xlab("(S_{n-1})^2")
15 enotvar_dplot=ggplot(data=results_e) +
16   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
17   xlab("(S_n)^2")
18 evar_dplot + enotvar_dplot

```

Figure 26: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Exponential distribution when $n = 50$




```

1 n = 100
2 # exponential
3 lambda = 2
4 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_e$var[i] = var(x) ) # variance
8   (results_e$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 #poisson
2 set.seed(7272)
3 evar_dplot=ggplot(data=results_e) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 enotvar_dplot=ggplot(data=results_e) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 evar_dplot + enotvar_dplot

```

Figure 27: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Exponential distribution when $n = 100$




```

1 n = 500
2 # exponential
3 lambda = 2
4 results_e = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_e$var[i] = var(x) ) # variance
8   (results_e$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 #poisson
2 set.seed(7272)
3 evar_dplot=ggplot(data=results_e) +
4   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
5   xlab("(S_(n-1))^2")
6 enotvar_dplot=ggplot(data=results_e) +
7   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
8   xlab("(S_(n))^2")
9
10 evar_dplot + enotvar_dplot

```

Figure 28: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Exponential distribution when $n = 500$




```

1  n = 20
2  # normal
3  lambda = 2
4  results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5  for(i in 1:n.sims){
6    x = rpois(n, lambda)
7    (results_n$var[i] = var(x) ) # variance
8    (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9  }

1  set.seed(7272)
2  nvar_dplot=ggplot(data=results_n) +
3    stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4    xlab("(S_{(n-1)})^2")
5  nnotvar_dplot=ggplot(data=results_n) +
6    stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7    xlab("(S_n)^2")
8
9  nvar_dplot + nnotvar_dplot

```

Figure 29: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Normal distribution when $n = 20$




```

1 n = 50
2 # normal
3 lambda = 2
4 results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_n$var[i] = var(x) ) # variance
8   (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 set.seed(7272)
2 nvar_dplot=ggplot(data=results_n) +
3   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4   xlab("(S_(n-1))^2")
5 nnotvar_dplot=ggplot(data=results_n) +
6   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7   xlab("(S_(n))^2")
8
9 nvar_dplot + nnotvar_dplot

```

Figure 30: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Normal distribution when $n = 50$




```

1 n = 100
2 # normal
3 lambda = 2
4 results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5 for(i in 1:n.sims){
6   x = rpois(n, lambda)
7   (results_n$var[i] = var(x) ) # variance
8   (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9 }

1 set.seed(7272)
2 nvar_dplot=ggplot(data=results_n) +
3   stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4   xlab("(S_(n-1))^2")
5 nnotvar_dplot=ggplot(data=results_n) +
6   stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7   xlab("(S_(n))^2")
8
9 nvar_dplot + nnotvar_dplot

```

Figure 31: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Normal distribution when $n = 100$




```

1  n = 500
2  # normal
3  lambda = 2
4  results_n = tibble(var=numeric(n.sims), not_var=numeric(n.sims))
5  for(i in 1:n.sims){
6    x = rpois(n, lambda)
7    (results_n$var[i] = var(x) ) # variance
8    (results_n$not_var[i] = (n-1)/n * var(x)) # the not variance - avg. squared dist from mean
9  }

1  set.seed(7272)
2  nvar_dplot=ggplot(data=results_n) +
3    stat_density(aes(x=var), color="chartreuse2", geom="line", linewidth=1) +
4    xlab("(S_{(n-1)})^2")
5  nnotvar_dplot=ggplot(data=results_n) +
6    stat_density(aes(x=not_var), color="chartreuse2", geom="line", linewidth=1) +
7    xlab("(S_n)^2")
8
9  nvar_dplot + nnotvar_dplot

```

Figure 32: Density plots of the $(s_n)^2$ and the $(s_{(n-1)})^2$ of the Normal distribution when $n = 500$



2.4 Bootstrap


Now, take *one* sample from each distribution. Instead of drawing a random sample at each iteration, sample *with replacement* from the one sample. Produce a density plot of the s_{n-1}^2 and s_n^2 . Do we get roughly the same information as we do in the full simulation? Try this over several seeds – what do you notice? Ensure to set your randomization seed.

Solution

```
1 set.seed(7272)
2 n = 20
3 b = 1000
4 original_sample = rpois(n, lambda = lambda)
5 sample_var = var(original_sample)
6 notvar = var(original_sample)*(n-1)/n
7 boot_var = numeric(b)
8 boot_notvar = numeric(b)
9 for (i in 1:b) {
10   resample = sample(original_sample, size = n, replace = TRUE)
11   boot_var[i] = var(resample)
12   boot_notvar[i] = var(resample)*(n-1)/n
13 }
14 bootstrap_b = tibble(bootstrap_var = boot_var, bootstrap_notvar = boot_notvar)

1 ggplot(data = bootstrap_b) +
2   stat_density(aes(x = boot_var, color = "(s_(n-1))^2"), geom = "line", linewidth = 1) +
3   stat_density(aes(x = bootstrap_notvar, color = "(s_n)^2"), geom = "line", linewidth = 1) +
4   scale_color_manual(values = c("(s_(n-1))^2" = "chartreuse2",
5   "(s_n)^2" = "darkorchid2")) +
6   labs(x = "Value")
```

Figure 33: Density plot of the bootstrap of the variance and the variance * (n-1)/n, where n = 20



```
1 set.seed(7272)
2 n = 50
3 b = 1000
4 original_sample = rpois(n, lambda = lambda)
5 sample_var = var(original_sample)
6 notvar = var(original_sample)*(n-1)/n
7 boot_var = numeric(b)
8 boot_notvar = numeric(b)
9 for (i in 1:b) {
```


```

10   resample = sample(original_sample, size = n, replace = TRUE)
11   boot_var[i] = var(resample)
12   boot_notvar[i] = var(resample)*(n-1)/n
13 }
14 bootstrap_b = tibble(bootstrap_var = boot_var, bootstrap_notvar = boot_notvar)

1 ggplot(data = bootstrap_b) +
2   stat_density(aes(x = boot_var, color = "(s_(n-1))^2"), geom = "line", linewidth = 1) +
3   stat_density(aes(x = bootstrap_notvar, color = "(s_n)^2"), geom = "line", linewidth = 1) +
4   scale_color_manual(values = c("(s_(n-1))^2" = "chartreuse2",
5                                 "(s_n)^2" = "darkorchid2")) +
6   labs(x = "Value")

```

Figure 34: Density plot of the bootstrap of the variance and the variance $\times (n-1)/n$, where $n = 50$




```

1 set.seed(7272)
2 n = 100
3 b = 1000
4 original_sample = rpois(n, lambda = lambda)
5 sample_var = var(original_sample)
6 notvar = var(original_sample)*(n-1)/n
7 boot_var = numeric(b)
8 boot_notvar = numeric(b)
9 for (i in 1:b) {
10   resample = sample(original_sample, size = n, replace = TRUE)
11   boot_var[i] = var(resample)
12   boot_notvar[i] = var(resample)*(n-1)/n
13 }
14 bootstrap_b = tibble(bootstrap_var = boot_var, bootstrap_notvar = boot_notvar)

1 ggplot(data = bootstrap_b) +
2   stat_density(aes(x = boot_var, color = "(s_(n-1))^2"), geom = "line", linewidth = 1) +
3   stat_density(aes(x = bootstrap_notvar, color = "(s_n)^2"), geom = "line", linewidth = 1) +
4   scale_color_manual(values = c("(s_(n-1))^2" = "chartreuse2",
5                                 "(s_n)^2" = "darkorchid2")) +
6   labs(x = "Value")

```

Figure 35: Density plot of the bootstrap of the variance and the variance $\times (n-1)/n$, where $n = 100$




```

1  set.seed(7272)
2  n = 500
3  b = 1000
4  original_sample = rpois(n, lambda = lambda)
5  sample_var = var(original_sample)
6  notvar = var(original_sample)*(n-1)/n
7  boot_var = numeric(b)
8  boot_notvar = numeric(b)
9  for (i in 1:b) {
10     resample = sample(original_sample, size = n, replace = TRUE)
11     boot_var[i] = var(resample)
12     boot_notvar[i] = var(resample)*(n-1)/n
13 }
14 bootstrap_b = tibble(bootstrap_var = boot_var, bootstrap_notvar = boot_notvar)

1  ggplot(data = bootstrap_b) +
2     stat_density(aes(x = boot_var, color = "(s_(n-1))^2"), geom = "line", linewidth = 1) +
3     stat_density(aes(x = bootstrap_notvar, color = "(s_n)^2"), geom = "line", linewidth = 1) +
4     scale_color_manual(values = c("(s_(n-1))^2" = "chartreuse2",
5                                   "(s_n)^2" = "darkorchid2")) +
6     labs(x = "Value")

```

Figure 36: Density plot of the bootstrap of the variance and the variance $\cdot (n-1)/n$, where $n = 100$



3 Executive Summary

Summarize the work you've done here to provide a brief (200-300 word) summary of the simulation and the results using your work in Lab 11. Your summary should be professional, clear, and all claims should be corroborated with statistical support.