

In lecture 16, we looked at precipitation amounts in Madison County (at Morrisville station). We found that the Weibull distribution had a good fit to the monthly precipitation amounts.

We found that the MLEs for the Weibull distribution were

$$\hat{a} = 2.1871$$

$$\hat{\sigma} = 3.9683$$

and

$$-\mathcal{L}(\{\hat{a}, \hat{\sigma}\} | \mathbf{x}) = 2166.496$$

is the realized negative log-likelihood. Note this means that the log-likelihood is

$$\mathcal{L}(\{\hat{a}, \hat{\sigma}\} | \mathbf{x}) = -2166.496,$$

and the usual likelihood is

$$L(\{\hat{a}, \hat{\sigma}\} | \mathbf{x}) = e^{[\mathcal{L}(\{\hat{a}, \hat{\sigma}\} | \mathbf{x})]} \approx e^{-2166.496},$$

which R cannot differentiate from 0.

1. Someone asked “why Weibull?” in class. That is, why wouldn’t we use another right-skewed distribution like the Gamma (see Lecture 15), or the Log-Normal (see Lecture 17).

- (a) Compute the MLEs for these data using a Gamma distribution.

The MLEs for the Gamma distribution are $\hat{\alpha} = 4.1742$, $\hat{\beta} = 0.8411$, and a negative log-likelihood ($-\mathcal{L}(\hat{\alpha}, \hat{\beta} | \mathbf{x})$) of 2151.149.

- (b) Compute the MLEs for these data using the Log-Normal distribution.

The MLEs for the Log-Normal distribution are $\hat{\mu} = 1.1314$, $\hat{\sigma} = 0.5333$, and a negative log-likelihood ($-\mathcal{L}(\hat{\mu}, \hat{\sigma} | \mathbf{x})$) of 2204.201.

- (c) Compute the likelihood ratio to compare the Weibull and the Gamma distribution. Which has a better fit according to the likelihood ratio?

$$Q = \frac{L(\{\hat{a}, \hat{\sigma}\} | \mathbf{x})}{L(\{\hat{\alpha}, \hat{\beta}\} | \mathbf{x})} = e^{[\mathcal{L}(\{\hat{a}, \hat{\sigma}\} | \mathbf{x}) - \mathcal{L}(\{\hat{\alpha}, \hat{\beta}\} | \mathbf{x})]}$$

```
log.lik.weib <- -2166.496
log.lik.gamma <- -2151.149
(q <- exp(log.lik.weib - log.lik.gamma))

## [1] 2.162134e-07
```

Since likelihood ratio, $Q < 1$, the Gamma distribution has a better fit for our data.

- (d) Compute the likelihood ratio to compare the Weibull and the Log-Normal distribution. Which has a better fit according to the likelihood ratio?

$$Q = \frac{L(\{\hat{a}, \hat{\sigma}\} | \mathbf{x})}{L(\{\hat{\mu}, \hat{\sigma}\} | \mathbf{x})} = e^{[\mathcal{L}(\{\hat{a}, \hat{\sigma}\} | \mathbf{x}) - \mathcal{L}(\{\hat{\mu}, \hat{\sigma}\} | \mathbf{x})]}$$

```
log.lik.weib <- -2166.496
log.lik.lognorm <- -2204.201
(q <- exp(log.lik.weib - log.lik.lognorm))

## [1] 2.371775e+16
```

Since likelihood ratio $Q > 1$, the Weibull distribution has a better fit for our data.

- (e) Compute the likelihood ratio to compare the Gamma and the Log-Normal distribution. Which has a better fit according to the likelihood ratio?

$$Q = \frac{L(\{\hat{\alpha}, \hat{\beta}\} | \mathbf{x})}{L(\{\hat{\mu}, \hat{\sigma}\} | \mathbf{x})} = e^{[\mathcal{L}(\{\hat{\alpha}, \hat{\beta}\} | \mathbf{x}) - \mathcal{L}(\{\hat{\mu}, \hat{\sigma}\} | \mathbf{x})]}$$

```
(q <- exp(log.lik.gamma - log.lik.lognorm))  
## [1] 1.09696e+23
```

Since likelihood ratio $Q > 1$, the Gamma distribution has a better fit for our data.

2. Optional Coding Challenge. Choose the “best” distribution and refit the model by season.

- (a) Fit the Distribution for Winter (December-February).
- (b) Fit the Distribution for Spring (March-May).
- (c) Fit the Distribution for Summer (June-August).
- (d) Fit the Distribution for Fall (September-November).
- (e) Plot the four distributions in one plot using **cyan3** for Winter, **chartreuse3** for Spring, **red3** for Summer, and **chocolate3** for Fall. Note any similarities/differences you observe across the seasons.