```
"id": "obj.1277",
"name": "Human[tax:9606] CAG repeats[S0:0000726] detection for Huntington's
disease[DOID:12858]",
"version": "1.0",
"createdby": "hadley king@gwmail.gwu.edu",
"created": "Jan 26, 2017 17:44:38",
"modified": "Mar 03, 2017 10:16:14",
"digital_signature": "QdB7vu4rLeLtMFmPHQ9S",
"verification_status": "unreviewed",
"publication_status": "draft",
"usability_domain": [
    "Determine CAG repeats[S0:0000726] in a sequence for the diagnosis of
    Huntingtons disease[ICD:G10] for htt gene[G0ID:G0272344]: 10 - 35 CAG
    trinucleotide repeats: No Risk, 36 - 40 CAG trinucleotide repeats:
   Moderate Risk, 40 CAG trinucleotide repeats: High Risk"
],
"authors": [
    {
        "orchid": "0000-0002-9920-565X"
    }
],
"description_domain": {
    "keywords": [
        "disease",
        "human disease"
    ],
    "xref": [
        "S0:0000726",
        "tax:9606",
        "DOID: 12858",
        "GOID:G0272344",
        "ICD:G10"
    ],
    "pipeline_steps": [
        {
            "tool_name": "HIVE_hexagon",
            "tool_desc": "Alignment of reads to a set of references",
            "tool_version": "1.3",
            "tool_package": "null",
            "step_number": "1",
            "input_uri_list": [
                "hive://nuc-read/514683",
                "hive://nuc-read/514682",
                "http://www.ncbi.nlm.nih.gov/nuccore/SRR2028290",
                "http://www.ncbi.nlm.nih.gov/nuccore/NM 0002111.8"
            ],
            "output_uri_list": [
                "hive://data/509234/allCount-aligned.csv"
            ]
```

{

```
},
            "tool_name": "HIVE_heptagon",
            "tool_desc": "variant calling",
            "tool_version": "1.3",
            "tool_package": "null",
            "step_number": "2",
            "input_uri_list": [
                "hive://data/509234/allCount-aligned.csv"
            ],
            "output_uri_list": [
                "hive://data/509234/SNPProofile.csv"
            ]
        }
    1
},
"execution_domain": {
    "script": "hive://workflows/huntingtons_disease_detection.hwf",
    "pipeline_version": "1.0",
    "pipeline_url": "https://hive.biochemistry.gwu.edu/workflows/",
    "platform": "HIVE",
    "driver": "hive://hive-driver",
    "prerequisites": [
        "{\"name\":\"HIVE_hexagon\",\"version\":\"}",
        "{\"name\":\"HIVE_heptagon\",\"version\":\"1.3\"}"
    ],
    "env_parameters": [
        "HIVEv1.3"
},
"parametric domain": {
    "divergence_threshold_percent": "10",
    "alignment_seed": "14 letters",
    "alignment min match len": "50",
    "variant_calling_min_coverage": "20",
    "variant_calling_frequency_cutoff": "0.2"
},
"io_domain": {
    "reference_uri_list": [
        "http://www.ncbi.nlm.nih.gov/nuccore/SRR2028290",
        "http://www.ncbi.nlm.nih.gov/nuccore/NM_0002111.8"
    ],
    "input_uri_list": [
        "hive://nuc-read/514683",
        "hive://nuc-read/514682"
    ],
    "output_uri_list": [
        "hive://data/509234/SNPProofile.csv"
    1
},
"error_domain": [
```

```
"false positive discovery <= 0.05",
    "false negative discovery <= 0.10"
]</pre>
```