

Ponatinib / Asciminib nella LMC – Newsletter interna (ultimi 15 giorni)

Generata il 16/06/2026

Generata il: 2026-06-16

Newsletter interna – sintesi operativa

Messaggi chiave (da leggere in 60 secondi):

- Nel periodo emergono soprattutto temi su **Resistenza / T315I / mutazioni** (n=1).
- Secondo filone: **TFR / stop terapia** (n=1).
- Terzo filone: **Safety e comorbidità** (n=1).

Cosa cambia per noi (azioni pratiche):

- Rivedere/rafforzare **monitoraggio cardiovascolare e interazioni** (es. anticoagulanti) nei pazienti in TKI.
- Valutare criteri interni per **consolidamento / de-escalation / TFR** nei pazienti in deep response.
- Verificare percorso di **test mutazionale** e criteri di switch nelle resistenze (incl. T315I).

Safety radar (segnali/interazioni da monitorare):

- Monitorare evoluzione su **TFR / stop terapia** (nuovi segnali/dati).
- Monitorare evoluzione su **Safety e comorbidità** (nuovi segnali/dati).
- Monitorare evoluzione su **Resistenza / T315I / mutazioni** (nuovi segnali/dati).

Top 5 da leggere subito (ordinati per impatto)

1) From Solid Organ Malignancies to Dual Leukaemias: Sequential CLL and CML in a Cancer Survivor.

Altro

European journal of case reports in internal medicine

Score: 4

Focus clinico: TFR / stop terapia, Safety e comorbidità, Sequencing / linee di terapia

PMID: 42266392 | 10.12890/2026_006813

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42266392/>

- **Tipo di studio/popolazione:** He began treatment with Ibrutinib in 2018 due to bulky symptomatic adenopathy and later switched to acalabrutinib due to arthralgias.
- **Risultato chiave:** Following a hospitalization in 2022 for coronavirus disease 2019, he was found to have chronic myeloid leukaemia (CML), confirmed by BCR-ABL positivity and bone marrow biopsy.
- **Rilevanza clinica:** Imatinib was started for CML but due to potential drug-induced pulmonary toxicity, switched to asciminib.

2) Selective HDAC6 inhibition perturbs autophagy and enhances integrated stress response-mediated immunogenic apoptosis in chronic myeloid leukemia.

Preclinico / laboratorio

Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie Score: 2

Focus clinico: Resistenza / T315I / mutazioni

PMID: 42284888 | 10.1016/j.biopha.2026.119652

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42284888/>

- **Tipo di studio/popolazione:** Chronic myeloid leukemia (CML) is driven by the BCR-ABL1 fusion oncoprotein and is managed with tyrosine kinase inhibitors (TKIs).
- **Risultato chiave:** Autophagy-mediated survival signaling contributes to therapeutic resistance in CML.
- **Rilevanza clinica:** These findings identify HDAC6-driven autophagy as a therapeutically exploitable vulnerability in CML that, when combined with asciminib, triggers ISR-dependent immunogenic apoptosis.

Evidenze per domande cliniche

Resistenza / T315I / mutazioni (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Selective HDAC6 inhibition perturbs autophagy and enhances integrated stress response-mediated immunogenic apoptosis in chronic myeloid leukemia.

Preclinico / laboratorio

Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie Score: 2

PMID: 42284888 | 10.1016/j.biopha.2026.119652

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42284888/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Chronic myeloid leukemia (CML) is driven by the BCR-ABL1 fusion oncoprotein and is managed with tyrosine kinase inhibitors (TKIs).
- **Risultato chiave:** Autophagy-mediated survival signaling contributes to therapeutic resistance in CML.
- **Rilevanza clinica:** These findings identify HDAC6-driven autophagy as a therapeutically exploitable vulnerability in CML that, when combined with asciminib, triggers ISR-dependent immunogenic apoptosis.

TFR / stop terapia (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

From Solid Organ Malignancies to Dual Leukaemias: Sequential CLL and CML in a Cancer Survivor.

Altro

European journal of case reports in internal medicine

Score: 4

PMID: 42266392 | 10.12890/2026_006813

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42266392/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** He began treatment with Ibrutinib in 2018 due to bulky symptomatic adenopathy and later switched to acalabrutinib due to arthralgias.
- **Risultato chiave:** Following a hospitalization in 2022 for coronavirus disease 2019, he was found to have chronic myeloid leukaemia (CML), confirmed by BCR-ABL positivity and bone marrow biopsy.
- **Rilevanza clinica:** Imatinib was started for CML but due to potential drug-induced pulmonary toxicity, switched to asciminib.

Safety e comorbidità (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

From Solid Organ Malignancies to Dual Leukaemias: Sequential CLL and CML in a Cancer Survivor.

Altro

European journal of case reports in internal medicine

Score: 4

PMID: 42266392 | 10.12890/2026_006813

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42266392/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** He began treatment with Ibrutinib in 2018 due to bulky symptomatic adenopathy and later switched to acalabrutinib due to arthralgias.
- **Risultato chiave:** Following a hospitalization in 2022 for coronavirus disease 2019, he was found to have chronic myeloid leukaemia (CML), confirmed by BCR-ABL positivity and bone marrow biopsy.
- **Rilevanza clinica:** Imatinib was started for CML but due to potential drug-induced pulmonary toxicity, switched to asciminib.

Sequencing / linee di terapia (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

From Solid Organ Malignancies to Dual Leukaemias: Sequential CLL and CML in a Cancer Survivor.

Altro

*European journal of case reports in internal
medicine*

Score: 4

PMID: 42266392 | 10.12890/2026_006813

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42266392/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** He began treatment with Ibrutinib in 2018 due to bulky symptomatic adenopathy and later switched to acalabrutinib due to arthralgias.
- **Risultato chiave:** Following a hospitalization in 2022 for coronavirus disease 2019, he was found to have chronic myeloid leukaemia (CML), confirmed by BCR-ABL positivity and bone marrow biopsy.
- **Rilevanza clinica:** Imatinib was started for CML but due to potential drug-induced pulmonary toxicity, switched to asciminib.