

Ponatinib / Asciminib nella LMC – Newsletter interna (ultimi 15 giorni)

Generata il 01/09/2026

Generata il: 2026-09-01

Newsletter interna – sintesi operativa

Messaggi chiave (da leggere in 60 secondi):

- Nel periodo emergono soprattutto temi su **Resistenza / T315I / mutazioni** (n=2).
- Secondo filone: **Sequencing / linee di terapia** (n=2).
- Terzo filone: **Safety e comorbidità** (n=1).

Cosa cambia per noi (azioni pratiche):

- Rivedere/rafforzare **monitoraggio cardiovascolare e interazioni** (es. anticoagulanti) nei pazienti in TKI.
- Verificare percorso di **test mutazionale** e criteri di switch nelle resistenze (incl. T315I).

Safety radar (segnali/interazioni da monitorare):

- Monitorare evoluzione su **Safety e comorbidità** (nuovi segnali/dati).
- Monitorare evoluzione su **Resistenza / T315I / mutazioni** (nuovi segnali/dati).

Top 5 da leggere subito (ordinati per impatto)

1) Sequential TKI Treatment in Patients With Chronic Phase Chronic Myeloid Leukemia Resistant to Frontline Nilotinib or Dasatinib Therapy. A Real-Life Survey From the Italian Campus CML Network.

Linee guida / consenso

Clinical lymphoma, myeloma & leukemia

Score: 12

Focus clinico: Resistenza / T315I / mutazioni, Sequencing / linee di terapia, Real-world (utilità pratica)

PMID: 42608256 | 10.1016/j.cml.2026.07.011

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42608256/>

- **Tipo di studio/popolazione:** We conducted a multicenter, real-world, retrospective analysis on 69 chronic phase (CP) CML patients who switched to second-line treatment in case of failure (according to the European LeukemiaNet 2020 recommendations) after first-line nilotinib or dasatinib.
- **Risultato chiave:** The median time to switch to second-line therapy was 15 months, and ponatinib was the most frequently used TKI (68%).
- **Rilevanza clinica:** As best molecular response (MR), 78% of patients achieved a BCR::ABL1^{IS} transcript level <1%.

2) PGAM5-mediated VDAC1 oligomerization facilitates ponatinib-induced cardiotoxicity via disrupting the mitophagy-mitochondrial unfolded protein response synergistic defense crosstalk.

Preclinico / laboratorio

Chemico-biological interactions

Score: 8

Focus clinico: Resistenza / T315I / mutazioni, Safety e comorbidità, Sequencing / linee di terapia

PMID: 42628894 | 10.1016/j.cbi.2026.112310

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42628894/>

- **Tipo di studio/popolazione:** Ponatinib is an effective tyrosine kinase inhibitor for chronic myeloid leukemia with the T315I mutation, but its clinical use is often limited by serious cardiovascular toxicity.
- **Risultato chiave:** We therefore investigated whether the PGAM5/VDAC1 axis mediates ponatinib-induced cardiac injury by coordinately disrupting mitophagy and the mitochondrial unfolded protein response (UPR^{mt}).
- **Rilevanza clinica:** Single-cell RNA sequencing, in vivo genetic loss-of-function models, and HL-1 cells with stable Pgam5 knockdown were used to define the underlying mechanism.

Evidenze per domande cliniche

Resistenza / T315I / mutazioni (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Sequential TKI Treatment in Patients With Chronic Phase Chronic Myeloid Leukemia Resistant to Frontline Nilotinib or Dasatinib Therapy. A Real-Life Survey From the Italian Campus CML Network.

Linee guida / consenso

Clinical lymphoma, myeloma & leukemia

Score: 12

PMID: 42608256 | 10.1016/j.cml.2026.07.011

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42608256/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** We conducted a multicenter, real-world, retrospective analysis on 69 chronic phase (CP) CML patients who switched to second-line treatment in case of failure (according to the European LeukemiaNet 2020 recommendations) after first-line nilotinib or dasatinib.
- **Risultato chiave:** The median time to switch to second-line therapy was 15 months, and ponatinib was the most frequently used TKI (68%).
- **Rilevanza clinica:** As best molecular response (MR), 78% of patients achieved a BCR::ABL1^{IS} transcript level <1%.

Mini-riassunto (da abstract, estrattivo):

PGAM5-mediated VDAC1 oligomerization facilitates ponatinib-induced cardiotoxicity via disrupting the mitophagy-mitochondrial unfolded protein response synergistic defense crosstalk.

Preclinico / laboratorio

Chemico-biological interactions

Score: 8

PMID: 42628894 | 10.1016/j.cbi.2026.112310

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42628894/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Ponatinib is an effective tyrosine kinase inhibitor for chronic myeloid leukemia with the T315I mutation, but its clinical use is often limited by serious cardiovascular toxicity.
- **Risultato chiave:** We therefore investigated whether the PGAM5/VDAC1 axis mediates ponatinib-induced cardiac injury by coordinately disrupting mitophagy and the mitochondrial unfolded protein response (UPR^{mt}).
- **Rilevanza clinica:** Single-cell RNA sequencing, in vivo genetic loss-of-function models, and HL-1 cells with stable Pgam5 knockdown were used to define the underlying mechanism.

Sequencing / linee di terapia (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Sequential TKI Treatment in Patients With Chronic Phase Chronic Myeloid Leukemia Resistant to Frontline Nilotinib or Dasatinib Therapy. A Real-Life Survey From the Italian Campus CML Network.

Linee guida / consenso

Clinical lymphoma, myeloma & leukemia

Score: 12

PMID: 42608256 | 10.1016/j.cml.2026.07.011

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42608256/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** We conducted a multicenter, real-world, retrospective analysis on 69 chronic phase (CP) CML patients who switched to second-line treatment in case of failure (according to the European LeukemiaNet 2020 recommendations) after first-line nilotinib or dasatinib.
- **Risultato chiave:** The median time to switch to second-line therapy was 15 months, and ponatinib was the most frequently used TKI (68%).
- **Rilevanza clinica:** As best molecular response (MR), 78% of patients achieved a BCR::ABL1^{IS} transcript level <1%.

Mini-riassunto (da abstract, estrattivo):

PGAM5-mediated VDAC1 oligomerization facilitates ponatinib-induced cardiotoxicity via disrupting the mitophagy-mitochondrial unfolded protein response synergistic defense crosstalk.

Preclinico / laboratorio

Chemico-biological interactions

Score: 8

PMID: 42628894 | 10.1016/j.cbi.2026.112310

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42628894/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Ponatinib is an effective tyrosine kinase inhibitor for chronic myeloid leukemia with the T315I mutation, but its clinical use is often limited by serious cardiovascular toxicity.
- **Risultato chiave:** We therefore investigated whether the PGAM5/VDAC1 axis mediates ponatinib-induced cardiac injury by coordinately disrupting mitophagy and the mitochondrial unfolded protein response (UPR^{mt}).
- **Rilevanza clinica:** Single-cell RNA sequencing, in vivo genetic loss-of-function models, and HL-1 cells with stable Pgam5 knockdown were used to define the underlying mechanism.

Safety e comorbidità (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

PGAM5-mediated VDAC1 oligomerization facilitates ponatinib-induced cardiotoxicity via disrupting the mitophagy-mitochondrial unfolded protein response synergistic defense crosstalk.

Preclinico / laboratorio

Chemico-biological interactions

Score: 8

PMID: 42628894 | 10.1016/j.cbi.2026.112310

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42628894/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Ponatinib is an effective tyrosine kinase inhibitor for chronic myeloid leukemia with the T315I mutation, but its clinical use is often limited by serious cardiovascular toxicity.
- **Risultato chiave:** We therefore investigated whether the PGAM5/VDAC1 axis mediates ponatinib-induced cardiac injury by coordinately disrupting mitophagy and the mitochondrial unfolded protein response (UPR^{mt}).
- **Rilevanza clinica:** Single-cell RNA sequencing, in vivo genetic loss-of-function models, and HL-1 cells with stable Pgam5 knockdown were used to define the underlying mechanism.

Real-world (utilità pratica) (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Sequential TKI Treatment in Patients With Chronic Phase Chronic Myeloid Leukemia Resistant to Frontline Nilotinib or Dasatinib Therapy. A Real-Life Survey From the Italian Campus CML Network.

Linee guida / consenso

Clinical lymphoma, myeloma & leukemia

Score: 12

PMID: 42608256 | 10.1016/j.cml.2026.07.011

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42608256/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** We conducted a multicenter, real-world, retrospective analysis on 69 chronic phase (CP) CML patients who switched to second-line treatment in case of failure (according to the European LeukemiaNet 2020 recommendations) after first-line nilotinib or dasatinib.
- **Risultato chiave:** The median time to switch to second-line therapy was 15 months, and ponatinib was the most frequently used TKI (68%).
- **Rilevanza clinica:** As best molecular response (MR), 78% of patients achieved a BCR::ABL1^{IS} transcript level <1%.