

Ponatinib / Asciminib nella LMC – Newsletter interna (ultimi 15 giorni)

Generata il 01/06/2026

Generata il: 2026-06-01

Newsletter interna – sintesi operativa

Messaggi chiave (da leggere in 60 secondi):

- Nel periodo emergono soprattutto temi su **Dose strategy / consolidamento** (n=3).
- Secondo filone: **Sequencing / linee di terapia** (n=3).
- Terzo filone: **Resistenza / T315I / mutazioni** (n=2).

Cosa cambia per noi (azioni pratiche):

- Rivedere/rafforzare **monitoraggio cardiovascolare e interazioni** (es. anticoagulanti) nei pazienti in TKI.
- Valutare criteri interni per **consolidamento / de-escalation / TFR** nei pazienti in deep response.
- Verificare percorso di **test mutazionale** e criteri di switch nelle resistenze (incl. T315I).

Safety radar (segnali/interazioni da monitorare):

- Monitorare evoluzione su **TFR / stop terapia** (nuovi segnali/dati).
 - Monitorare evoluzione su **Safety e comorbidità** (nuovi segnali/dati).
 - Monitorare evoluzione su **Resistenza / T315I / mutazioni** (nuovi segnali/dati).
-

Top 5 da leggere subito (ordinati per impatto)

1) Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

Focus clinico: Resistenza / T315I / mutazioni, Dose strategy / consolidamento, Safety e comorbidità, Sequencing / linee di terapia, Real-world (utilità pratica)

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

2) Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

Focus clinico: Resistenza / T315I / mutazioni, TFR / stop terapia, Dose strategy / consolidamento, Sequencing / linee di terapia, Sintesi (review/linee guida)

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

3) Mechanisms, Clinical Phenotype and Potential Risk Prediction for Cardiovascular Toxicity Induced by Tyrosine Kinase Inhibitors In Chronic Myeloid Leukemia.

Review

Dose-response : a publication of International
Hormesis Society

Score: 10

Focus clinico: Safety e comorbidità, Sintesi (review/linee guida)

PMID: 42182616 | 10.1177/15593258261454350

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42182616/>

- **Tipo di studio/popolazione:** Tyrosine kinase inhibitors (TKIs) form the cornerstone of targeted therapy for chronic myeloid leukemia (CML), significantly improving patient prognosis.
- **Risultato chiave:** However, long-term use of TKIs, particularly second-generation (e.g., nilotinib) and third-generation (e.g., ponatinib) drugs, has led to adverse cardiovascular events (CV-AEs).
- **Rilevanza clinica:** CV-AEs pose a critical challenge to patients' long-term quality of life and treatment adherence.

4) Assessment of Transitioning from High-Potency to Low-Potency Inhibitors in Chronic Myeloid Leukemia (CML) Patients: The Downgrading-Impact (D-IMPACT) Project.

Altro

Cancers

Score: 8

Focus clinico: TFR / stop terapia, Dose strategy / consolidamento, Sequencing / linee di terapia

PMID: 42193014 | 10.3390/cancers18101656

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42193014/>

- **Tipo di studio/popolazione:** **Background:** Since the advent of imatinib, more potent next-generation tyrosine kinase inhibitors (TKIs) and asciminib have expanded therapeutic options for chronic myeloid leukemia (CML).
- **Risultato chiave:** Treatment-free remission (TFR) is an important goal in CML management, but only ~30% of patients can achieve it, leaving many on lifelong therapy.
- **Rilevanza clinica:** **Methods:** We analyzed this strategy in 157 patients across 29 Italian CML Campus centres.

5) Neutrophilic Panniculitis in a Pediatric Patient Receiving Ponatinib.

Altro

Pediatric dermatology

Score: 3

Focus clinico: Altro

PMID: 42152725 | 10.1111/pde.70275

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42152725/>

- **Tipo di studio/popolazione:** Ponatinib is a tyrosine kinase inhibitor used in children with chronic myeloid leukemia.
- **Risultato chiave:** We report a case of neutrophilic panniculitis following ponatinib administration in a child.
- **Rilevanza clinica:** With the expanded use of immunomodulatory medications in the pediatric population, this report aims to expand knowledge regarding potential drug-related adverse events and the inclusion of ponatinib-related adverse events in the differential diagnosis of neutrophilic panniculitis.

Evidenze per domande cliniche

Dose strategy / consolidamento (3)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

Mini-riassunto (da abstract, estrattivo):

Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

Mini-riassunto (da abstract, estrattivo):

Assessment of Transitioning from High-Potency to Low-Potency Inhibitors in Chronic Myeloid Leukemia (CML) Patients: The Downgrading-Impact (D-IMPACT) Project.

Altro

Cancers

Score: 8

PMID: 42193014 | 10.3390/cancers18101656

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42193014/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Background: Since the advent of imatinib, more potent next-generation tyrosine kinase inhibitors (TKIs) and asciminib have expanded therapeutic options for chronic myeloid leukemia (CML).
- **Risultato chiave:** Treatment-free remission (TFR) is an important goal in CML management, but only ~30% of patients can achieve it, leaving many on lifelong therapy.
- **Rilevanza clinica:** Methods: We analyzed this strategy in 157 patients across 29 Italian CML Campus centres.

Sequencing / linee di terapia (3)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

Mini-riassunto (da abstract, estrattivo):

Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

Mini-riassunto (da abstract, estrattivo):

Assessment of Transitioning from High-Potency to Low-Potency Inhibitors in Chronic Myeloid Leukemia (CML) Patients: The Downgrading-Impact (D-IMPACT) Project.

Altro

Cancers

Score: 8

PMID: 42193014 | 10.3390/cancers18101656

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42193014/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Background: Since the advent of imatinib, more potent next-generation tyrosine kinase inhibitors (TKIs) and asciminib have expanded therapeutic options for chronic myeloid leukemia (CML).
- **Risultato chiave:** Treatment-free remission (TFR) is an important goal in CML management, but only ~30% of patients can achieve it, leaving many on lifelong therapy.
- **Rilevanza clinica:** Methods: We analyzed this strategy in 157 patients across 29 Italian CML Campus centres.

Resistenza / T315I / mutazioni (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

Mini-riassunto (da abstract, estrattivo):

Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

TFR / stop terapia (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

Mini-riassunto (da abstract, estrattivo):

Assessment of Transitioning from High-Potency to Low-Potency Inhibitors in Chronic Myeloid Leukemia (CML) Patients: The Downgrading-Impact (D-IMPACT) Project.

Altro

Cancers

Score: 8

PMID: 42193014 | 10.3390/cancers18101656

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42193014/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Background: Since the advent of imatinib, more potent next-generation tyrosine kinase inhibitors (TKIs) and asciminib have expanded therapeutic options for chronic myeloid leukemia (CML).
- **Risultato chiave:** Treatment-free remission (TFR) is an important goal in CML management, but only ~30% of patients can achieve it, leaving many on lifelong therapy.
- **Rilevanza clinica:** Methods: We analyzed this strategy in 157 patients across 29 Italian CML Campus centres.

Sintesi (review/linee guida) (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Chronic myeloid leukemia: incorporation of recent advances into current treatment algorithms.

Review

Blood cancer journal

Score: 10

PMID: 42209462 | 10.1038/s41408-026-01527-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42209462/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Several approved and investigational BCR::ABL1 tyrosine kinase inhibitors (TKIs) and STAMP inhibitors are used for the treatment of chronic myeloid leukemia in chronic phase (CML-CP).
- **Risultato chiave:** Achieving a complete cytogenetic response (BCR::ABL1 transcripts on the International Scale [IS] <1%) with TKI therapy within the first year is associated with normalization of survival, whereas achieving a deeper molecular response (BCR::ABL1 transcripts < 0.01% [IS]) may allow for treatment discontinuation with the possibility of treatment-free remission.
- **Rilevanza clinica:** In patients experiencing failure of frontline therapy due to resistance or intolerance, multiple second- and third-line options are available, including second-generation TKIs, ponatinib, and asciminib, as well as novel investigational agents, including the ABL1 kinase domain inhibitors olverembatinib and ELVN-001 and the STAMP inhibitors TGRX-678 and TERN-701.

Mini-riassunto (da abstract, estrattivo):

Riassunto aggiuntivo (full text Open Access, estrattivo):

Mechanisms, Clinical Phenotype and Potential Risk Prediction for Cardiovascular Toxicity Induced by Tyrosine Kinase Inhibitors In Chronic Myeloid Leukemia.

Review

Dose-response : a publication of International
Hormesis Society

Score: 10

PMID: 42182616 | 10.1177/15593258261454350

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42182616/>

Open Access (Europe PMC/PMC): sì (PMC13195214)

- **Tipo di studio/popolazione:** Tyrosine kinase inhibitors (TKIs) form the cornerstone of targeted therapy for chronic myeloid leukemia (CML), significantly improving patient prognosis.
- **Risultato chiave:** However, long-term use of TKIs, particularly second-generation (e.g., nilotinib) and third-generation (e.g., ponatinib) drugs, has led to adverse cardiovascular events (CV-AEs).
- **Rilevanza clinica:** CV-AEs pose a critical challenge to patients' long-term quality of life and treatment adherence.
- **Metodi (full text OA):** non riportato.
- **Risultato principale (full text OA):** non riportato.
- **Dettaglio utile (full text OA):** non riportato.
- **Messaggio clinico (full text OA):** Future research in this field should prioritize addressing the clearly defined knowledge gaps identified in this review, such as elucidating the molecular basis of ponatinib's unique vascular risk, validating predictive biomarkers, and prospectively evaluating existing risk prediction models in real-world chronic myeloid leukemia (CML) cohorts.

Safety e comorbidità (2)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

Mini-riassunto (da abstract, estrattivo):

Riassunto aggiuntivo (full text Open Access, estrattivo):

Mechanisms, Clinical Phenotype and Potential Risk Prediction for Cardiovascular Toxicity Induced by Tyrosine Kinase Inhibitors In Chronic Myeloid Leukemia.

Review

Dose-response : a publication of International Hormesis Society

Score: 10

PMID: 42182616 | 10.1177/15593258261454350

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42182616/>

Open Access (Europe PMC/PMC): sì (PMC13195214)

- **Tipo di studio/popolazione:** Tyrosine kinase inhibitors (TKIs) form the cornerstone of targeted therapy for chronic myeloid leukemia (CML), significantly improving patient prognosis.
- **Risultato chiave:** However, long-term use of TKIs, particularly second-generation (e.g., nilotinib) and third-generation (e.g., ponatinib) drugs, has led to adverse cardiovascular events (CV-AEs).
- **Rilevanza clinica:** CV-AEs pose a critical challenge to patients' long-term quality of life and treatment adherence.
- **Metodi (full text OA):** non riportato.
- **Risultato principale (full text OA):** non riportato.
- **Dettaglio utile (full text OA):** non riportato.
- **Messaggio clinico (full text OA):** Future research in this field should prioritize addressing the clearly defined knowledge gaps identified in this review, such as elucidating the molecular basis of ponatinib's unique vascular risk, validating predictive biomarkers, and prospectively evaluating existing risk prediction models in real-world chronic myeloid leukemia (CML) cohorts.

Real-world (utilità pratica) (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Real-world outcomes of ponatinib in heavily pretreated patients with chronic myeloid leukemia and Philadelphia chromosome-positive acute lymphoblastic leukemia.

Real-world / osservazionale

Annals of hematology

Score: 19

PMID: 42174310 | 10.1007/s00277-026-07053-6

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42174310/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** To evaluate the real-world efficacy and safety of ponatinib, we conducted a multicenter, prospective post-marketing surveillance study in patients with chronic phase (CP), accelerated phase (AP), or blast phase (BP) chronic myeloid leukemia (CML) or Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph + ALL), who were resistant or intolerant to prior tyrosine kinase inhibitors (TKIs) or harbored the T315I mutation.
- **Risultato chiave:** In CP-CML, the overall major molecular response (MMR) rate was 43.2% (35/81), with a cumulative MMR incidence of 23.9% at week 24 among patients without prior MMR.
- **Rilevanza clinica:** CP-CML patients receiving ponatinib as earlier-line therapy achieved higher MR4.5 rates than those treated later (47.4% vs.

Altro (1)

Score: 0

Mini-riassunto (da abstract, estrattivo):

Neutrophilic Panniculitis in a Pediatric Patient Receiving Ponatinib.

Altro

Pediatric dermatology

Score: 3

PMID: 42152725 | 10.1111/pde.70275

Link PubMed: <https://pubmed.ncbi.nlm.nih.gov/42152725/>

Open Access (Europe PMC/PMC): non rilevato

- **Tipo di studio/popolazione:** Ponatinib is a tyrosine kinase inhibitor used in children with chronic myeloid leukemia.
- **Risultato chiave:** We report a case of neutrophilic panniculitis following ponatinib administration in a child.
- **Rilevanza clinica:** With the expanded use of immunomodulatory medications in the pediatric population, this report aims to expand knowledge regarding potential drug-related adverse events and the inclusion of ponatinib-related adverse events in the differential diagnosis of neutrophilic panniculitis.