Ponatinib / Asciminib nella LMC – Newsletter interna (ultimi 15 giorni)

Generata il 16/10/2026

Generata il: 2026-10-16

Newsletter interna – sintesi operativa

Messaggi chiave (da leggere in 60 secondi):

- • Nel periodo emergono soprattutto temi su **Resistenza / T315I / mutazioni** (n=3).
- • Secondo filone: **TFR / stop terapia** (n=2).
- • Terzo filone: **Safety e comorbidità** (n=2).

Cosa cambia per noi (azioni pratiche):

- • Rivedere/rafforzare **monitoraggio cardiovascolare e interazioni** (es. anticoagulanti) nei pazienti in TKI.
- • Valutare criteri interni per **consolidamento / de-escalation / TFR** nei pazienti in deep response.
- • Verificare percorso di **test mutazionale** e criteri di switch nelle resistenze (incl. T315I).

Safety radar (segnali/interazioni da monitorare):

- • Monitorare evoluzione su **TFR / stop terapia** (nuovi segnali/dati).
- • Monitorare evoluzione su **Safety e comorbidità** (nuovi segnali/dati).
- • Monitorare evoluzione su **Resistenza / T315I / mutazioni** (nuovi segnali/dati).

Top 5 da leggere subito (ordinati per impatto)

1) Successful Pregnancy After Planned Pregnancy-Oriented TKI Interruption and Embryo Transfer Following Ponatinib Re-Consolidation in Chronic-Phase Chronic Myeloid Leukemia.

Trial clinico *EJHaem* *Score:* ***18***

***Focus clinico**: TFR / stop terapia, Dose strategy / consolidamento*

PMID: 42840123 | 10.1002/jha2.70426

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42840123/`

- • **Tipo di studio/popolazione:** Pregnancy after unsuccessful treatment-free remission (TFR) in chronic-phase chronic myeloid leukemia (CP-CML) remains challenging.
- • **Risultato chiave:** Ponatinib re-consolidation enabled a second TKI-free interval, with MMR maintained but deep molecular response unstable.
- • **Rilevanza clinica:** Further ponatinib re-consolidation restored MR4.5; ponatinib was stopped before embryo transfer, resulting in pregnancy and live birth without TKI exposure.

2) The evolving landscape of tyrosine kinase inhibitors in chronic-phase chronic myeloid leukemia: A 25-year journey.

Altro *Blood reviews* Score: ***11***

***Focus clinico**: Resistenza / T315I / mutazioni, TFR / stop terapia, Sequencing / linee di terapia*

PMID: 42875330 | 10.1016/j.blre.2026.101436

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42875330/`

- • **Tipo di studio/popolazione:** BCR::ABL1 tyrosine kinase inhibitors (TKIs) have transformed the treatment landscape of chronic-phase chronic myeloid leukemia, normalizing life expectancy for most patients.
- • **Risultato chiave:** For frontline therapy, generically available first-generation BCR::ABL1 TKI imatinib is a rational and cost-effective approach to optimize survival.
- • **Rilevanza clinica:** Considerable finesse is warranted to tailor a range of efficacious and well-tolerated therapies to the individual needs of patients.

3) Perioperative care of patients with chronic myeloid leukemia on tyrosine kinase inhibitors.

Review *Frontiers in oncology* Score: ***10***

***Focus clinico**: Safety e comorbidità, Sintesi (review/linee guida)*

PMID: 42857533 | 10.3389/fonc.2026.1899207

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42857533/`

- • **Tipo di studio/popolazione:** Patients with chronic myeloid leukemia (CML) are increasingly undergoing surgical procedures as long-term survival improves with tyrosine kinase inhibitor (TKI) therapy.
- • **Risultato chiave:** This narrative review aims to synthesize current evidence regarding perioperative risk assessment and management in patients with CML receiving TKIs.
- • **Rilevanza clinica:** Perioperative management in CML requires individualized multidisciplinary planning integrating hematologic, anesthetic, surgical, and pharmacologic considerations to optimize surgical safety while maintaining disease control.

4) Rethinking cardiovascular risk stratification in TKI-treated chronic myeloid leukemia.

Altro *Cancer treatment and research communications* Score: ***9***

***Focus clinico**: Safety e comorbidità*

PMID: 42833183 | 10.1016/j.ctarc.2026.101473

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42833183/`

- • **Tipo di studio/popolazione:** Current cardiovascular management relies on general population risk scores and ankle-brachial index (ABI) measurement, tools not designed to capture TKI-specific vascular biology or detect subclinical arterial damage.
- • **Risultato chiave:** We appraise cardiovascular risk stratification tools and vascular imaging modalities in the context of TKI-treated CML, and propose an integrated algorithm combining dual risk scoring, ABI, and targeted femoral duplex ultrasound (DUS).
- • **Rilevanza clinica:** Vascular outcome reporting in TKI trials is hampered by terminological inconsistency: Major Adverse Limb Events (MALE), the standard limb-specific endpoint in contemporary cardiovascular trials, has never been adopted in any TKI/CML publication.

5) Tyrosine kinase inhibitor use and risk of neurodegenerative disease: A Swedish population-based cohort study.

Real-world / osservazionale *Alzheimer's & dementia : the journal of the Alzheimer's Association* Score: ***6***

***Focus clinico**: Altro*

PMID: 42860391 | 10.1002/alz.71916

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42860391/`

- • **Tipo di studio/popolazione:** CML patients ≥50 years at diagnosis initiating TKI therapy were identified, alongside five comparators without CML matched on sex, age, and residence at diagnosis.
- • **Risultato chiave:** A total of 1322 CML cases were included in the study.
- • **Rilevanza clinica:** 1188 initiated imatinib (of which 41 experienced an NDD event), 541 dasatinib (15 events), 390 nilotinib (five events), 174 bosutinib, 71 ponatinib, and 41 asciminib.

Evidenze per domande cliniche

Resistenza / T315I / mutazioni (3)

Score: ***0***

Mini-riassunto (da abstract, estrattivo):

The evolving landscape of tyrosine kinase inhibitors in chronic-phase chronic myeloid leukemia: A 25-year journey.

Altro *Blood reviews* *Score:* ***11***

PMID: 42875330 | 10.1016/j.blre.2026.101436

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42875330/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** BCR::ABL1 tyrosine kinase inhibitors (TKIs) have transformed the treatment landscape of chronic-phase chronic myeloid leukemia, normalizing life expectancy for most patients.
- • **Risultato chiave:** For frontline therapy, generically available first-generation BCR::ABL1 TKI imatinib is a rational and cost-effective approach to optimize survival.
- • **Rilevanza clinica:** Considerable finesse is warranted to tailor a range of efficacious and well-tolerated therapies to the individual needs of patients.

Mini-riassunto (da abstract, estrattivo):

A novel BCR::ABL1 I502T mutation associated with acquired asciminib resistance in CML.

Altro *Leukemia research* *Score:* ***4***

PMID: 42837802 | 10.1016/j.leukres.2026.108335

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42837802/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** non riportato (abstract non disponibile).
- • **Risultato chiave:** non riportato.
- • **Rilevanza clinica:** non valutabile senza abstract.

Mini-riassunto (da abstract, estrattivo):

Significance of BCR-ABL P-loop point mutations (L248R/V) on binding to FDA approved drugs: molecular dynamics simulations and DFT studies.

Altro *Journal of biomolecular structure & dynamics* Score: ***4***

PMID: 42834848 | 10.1080/07391102.2026.2735399

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42834848/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** The BCR-ABL fusion protein with dysregulated tyrosine kinase activity is the cause of chronic myelogenous leukemia (CML).
- • **Risultato chiave:** This work aims to study the structural, dynamical, and functional effects of the P-loop point mutations on BCR-ABL kinase.
- • **Rilevanza clinica:** The MD simulations and QM studies suggest that L248 residue and its point mutations (L248R/V) are responsible for high affinity binding of BCR-ABL kinase with the drugs considered for the study.

TFR / stop terapia (2)

Score: ***0***

Mini-riassunto (da abstract, estrattivo):

Successful Pregnancy After Planned Pregnancy-Oriented TKI Interruption and Embryo Transfer Following Ponatinib Re-Consolidation in Chronic-Phase Chronic Myeloid Leukemia.

Trial clinico *EJHaem* Score: ***18***

PMID: 42840123 | 10.1002/jha2.70426

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42840123/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** Pregnancy after unsuccessful treatment-free remission (TFR) in chronic-phase chronic myeloid leukemia (CP-CML) remains challenging.
- • **Risultato chiave:** Ponatinib re-consolidation enabled a second TKI-free interval, with MMR maintained but deep molecular response unstable.
- • **Rilevanza clinica:** Further ponatinib re-consolidation restored MR4.5; ponatinib was stopped before embryo transfer, resulting in pregnancy and live birth without TKI exposure.

Mini-riassunto (da abstract, estrattivo):

The evolving landscape of tyrosine kinase inhibitors in chronic-phase chronic myeloid leukemia: A 25-year journey.

Altro *Blood reviews* *Score: **11***

PMID: 42875330 | 10.1016/j.blre.2026.101436

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42875330/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** BCR::ABL1 tyrosine kinase inhibitors (TKIs) have transformed the treatment landscape of chronic-phase chronic myeloid leukemia, normalizing life expectancy for most patients.
- • **Risultato chiave:** For frontline therapy, generically available first-generation BCR::ABL1 TKI imatinib is a rational and cost-effective approach to optimize survival.
- • **Rilevanza clinica:** Considerable finesse is warranted to tailor a range of efficacious and well-tolerated therapies to the individual needs of patients.

Safety e comorbidità (2)

*Score: **0***

Mini-riassunto (da abstract, estrattivo):

Perioperative care of patients with chronic myeloid leukemia on tyrosine kinase inhibitors.

Review *Frontiers in oncology* *Score: **10***

PMID: 42857533 | 10.3389/fonc.2026.1899207

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42857533/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** Patients with chronic myeloid leukemia (CML) are increasingly undergoing surgical procedures as long-term survival improves with tyrosine kinase inhibitor (TKI) therapy.
- • **Risultato chiave:** This narrative review aims to synthesize current evidence regarding perioperative risk assessment and management in patients with CML receiving TKIs.
- • **Rilevanza clinica:** Perioperative management in CML requires individualized multidisciplinary planning integrating hematologic, anesthetic, surgical, and pharmacologic considerations to optimize surgical safety while maintaining disease control.

Mini-riassunto (da abstract, estrattivo):

Rethinking cardiovascular risk stratification in TKI-treated chronic myeloid leukemia.

Altro *Cancer treatment and research communications* *Score: **9***

PMID: 42833183 | 10.1016/j.ctarc.2026.101473

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42833183/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** Current cardiovascular management relies on general population risk scores and ankle-brachial index (ABI) measurement, tools not designed to capture TKI-specific vascular biology or detect subclinical arterial damage.
- • **Risultato chiave:** We appraise cardiovascular risk stratification tools and vascular imaging modalities in the context of TKI-treated CML, and propose an integrated algorithm combining dual risk scoring, ABI, and targeted femoral duplex ultrasound (DUS).
- • **Rilevanza clinica:** Vascular outcome reporting in TKI trials is hampered by terminological inconsistency: Major Adverse Limb Events (MALE), the standard limb-specific endpoint in contemporary cardiovascular trials, has never been adopted in any TKI/CML publication.

Sequencing / linee di terapia (1)

*Score: **0***

Mini-riassunto (da abstract, estrattivo):

The evolving landscape of tyrosine kinase inhibitors in chronic-phase chronic myeloid leukemia: A 25-year journey.

Altro *Blood reviews* *Score: **11***

PMID: 42875330 | 10.1016/j.blre.2026.101436

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42875330/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** BCR::ABL1 tyrosine kinase inhibitors (TKIs) have transformed the treatment landscape of chronic-phase chronic myeloid leukemia, normalizing life expectancy for most patients.
- • **Risultato chiave:** For frontline therapy, generically available first-generation BCR::ABL1 TKI imatinib is a rational and cost-effective approach to optimize survival.
- • **Rilevanza clinica:** Considerable finesse is warranted to tailor a range of efficacious and well-tolerated therapies to the individual needs of patients.

Altro (1)

*Score: **0***

Mini-riassunto (da abstract, estrattivo):

Tyrosine kinase inhibitor use and risk of neurodegenerative disease: A Swedish population-based cohort study.

Real-world / osservazionale *Alzheimer's & dementia : the journal of the Alzheimer's Association* Score: **6**

PMID: 42860391 | 10.1002/alz.71916

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42860391/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** CML patients ≥50 years at diagnosis initiating TKI therapy were identified, alongside five comparators without CML matched on sex, age, and residence at diagnosis.
- • **Risultato chiave:** A total of 1322 CML cases were included in the study.
- • **Rilevanza clinica:** 1188 initiated imatinib (of which 41 experienced an NDD event), 541 dasatinib (15 events), 390 nilotinib (five events), 174 bosutinib, 71 ponatinib, and 41 asciminib.

Sintesi (review/linee guida) (1)

Score: **0**

Mini-riassunto (da abstract, estrattivo):

Perioperative care of patients with chronic myeloid leukemia on tyrosine kinase inhibitors.

Review *Frontiers in oncology* Score: **10**

PMID: 42857533 | 10.3389/fonc.2026.1899207

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42857533/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** Patients with chronic myeloid leukemia (CML) are increasingly undergoing surgical procedures as long-term survival improves with tyrosine kinase inhibitor (TKI) therapy.
- • **Risultato chiave:** This narrative review aims to synthesize current evidence regarding perioperative risk assessment and management in patients with CML receiving TKIs.
- • **Rilevanza clinica:** Perioperative management in CML requires individualized multidisciplinary planning integrating hematologic, anesthetic, surgical, and pharmacologic considerations to optimize surgical safety while maintaining disease control.

Dose strategy / consolidamento (1)

Score: **0**

Mini-riassunto (da abstract, estrattivo):

Successful Pregnancy After Planned Pregnancy-Oriented TKI Interruption and Embryo Transfer Following Ponatinib Re-Consolidation in Chronic-Phase Chronic Myeloid Leukemia.

Trial clinico *EJHaem* *Score:* ***18***

PMID: 42840123 | 10.1002/jha2.70426

`Link PubMed: https://pubmed.ncbi.nlm.nih.gov/42840123/`

Open Access (Europe PMC/PMC): non rilevato

- • **Tipo di studio/popolazione:** Pregnancy after unsuccessful treatment-free remission (TFR) in chronic-phase chronic myeloid leukemia (CP-CML) remains challenging.
- • **Risultato chiave:** Ponatinib re-consolidation enabled a second TKI-free interval, with MMR maintained but deep molecular response unstable.
- • **Rilevanza clinica:** Further ponatinib re-consolidation restored MR4.5; ponatinib was stopped before embryo transfer, resulting in pregnancy and live birth without TKI exposure.