

IE 521 Convex Optimization

Lecture 11: Center of Gravity, Ellipsoid Method

Niao He

7th April 2019

Outline

Complexity vs
Convergence

Complexity vs Convergence

Cutting Plane
Methods

Cutting Plane Methods

Center of Gravity
Method

Center of Gravity Method

Ellipsoid Method

Ellipsoid Method

Complexity

Given an input $\epsilon > 0$, a problem instance P ,

- ▶ *Oracle complexity*: number of oracles required to solve the problem (P) up to accuracy $\epsilon > 0$
- ▶ *Arithmetic complexity*: number of arithmetic operation (bit-wise operation) requirement to solve the problem (P) up to accuracy $\epsilon > 0$

Convergence

Given solutions $\{x_t\}$ and accuracy measure $\mathcal{E}(x_t)$

$$\lim_{t \rightarrow \infty} \frac{\mathcal{E}(x_{t+1})}{\mathcal{E}(x_t)^p} = q$$

- ▶ **Linear convergence:** $p = 1, q \in (0, 1)$
 - ▶ E.g., $\mathcal{E}(x_t) = O(e^{-\alpha t})$, where $\alpha > 0$
- ▶ **Sublinear convergence:** $p = 1, q = 1$
 - ▶ E.g., $\mathcal{E}(x_t) = \frac{1}{t^\beta}$, where $\beta > 0$
- ▶ **Superlinear convergence:** $p = 1, q = 0$
 - ▶ E.g., $\mathcal{E}(x_t) = O(e^{-\alpha t^2})$, where $\alpha > 0$
- ▶ **Convergence of order p :** $p > 1, q > 0$
 - ▶ When $p = 2$, called quadratic convergence.
 - ▶ E.g., $\mathcal{E}(x_t) = O(e^{-\alpha p^t})$, where $\alpha > 0$

Illustration: Convergence




Figure: sublinear, linear, quadratic convergence

Solving Convex Program

We focus on the following general convex problem


$$\min_{x \in X} f(x)$$

Problem Setting:


- ▶ f is convex and admits *zero- and first-order oracles*;
- ▶ $X \subset \mathbb{R}^n$ is a convex body (convex, compact, with nonempty interior) and admits *separation oracle*.

Cutting Plane Methods

$$\min_{x \in X} f(x)$$



(a) Separation oracle



(b) First-order oracle

- (a) $X \subseteq \{y : a^T(y - x) \leq 0\}$ if $x \notin X$;
- (b) $X^* \subseteq \{y : g^T(y - x) \leq 0\}$ if x is not optimal.

Cutting Plane Methods




Figure: Localization Polyhedron

$$\mathcal{P}_1 \supseteq \cdots \supseteq \mathcal{P}_k \supseteq X^*$$

Q. How to choose the query point to cut the most off?

Cutting Plane Methods

- ▶ *Center of gravity method*: choose the query to be the center of the gravity of \mathcal{P}_k .
- ▶ *Maximum volume ellipsoid cutting plane method*: choose the query to be the center of the maximum volume ellipsoid contained in \mathcal{P}_k .
- ▶ *Chebyshev center cutting-plane method*: choose the query point to be the Chebyshev center of \mathcal{P}_k , i.e., the center of the largest Euclidean ball that lies in \mathcal{P}_k .

Center of Gravity Method

(Levin, 1965; Newman, 1965)

- ▶ Initialize $G_0 = X$
 - ▶ At iteration $t = 1, 2, \dots, T$, do
 - ▶ Compute the center of gravity:

$$x_t = \frac{1}{\text{Vol}(G_{t-1})} \int_{x \in G_{t-1}} x dx$$
 - ▶ Call the first order oracle and obtain $g_t \in \partial f(x_t)$
 - ▶ Set $G_t = G_{t-1} \cap \{y : g_t^T(y - x_t) \leq 0\}$
 - ▶ Output $\hat{x}_T \in \arg \min_{x \in \{x_1, \dots, x_T\}} f(x)$
-

Center of Gravity Method

Lemma (Grünbaum [1960]). Let $C \subset \mathbb{R}^n$ be a convex body with $\int_C x dx = 0$. Then $\forall a \neq 0$

$$\begin{aligned}\text{Vol}(C \cap \{x : a^T x \leq 0\}) &\leq \left(1 - \left(\frac{n}{n+1}\right)^n\right) \text{Vol}(C) \\ &\leq \left(1 - \frac{1}{e}\right) \text{Vol}(C) \approx 0.63 \text{Vol}(C)\end{aligned}$$

Remark. It follows that

$$\text{Vol}(G_t) \leq \left(1 - \frac{1}{e}\right)^t \text{Vol}(X), t \geq 1$$

Convergence of Center of Gravity Method

Theorem. The center of gravity method return $\hat{x}_T \in X$:

$$f(\hat{x}_T) - f^* \leq \left(1 - \frac{1}{e}\right)^{\frac{T}{n}} \cdot \text{Var}_X(f)$$

where $\text{Var}_X(f) = \max_{x \in X} f(x) - \min_{x \in X} f(x)$.

- ▶ Linear convergence rate
- ▶ Oracle complexity: $N(\epsilon) = \mathcal{O}\left(n \log\left(\frac{\text{Var}_X(f)}{\epsilon}\right)\right)$
- ▶ Main disadvantage: computing the center of gravity is extremely difficult, even for polytopes.

Proof of Convergence

- ▶ Note $x^* \in G_t, \forall t \geq 1$ and $\text{Vol}(G_t) \leq (1 - \frac{1}{e})^t \text{Vol}(X)$.
- ▶ Consider the neighborhood of x^* :
 $X_\delta = \{x^* + \delta(x - x^*) : x \in X\}$, where $\delta \in ((1 - \frac{1}{e})^{\frac{T}{n}}, 1)$.
- ▶ Observe that $X_\delta / G_T \neq 0$.

$$\text{Vol}(X_\delta) = \delta^n \text{Vol}(X) > (1 - \frac{1}{e})^T \text{Vol}(X) \geq \text{Vol}(G_T)$$

- ▶ Let $y = x^* + \delta(z - x^*) \in X_\delta / G_T$ for some $z \in X$.
Thus, for certain $t^* \leq T$, we have $y \in G_{t^*-1} / G_{t^*}$.
- ▶ Since $y \notin G_{t^*}$, we have $g_{t^*}^T(y - x_{t^*}) > 0$, so $f(y) > f(x_{t^*})$.
- ▶ Since $y = x^* + \delta(z - x^*)$, by convexity of f ,

$$\begin{aligned} f(y) &= f(\delta z + (1 - \delta)x^*) \leq \delta f(z) + (1 - \delta)f(x^*) \\ &= f(x^*) + \delta[f(z) - f(x^*)] \\ &\leq f(x^*) + \delta \text{Var}_X(f) \end{aligned}$$

Hence $f(\hat{x}_T) \leq f(x_{t^*}) \leq f^* + \delta \text{Var}_X(f)$.

Ellipsoid as Localizer


Definition. Let $Q \succ 0$ be symmetric, and c be the center, an ellipsoid is uniquely characterized by (c, Q) :

$$\begin{aligned} E(c, Q) &= \left\{ x \in \mathbb{R}^n : (x - c)^T Q^{-1} (x - c) \leq 1 \right\} \\ &= \left\{ x = c + Q^{\frac{1}{2}} u : u^T u \leq 1 \right\} \end{aligned}$$

- $\text{Vol}(E(c, Q)) = \text{Det}(Q^{\frac{1}{2}}) \text{Vol}(B_n)$, where B_n is a unit Euclidean ball in \mathbb{R}^n with $\text{Vol}(B_n) = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2}+1)}$.

Half Ellipsoid

Let $H_+ = \{x : \omega^T x \leq \omega^T c\}$ be a half space with $\omega \neq 0$ that pass through the center c of the ellipsoid $E(c, Q)$.



Half Ellipsoid

Let $H_+ = \{x : \omega^T x \leq \omega^T c\}$ be a half space with $\omega \neq 0$ that pass through the center c of the ellipsoid $E(c, Q)$.

- ▶ $E \cap H_+ \subseteq E^+ = E(c^+, Q^+)$ with

$$c^+ = c - \frac{1}{n+1}q, \text{ where } q = \frac{Q\omega}{\sqrt{\omega^T Q \omega}},$$

$$Q^+ = \frac{n^2}{n^2+1} \left(Q - \frac{2}{n+1} q q^T \right).$$

- ▶ Volume decrease:

$$\text{Vol}(E^+) \leq \exp \left\{ -\frac{1}{2n} \right\} \text{Vol}(E)$$

Ellipsoid Method

(Shor; Nemirovsky, Yudin, 1970s)

- ▶ Initialize $E(c_0, Q_0)$ with $c_0 = 0, Q_0 = R^2 I$
- ▶ At iteration $t = 1, 2, \dots, T$, do
 - ▶ Call separation oracle with the input c_{t-1}
 - ▶ If $c_{t-1} \notin X$, call separation oracle and obtain $\omega \neq 0$
 - ▶ If $c_{t-1} \in X$, call first order oracle and obtain $\omega \in \partial f(c_t)$
 - ▶ Set the new ellipsoid $E(c_t, Q_t)$ with

$$c_t = c_{t-1} - \frac{1}{n+1} \frac{Q_{t-1}\omega}{\sqrt{\omega^T Q_{t-1}\omega}}$$
$$Q_t = \frac{n^2}{n^2 - 1} \left(Q_{t-1} - \frac{2}{n+1} \frac{Q_{t-1}\omega\omega^T Q_{t-1}}{\omega^T Q_{t-1}\omega} \right)$$

- ▶ Output $\hat{x}_T = \arg \min_{c \in \{c_1, \dots, c_T\} \cap X} f(c)$
-

Illustration of Ellipsoid Method




Figure: Illustration

Figure from Boyd, EE364b lecture notes

Convergence of Ellipsoid Method

Theorem. Assume $B(\bar{x}, r^2 I) \subseteq X \subseteq B(0, R^2 I)$. The Ellipsoid method after T steps satisfies:

$$f(\hat{x}_T) - f^* \leq \frac{R}{r} \cdot \text{Var}_X(f) \exp \left\{ -\frac{T}{2n^2} \right\}$$

- ▶ Linear convergence rate
- ▶ Oracle complexity: $N(\epsilon) = \mathcal{O} \left(n^2 \log \left(\frac{\text{Var}_X(f)}{\epsilon} \right) \right)$
- ▶ Modest per iteration computation cost: $\mathcal{O}(n^2)$
- ▶ Polynomial solvability: as long as it takes polynomial time to call the separation and first-order oracles

Proof of Convergence

- ▶ Similar as the proof for the center of gravity method.
- ▶ Consider the neighborhood of x^* :
 $X_\delta = \{x^* + \delta(x - x^*) : x \in X\}$, $\delta \in (\frac{R}{r} \exp\{-\frac{T}{2n^2}\}, 1)$.
- ▶ Note $X_\delta/E(c_T, Q_T) \neq \emptyset$, because

$$\begin{aligned}\text{Vol}(X_\delta) &= \delta^n \text{Vol}(X) \\ &\geq \delta^n r^n \text{Vol}(B_n) \\ &> R^n \exp\left\{-\frac{T}{2n}\right\} \text{Vol}(B_n) \\ &\geq \text{Vol}(E(c_T, Q_T))\end{aligned}$$

- ▶ Rest is the same as proof of center of gravity method.

Stopping Criterion

- ▶ In practice, f^* is often unknown and it is impossible to compute $f(x_t) - f^*$.
- ▶ Construct online lower bounds for f^* : $\ell_t \leq f^*$

$$\begin{aligned} f^* &\geq f(x_t) + \omega_t^T (x^* - x_t), \quad \omega_t \in \partial f(x_t) \\ &\geq f(x_t) + \inf_{x \in E(c_t, Q_t)} \omega_t^T (x - x_t) \\ &= f(x_t) - \sqrt{\omega_t^T Q_t \omega_t} \end{aligned}$$

- ▶ Hence, $\sqrt{\omega_t^T Q_t \omega_t} \leq \epsilon \implies f(x_t) - f^* \leq \epsilon$
- ▶ Tighter lower bound:

$$\ell_t = \max_{1 \leq \tau \leq t} (f(x_\tau) - \sqrt{\omega_\tau^T Q_\tau \omega_\tau})$$

Discussion

Advantages and disadvantages of the Ellipsoid method:

- + : universal
- + : simple to implement
- + : steady for small size problems
- + : low order dependence on the number of constraints
- : quadratic growth on the size of problem
- : inefficient for large-scale problems

Experiment on SVM

$$\min_{w,b} \frac{1}{m} \sum_{i=1}^m \max(1 - y_i(w^T x_i + b), 0) + \lambda \|w\|_2^2$$




Figure: Ellipsoid Method for SVM on WBDC dataset (n=30)

References

Complexity vs
Convergence

Cutting Plane
Methods

Center of Gravity
Method

Ellipsoid Method

- Ben-Tal & Nemirovski, Chapter 7